Genetic variant rs3750625 in the 3'UTR of *ADRA2A* affects stress-dependent acute pain severity after trauma and alters a microRNA-34a regulatory site

Sarah D. Linnstaedt^{a,b,*}, Margaret G. Walker^{a,b}, Kyle D. Riker^{a,b}, Jennifer E. Nyland^c, JunMei Hu^{a,b}, Catherine Rossi^d, Robert A. Swor^e, Jeffrey S. Jones^f, Luda Diatchenko^g, Andrey V. Bortsov^{a,b}, David A. Peak^h, Samuel A. McLean^{a,b,i}

Abstract

α 2A adrenergic receptor (α 2A-AR) activation has been shown in animal models to play an important role in regulating the balance of acute pain inhibition vs facilitation after both physical and psychological stress. To our knowledge, the influence of genetic variants in the gene encoding α 2A-AR, *ADRA2A*, on acute pain outcomes in humans experiencing traumatic stress has not been assessed. In this study, we tested whether a genetic variant in the 3'UTR of *ADRA2A*, rs3750625, is associated with acute musculoskeletal pain (MSP) severity following motor vehicle collision (MVC, $n = 948$) and sexual assault ($n = 84$), and whether this influence was affected by stress severity. We evaluated rs3750625 because it is located in the seed binding region of miR-34a, a microRNA (miRNA) known to regulate pain and stress responses. In both cohorts, the minor allele at rs3750625 was associated with increased musculoskeletal pain in distressed individuals (stress*rs3750625 $P = 0.043$ for MVC cohort and $P = 0.007$ for sexual assault cohort). We further found that (1) miR-34a binds the 3'UTR of *ADRA2A*, (2) the amount of repression is greater when the minor (risk) allele is present, (3) miR-34a in the IMR-32 adrenergic neuroblastoma cell line affects *ADRA2A* expression, (4) miR-34a and *ADRA2A* are expressed in tissues known to play a role in pain and stress, (5) following forced swim stress exposure, rat peripheral nerve tissue expression changes are consistent with miR-34a regulation of *ADRA2A*. Together, these results suggest that *ADRA2A* rs3750625 contributes to poststress musculoskeletal pain severity by modulating miR-34a regulation.

Keywords: Motor vehicle collision, Sexual assault, Stress, Trauma, Musculoskeletal pain, Polymorphism, Single nucleotide, microRNA, miR-34a, Adrenergic receptor

1. Introduction

Exposure to traumatic stress is, unfortunately, common in life.⁴² For example, in America, each year more than eleven million individuals experience a motor vehicle collision (MVC) and nearly 700,000 women are sexually assaulted.^{1,49} The precise molecular mechanisms influencing acute musculoskeletal pain (MSP)

severity after stressful events such as MVC and sexual assault remain poorly understood.

α 2A adrenergic receptor (α 2A-AR) activation has been shown to influence pain outcomes in both preclinical and human studies.^{1–10} These data include animal model data demonstrating that α 2A-AR activation plays an important role in regulating the balance of acute pain inhibition vs facilitation after both physical and psychological stress.¹⁹ However, to our knowledge, the influence of α 2A-AR activation on acute MSP outcomes in humans experiencing traumatic stress has not been assessed. One way to assess the potential influence of α 2A-AR activation on acute MSP in trauma-exposed individuals is to evaluate for associations between genetic variants influencing α 2A-AR function and acute MSP outcomes. If α 2A-AR activation influences acute MSP severity in humans experiencing traumatic stress, then inherited differences in the gene for α 2A-AR, *ADRA2A*, should be associated with individual differences in acute MSP severity. Few studies have examined associations between *ADRA2A* polymorphisms and responses to stress generally or pain outcomes specifically. One 2004 study of healthy volunteers found that a polymorphism in the 3'UTR (3'untranslated region) of *ADRA2A* modulates autonomic responses to physiological and environmental stress,²³ and another study of healthy volunteers found that *ADRA2A* polymorphisms predict evoked pain responses.³² To our knowledge, no prospective cohort studies have evaluated associations between *ADRA2A* genetic

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

Scientific Meeting Presentation: American Pain Society Meeting 2014, Tampa, FL.

^a TRYUMPH Research Program, ^b Department of Anesthesiology, University of North Carolina, Chapel Hill, NC, USA, ^c Department of Battlefield Pain Research, Fort Sam, TX, USA, ^d Forensic Nursing Program, Family Medicine, Cone Health System, Greensboro, NC, USA, ^e Department of Emergency Medicine, William Beaumont Hospital, Royal Oak, MI, USA, ^f Department of Emergency Medicine, Spectrum Health Butterworth Campus, Grand Rapids, MI, USA, ^g The Allan Edwards Centre for Research on Pain, McGill University, Montreal, QC, Canada, ^h Department of Emergency Medicine, Massachusetts General Hospital, Boston, MA, USA, ⁱ Department of Emergency Medicine, University of North Carolina, Chapel Hill, NC, USA

*Corresponding author. Address: University of North Carolina, Medical School Wing C, CB#7010, Chapel Hill, NC 27599-7010, USA. Tel.: 919-576-6179; fax: 919-966-7193. E-mail address: slinnstaedt@aims.unc.edu (S. D. Linnstaedt).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's Web site (www.painjournalonline.com).

PAIN 158 (2017) 230–239

© 2016 International Association for the Study of Pain

<http://dx.doi.org/10.1097/j.pain.0000000000000742>

variants and clinical acute MSP outcomes following trauma/stress exposure.

In this study, we assessed the association between a single-nucleotide polymorphism (SNP) in the 3'UTR of *ADRA2A*, rs3750625, and acute MSP outcomes among individuals experiencing MVC and sexual assault. We evaluated this SNP because bioinformatics analyses indicate that it is in the seed binding region of miR-34a, a miRNA known to affect pain and stress responses.^{27,60} Increasing evidence indicates that one common mechanism by which genetic polymorphisms influence cellular function is by altering miRNA seed binding regions, thus altering the ability of miRNA to bind and regulate mRNA and the amount of protein product produced.² When assessing the association between rs3750625 and acute MSP outcomes, we included potential gene variant $\times$ sex and gene variant $\times$ stress interactions because genetic influences on MSP after stress exposure are often sex and/or stress severity dependent (eg,^{8,39,58}). We then experimentally tested whether the allele present at rs3750625 affects miR-34a binding to the 3'UTR of *ADRA2A*, and whether miR-34a and *ADRA2A* expression levels change in key tissues involved in stress-induced hyperalgesia in an animal model of stress exposure.

2. Methods

2.1. Motor vehicle collision cohort

2.1.1. Study design and population

The methods of the MVC study have been described.⁴⁵ In brief, individuals ≥ 18 and ≤ 65 years of age presenting to 1 of 8 emergency departments (EDs) in 4 no-fault insurance states within 24 hours of MVC who did not have fracture or other injury requiring hospital admission were enrolled. Patients who were not alert and oriented were excluded, as were pregnant patients, prisoners, patients unable to read and understand English, patients taking a β -adrenoreceptor antagonist, or patients taking opioids above a total daily dose of 30 mg of oral morphine or equivalent. In addition, enrollment was limited to non-Hispanic whites (the most common ethnicity at study sites). Informed consent was obtained from all participants and Institutional Review Board (IRB) approval was obtained at all study sites.

2.1.2. DNA collection and genotyping

Study personnel collected blood samples at the time of enrollment using PAXgene DNA tubes. Following DNA purification (PAXgene blood DNA kit, QIAGEN), genotyping using the Sequenom platform was performed at rs3750625. As described above, this SNP was chosen for analysis based on its location in a regulatory region of the pain and stress-associated gene, *ADRA2A*, and because bioinformatics analyses suggest that this allele might affect miR-34a binding (^{12,37,50,62}). Two HapMap samples and 2 repeat samples were included in each genotyping batch of 96 samples to ensure genotyping accuracy and reliability. rs3750625 was in Hardy–Weinberg equilibrium ($P > 0.05$), and repeat genotyping demonstrated greater than 98% call agreement.

2.1.3. Assessments

Acute MSP severity was assessed in the ED using a verbal 0 to 10 numeric rating scale (NRS).⁹ Distress in the ED was measured using the peritraumatic distress inventory.¹¹ A validated cutoff of 23 was used to identify those with substantial distress.⁴¹

2.2. Sexual assault cohort

2.2.1. Study design and population

The methods of the sexual assault study have been reported.^{4,38} In brief, women 18 years of age or older who presented to 1 of 10 sexual assault nurse examiner (SANE) programs in 4 states for medical care within 72 hours of sexual assault were recruited. Women unable to give informed consent (eg, due to intoxication) were excluded, as were women who were hospitalized after sexual assault, lived with their assailant, were prisoners, were pregnant, did not have a telephone, and/or did not live within driving distance for follow-up interviews. Institutional Review Board (IRB) approval was obtained at all study sites and all study participants provided written informed consent.

2.2.2. DNA collection and genotyping

Saliva specimens were obtained at 1-week follow-up evaluations using Oragene DNA self-collection kits. Following DNA purification, genotyping was performed within the same batches as MVC cohort samples. HapMap samples and 2 repeat samples were included in each genotyping batch to ensure genotypic accuracy and reliability; repeated genotyping demonstrated greater than 98% call agreement, and rs3750625 was in Hardy–Weinberg equilibrium ($P > 0.05$).

2.2.3. Assessments

Acute MSP severity was assessed in the ED using a verbal 0 to 10 numeric rating scale. One week after the assault, sexual assault survivors completed an in-person computerized self-report questionnaire that included an assessment of the individual's distress levels at the time of assault using the acute stress disorder (ASD) questionnaire.¹³ The ASD is a 19-item questionnaire assessing feelings that were experienced during or immediately after the assault. Each item on the questionnaire was evaluated via numeric rating scale from 0 (no distress) to 5 (high distress). A cutoff of 56 for substantial acute stress (ASD scale) has been reported previously.¹³ However, over 90% of the participants in the sexual assault cohort reported distress levels above this threshold; thus, we used the median score of 77 to distinguish more vs less distressed sexual assault survivors.

2.2.4. Statistical analyses

Sociodemographic characteristics of the sample were summarized using standard descriptive statistics. General linear models were used to evaluate the association between rs3750625 with acute MSP outcomes and for potential sex $\times$ rs3750625 and stress $\times$ rs3750625 interactions, adjusting for potential confounding by age and study site. Sex and stress-dependent effects were evaluated because of increasing evidence that such interactions are frequently present^{8,39,58} and because such effects have been found in *ADRA2A* previously.^{19,23} A dominant genetic model (homozygous for major allele vs other) was used in all analyses due to the low minor allele frequency (MAF = 0.05) for rs3750625 in European Americans. Mean levels of MSP in distressed and nondistressed individuals homozygous for the major allele (CC) or individuals heterozygous or homozygous for the minor allele (CA/AA) were obtained from general linear models. All statistical analyses were completed using SPSS software (v.23; SPSS Inc, Chicago, IL).

2.2.5. Bioinformatics analyses

The miRdSNP online algorithm (<http://mirdsnp.ccr.buffalo.edu/search.php>) was used to identify whether rs3750625 was predicted to affect microRNA (miRNA) binding.¹² The UCSC genome browser (<https://genome.ucsc.edu/>) was used to determine species conservation.²⁸ RNA hybrid (<http://bibiserv.techfak.uni-bielefeld.de/rnahybrid/>) was used to predict the secondary structure of the miR-34a-ADRA2A binding event.³³

2.2.6. Constructs

Lentiviral constructs containing either a firefly luciferase gene (pL-SV40-GL3) or a renilla luciferase gene (pL-SV40-RLUC) were used for dual luciferase assays.²⁶ The 3'UTR of ADRA2A was amplified from human genomic DNA using primers 1-F and 1-R (Supplementary Table 1, available online at <http://links.lww.com/PAIN/A354>). The resulting 1360bp product was cloned downstream of the firefly luciferase gene in pL-SV40-GL3 using XhoI and EcoRI restriction enzyme sites. This newly created construct, pL-GL3-ADRA2A-maj, was sequenced to confirm the major allele, C, at position rs3750625 in the ADRA2A 3'UTR and was then used as template to mutate the major allele to the minor allele, A, using the QuickChange II Site-Directed Mutagenesis Kit (Promega) and primers 2-F and 2-R (Supplementary Table 1, available online at <http://links.lww.com/PAIN/A354>). The construct containing the minor allele was named pL-GL3-ADRA2A-min.

Seed region and compensatory region mutations were made in the ADRA2A 3'UTR using 2-step PCR. pL-GL3-ADRA2A-maj was used as the parent construct, and primers containing mutant sequence at either the seed region or compensatory site were used to create constructs pL-GL3-ADRA2A-seed (Primers 1-F, 1-R, 7-F, and 7-R, Supplementary Table 1, available online at <http://links.lww.com/PAIN/A354>) and pL-GL3-ADRA2A-comp (Primers 1-F, 1-R, 6-F, 6-R, Supplementary Table 1, available online at <http://links.lww.com/PAIN/A354>), respectively. The pcDNA-based miR-34a expression construct (pC-34a) was generously donated by Moshe Oren (The Weizmann Institute of Science).

pLCE-s34 and associated controls, pLCE and pLCE sCXCR4 were generously donated from the Luftig Lab (Duke University). Generation of these constructs and the efficacy with which pLCE-s34 knocks down miR-34a expression have been previously described.²⁴

2.2.7. Cell culture and generation of neuroblastoma cells stably knocked down for miR-34a expression

HEK293T cells were grown in Dulbecco's Modified Eagle's Medium (DMEM) with 10% fetal bovine serum (FBS) and 1% gentamicin. IMR-32 cells are human adrenergic neuroblastoma cells with high levels of the catecholamine precursor, tyrosine hydroxylase.⁶¹ They were cultured in Eagle's Minimum Essential Medium (EMEM) with 10% FBS and 1% gentamicin. All cells were incubated at 37°C, 5% CO₂.

To address whether miR-34a plays a significant role in regulating endogenous levels of ADRA2A, we suppressed miR-34a activity in the IMR-32 neuroblastoma cell line. miR-34a activity was inhibited by transduction of these cells with a GFP-based lentiviral miRNA sponge specific for miR-34a (s34). miRNA sponges represent a well-established technique^{21,36,59} for specifically and stably blocking the activity of an miRNA through overexpression of an mRNA containing the GFP indicator gene

linked to multiple (in this case, 4) copies of an incompletely complementary artificial target site for the miRNA of interest. This particular miR-34a sponge has previously been shown to result in robust knockdown of miR-34a.²⁴

We also separately transduced a previously described²⁶ control lentiviral miRNA sponge vector (sCXCR4) that expresses a sponge specific for a small interfering RNA that inhibits CXCR4 mRNA expression. The second control vector represents the parental lentiviral vector expressing only GFP (pLCE). Viral transductants for pLCE-s34 were produced in HEK293T cells using a third-generation lentiviral packaging system²⁰ consisting of pMD2-VSVG, pRSV-REV, pMDLgp, and either pLCE-s34 ("s34"), pLCE (GFP control), or pLCE-sCXCR4 ("sCXCR4," a control sponge). HEK293T cells were transfected using Eugene 6 reagent (Promega). Forty-eight hours after transfection, viral media was collected, filtered, and concentrated using Amicon Ultra centrifugal columns, and then added to the culture medium atop IMR-32 cells, and integration of sponge sequence was monitored by GFP expression. Cells with the top 30% mean GFP fluorescence were sorted using FACS analysis (ARIA II) and grown in culture until sufficient numbers of cells were available for RNA isolation via TRIzol (Life Technologies) or for protein isolation via RIPA buffer (Pierce).

2.2.8. Dual luciferase assays

Binding of miR-34a to ADRA2A was assessed using the above-described FLUC and RLUC-based indicator vectors, pL-SV40-GL3 and pL-SV40-RLUC. HEK293T cells were cotransfected with pC-34a, pL-SV40-RLUC, and either pL-GL3-ADRA2A-maj or pL-GL3-ADRA2A-min using Eugene 6 transfection reagent (Promega). To assess dose-dependent binding, 20 fmol, 60 fmol, and 120 fmol of pC-34a were used. For seed vs compensatory binding experiments, 150 fmol of pC-34 was used.

Seventy-two hours after transfection, cells were collected and lysed. Renilla luciferase and FLUC levels were measured on a SpectraMax microplate reader (Molecular Devices) using substrates from a dual luciferase reporter assay kit as described (Promega); FLUC values were normalized to RLUC values.

2.2.9. RT-qPCR and western blotting

miRNA expression was measured by stem-loop reverse transcription quantitative PCR (RT-qPCR). Total RNA was prepared with TRIzol reagent according to the manufacturer's instructions and 10 ng of total RNA per reverse transcription reaction was used. Primers for RT and probes for quantitative PCR (TaqMan) were ordered from Life Technologies and used as directed. miR-34a expression levels (microRNA assay number 000,426; Life Technologies) were normalized to RNU48 (human samples) or U87 (rat samples) (assay numbers 001,006 and 001,712, respectively; Life Technologies).

For analysis of ADRA2A mRNA expression, RNA was first treated with DNaseI (New England Biolabs) to remove genomic DNA. Random primers were used for reverse transcription as described (High Capacity Reverse Transcription Kit, Life Technologies) and SYBR reagents were used for qPCR with transcript specific primers as instructed (Life Technologies). Primer sequences for detecting ADRA2A mRNA in humans and rats are shown in Supplementary Table 1 (available online at <http://links.lww.com/PAIN/A354>). Endogenous control primers were designed against B2M (human) and Beta actin (Rat); their sequences are also shown in Supplementary Table 1 (available online at <http://links.lww.com/PAIN/A354>).

To compare α 2A-AR protein levels after miR-34a knockdown, protein was isolated from cells using RIPA buffer (ThermoFisher Scientific) and protease inhibitors (ThermoFisher Scientific). Total protein concentrations were measured with a BCA protein assay (Pierce) and equal amounts of protein (26 μ g per lane) were loaded onto a 4% to 12% Bis-Tris gel (Life Technologies) and analyzed by western blotting. Primary rabbit anti-ADRA2A was from Boster Antibody and ELISA Experts (PA2197), and primary rabbit anti- β -actin was from Cell Signaling Technology (13E5, #4970). Secondary anti-rabbit IgG HRP was from Cell Signaling Technology (#7074). Signals were developed using ECL western blotting substrate (Pierce) and in accordance with product literature, bands corresponding to ~50 kDa were identified as ADRA2A.

2.2.10. Animal studies

The US Army Institute of Surgical Research Institutional Animal Care and Use Committee approved the described animal study, which was conducted in compliance with the Animal Welfare Act, Animal Welfare Regulations, and the principles of the Guide for the Care and Use of Laboratory Animals and conformed to federal guidelines. All efforts were made to minimize the number of animals used in these experiments and to minimize potential suffering.

Six adult (200–225g) male Sprague-Dawley rats (Charles River Laboratories, Wilmington, MA) were pair-housed in a 12:12 hour light/dark cycle with ad libitum access to food and water. To induce stress, rats were individually placed in cylindrical polyethylene containers measuring 30-cm in diameter and 50-cm in height and containing 20-cm of room temperature water for 20 minutes, as previously described.⁴⁶ This was sufficient to force the rats to swim, as they were unable to reach the bottom of the cylinder. Baseline (no stress) subjects were placed in empty cylinders. Swim sessions were performed once daily for three consecutive days. All rats were towel-dried and returned to clean cages following each session and were observed for behavioral signs of distress.

Twenty-four hours following stress/sham exposure, rats were given an overdose of pentobarbital sodium. The following tissues were collected: adrenal glands, peripheral nerves, and dorsal root ganglion from L2 to L5. Each tissue was isolated and immediately stored in RNA later (ThermoFisher) at -80°C . RNA was isolated using TRIzol and RNA concentration was measured using a NanoDrop 2000 (Thermo Scientific).

3. Results

3.1. Cohort characteristics

Characteristics of study participants in the initial cohort (MVC, $n = 948$) and replication cohort (sexual assault, $n = 84$) are shown in **Table 1**. In the MVC cohort, all participants were European American and the majority of the participants were women. For the sexual assault cohort, the majority of the participants were European American and all study participants were women. In general, individuals experiencing MVC were older than sexual assault participants, reported less overall MSP in the ED, and had less distress at the time of trauma.

3.2. ADRA2A rs3750625 is located within the seed region of a predicted binding site for miR-34a

Based on computational data from the online algorithm, miRdSNP, ADRA2A rs3750625 is located in the seed binding region for miR-34a.¹² The minor allele, A, at ADRA2A rs3750625 is predicted to create an A-U basepair between miR-34a and ADRA2A, increasing seed region binding affinity between

Table 1

Baseline characteristics of study participants.

Characteristic	MVC	SA
Enrolled, n	948	84
Age [mean (SD)], y	36 (13)	26 (8)
Females, n (%)	575 (61)	84 (100)
Ethnicity		
European American	948 (100)	54 (64)
African American	—	30 (36)
Education, n (%)		
8–11 y	42 (4)	11 (13)
HS	184 (19)	16 (19)
Post-HS training (not college)	57 (6)	1 (1)
Some college	311 (33)	43 (51)
College	237 (25)	9 (11)
Postcollege	113 (12)	3 (4)
Overall ED pain, 0–10 NRS, mean (SD)	5.5 (2.4)	6.8 (2.9)
Distress in the early aftermath of trauma, PDI scale (MVC) and ASD (SA), median	18	77

ASD, acute stress disorder scale; ED, emergency department; HS, high-school; MVC, motor vehicle collision; NRS, numeric rating scale; PDI, peritraumatic distress inventory; SA, sexual assault; SD, standard deviation.

miR-34a and ADRA2A. This position is unpaired if miR-34a binds the ADRA2A major allele (C, **Fig. 1A**). Examination of the full miR-34a binding region of ADRA2A revealed that this region in general, and the seed region in particular, is highly conserved across mammalian species, suggesting that this is an important miRNA binding site (**Fig. 1B**).

3.3. Acute MSP following MVC and sexual assault

In initial general linear models, the influence of ADRA2A rs3750625 on acute MSP severity after both MVC and sexual assault depended on the level of peritraumatic stress (**Table 2**). Because of this interaction, the influence of ADRA2A rs3750625 was subsequently evaluated among individuals with higher and lower levels of peritraumatic stress separately.

3.4. ADRA2A rs3750625 influences acute MSP severity among individuals with higher levels of peritraumatic distress

Individuals who reported higher levels of peritraumatic stress during and after sexual assault and had one or more copies of the ADRA2A rs3750625 minor allele (CA/AA) experienced more severe acute MSP following sexual assault than individuals homozygous for the major allele (CC = 9.3 vs CA/AA = 8.4, $P = 0.020$, **Fig. 2**). The same direction of effect and very similar effect size was observed in the MVC cohort among those who reported higher levels of peritraumatic stress during and after the MVC although this difference did not reach statistical significance (CC = 7.1 vs CA/AA = 6.3, $P = 0.120$, **Fig. 2**).

3.5. miR-34a binds to the 3'UTR of ADRA2A

To experimentally test whether miR-34a binds to the 3'UTR of ADRA2A, we cloned the full ADRA2A 3'UTR downstream of a firefly luciferase reporter gene and performed dual luciferase reporter assays.²⁶ Somewhat surprisingly, given imperfect predicted seed binding, miR-34a binding to the 3'UTR of ADRA2A caused a dose-dependent decrease in luciferase production with the major allele (**Fig. 3A**).

Figure 1. rs3750625 is in the seed binding region of miR-34a (A) rs3750625 (black arrow, bolded nucleotides) maps within a potential seed binding site for miR-34a. Predicted base pairing between miR-34a and the *ADRA2A* 3'UTR with either the major or minor allele is indicated by a solid vertical line (Watson-Crick) or by a gray dot (wobble). The seed region of miR-34a (nucleotides 2-8) is underlined. (B) Nucleotide conservation among 7 mammals in the miR-34a binding region of *ADRA2A*. The arrow indicates the location of rs3750625. Conserved nucleotides are grayed.

3.6. The *ADRA2A* rs3750625 allele affects miR-34a binding in vitro

We next compared binding of miR-34a to the 3'UTR of *ADRA2A* in the presence of the minor allele. Consistent with miRdSNP prediction of increased base pairing of miR-34a to the *ADRA2A* 3'UTR seed region in the presence of the minor allele (Fig. 1A), decreased luciferase production was observed in the presence of the minor allele vs the major allele at all concentrations of miR-34a (Fig. 3B).

3.7. The seed region of the miR-34a binding site in the 3'UTR of *ADRA2A* confers the majority of miR-34a binding specificity in vitro

Because we observed binding of miR-34a to the 3'UTR of *ADRA2A* with the major allele present, which is not predicted to form perfect seed pairing (suggesting a possible 3'compensatory binding mechanism⁵), we made mutations in both the seed region and the compensatory region of the *ADRA2A* 3'UTR and

tested miR-34a binding (Fig. 3C). Dual luciferase reporter assays demonstrated that the majority of binding between miR-34a and the *ADRA2A* 3'UTR is conferred through the seed sequence (Fig. 3D), and that this binding site is the only miR-34a site in the *ADRA2A* 3'UTR. (No binding was observed when this seed site was mutated).

3.8. miR-34a regulates endogenous expression of *ADRA2A* in cell culture

ADRA2A mRNA and protein levels were assayed in an IMR-32 neuroblastoma cell line stably expressing a miR-34a sponge. IMR-32 cells exhibit adrenergic neurotransmitter properties^{47,61} and express both miR-34a and *ADRA2A* (Supplementary Fig. 1, available online at <http://links.lww.com/PAIN/A354>). *ADRA2A* mRNA levels (measured via RT-qPCR) increased by 2.5-fold in the absence of miR-34a (Fig. 4A, $P < 0.05$, Mann-Whitney U test). Consistent with this result, using western blotting, we also detected an increase in the level of *ADRA2A* protein in the presence of the miR-34a sponge as compared to pLCE and sCXCR4 controls (Fig. 4B).

3.9. miR-34a and *ADRA2A* are coexpressed in rat tissues relevant to pain and stress and their expression levels change in response to forced swim stress exposure

For regulation to occur in vivo, miR-34a and *ADRA2A* must both be expressed in stress-relevant tissues. Previous animal model studies have shown that pain development following stress exposure is influenced by the release of catecholamines from the adrenal gland that then sensitize peripheral nociceptors.^{19,29} We therefore examined miR-34a and *ADRA2A* mRNA expression in the adrenal gland, peripheral nerve, and dorsal root ganglion (DRG), both before and after forced swim stress.

3.9.1. Peripheral nerve and DRG

Consistent with previous reports, *ADRA2A* mRNA and miR-34a were both expressed in the DRG^{3,25,55,56} and peripheral nerve tissue, with higher levels of each RNA in the DRG (Supplementary

Table 2

General linear model examining the relationship between candidate predictors and overall acute pain severity following motor vehicle collision (n = 922) and sexual assault (n = 80).

Variable*	Motor vehicle collision		Sexual assault	
	F	P	F	P
Age	3.83	0.051	<0.001	0.999
Sex	0.037	0.947	N/A	N/A
Peritraumatic distress†	34.19	<0.001	0.011	0.918
rs3750625‡	3.224	0.073	5.432	0.007
rs3750625 × sex	0.102	0.750	N/A	N/A
rs3750625 × distress	4.103	0.043	5.467	0.007

* Study site was also included as a categorical variable.

† For the MVC, cohort distress was assessed in the emergency department using the peritraumatic distress inventory.¹¹ For the sexual assault, cohort distress was assessed using the Acute Stress Disorder Questionnaire.¹³

‡ Dominant genetic model used for rs3750625.

Fig. 2, available online at <http://links.lww.com/PAIN/A354>. Following forced swim stress exposure, *ADRA2A* mRNA levels decreased in the peripheral nerve (Fig. 5). In contrast, in peripheral nerve tissues, miR-34a expression levels increased. No change in *ADRA2A* mRNA or miR-34a expression was detected in the DRG.

3.9.2. Adrenal gland

ADRA2A and miR-34a were also expressed in the adrenal gland at lower levels than in the peripheral nerve and DRG (Supplementary Fig. 2, available online at <http://links.lww.com/PAIN/A354>). Swim stress exposure had little to no effect on adrenal gland *ADRA2A* or miR-34a expression levels (Fig. 5).

4. Discussion

To our knowledge, the present report is the first to provide evidence that acute MSP severity following traumatic stress exposure is influenced by microRNA regulation of the expression of an important stress/pain gene (ie, *ADRA2A*). In addition, our results indicate that a specific SNP in the *ADRA2A*

gene nucleotide sequence, where miR-34a binds, affects acute MSP variability. Specifically, our results suggest that distressed individuals with the rs3750625 minor allele experience an approximately 1-point increase (on a 0-10 numeric rating scale) in acute MSP after MVC trauma or sexual assault due to the fact that the 3'UTR of the *ADRA2A* gene is bound more efficiently by miR-34a when the minor allele is present. Importantly, the influence of the rs3750625 allele was stress dependent. This finding is consistent with the known regulation of miR-34a by stress^{18,27} and suggests that activation of the same physiologic systems that result in subjective symptoms of stress/distress also increase the expression of miR-34a and enhance the impact of the rs3750625 allele on the cell physiology of stress-relevant tissues and resulting phenotype. These data also provide an example of the important influence of stress-induced hyperalgesia on clinical pain outcomes after trauma exposure and provide data supporting a physiologic mechanism by which increased stress causes increased acute MSP, distinct from traditionally hypothesized mechanisms such as muscle tension.⁴³

Donello et al previously showed that, in the absence of *ADRA2A*, animals exhibit acute hyperalgesia (vs analgesia) following exposure to sound and footshock stressors.¹⁹ Our results showing that trauma-exposed individuals with the minor allele at rs3750625 have increased miR-34a binding to *ADRA2A*, reduced *ADRA2A* transcript, and increased acute MSP are consistent with these preclinical data. In addition, data from Donello et al., together with data from Levine et al., indicate that low levels of peripherally located $\alpha 2A$ -AR receptors lead to reduced feedback inhibition of norepinephrine release in response to stress.^{19,30} Norepinephrine can sensitize sensory afferents; thus, high levels of this neurotransmitter (due to low levels of $\alpha 2A$ -AR) result in increased hyperalgesia.³¹ Our data indicate that expression levels of *ADRA2A* and miR-34a RNAs change in peripheral nerves in response to stress. If miR-34a binds *ADRA2A* more efficiently in individuals with the risk allele, then reduced levels of $\alpha 2A$ -AR in such individuals would be anticipated to result in higher circulating levels of norepinephrine and greater afferent sensitization. Future studies measuring $\alpha 2A$ -AR gene expression and circulating levels of catecholamines in trauma-exposed individuals with the major vs minor allele could test this hypothesis.

Our results and those from other groups suggest that miR-34a regulation of *ADRA2A* is likely to be tissue specific. In our study, we assayed 3 stress-sensitive rat tissues (the adrenal gland, DRG, and peripheral nerves) for biologically relevant overlapping expression of miR-34a and *ADRA2A* mRNA and for expression level changes following stress exposure. Though the two RNAs were collocated in all three tissues, change in expression following stress exposure was only consistent with direct miR-34a regulation of *ADRA2A* in the peripheral nerves. These data are consistent with previous reports, which have also suggested that miR-34a and *ADRA2A* expression changes following stress exposure are tissue (and perhaps stress) specific. For example, *ADRA2A* expression in the locus coeruleus of Wistar and Wistar-Kyoto rats exhibited no change following acute restraint stress,⁵¹ whereas *ADRA2A* was significantly downregulated in the bed nucleus of the stria terminalis in response to repeated footshock stressors.³⁴ The fact that miR-34a and *ADRA2A* interaction is likely tissue specific suggests important implications regarding potential therapeutic targeting of miRNA, since miR-34a has also been shown to play roles in tumorigenesis and cardiac function.^{6,7,17,24,48,53,57} In other words, to avoid off-target effects, such treatment would likely need to be specifically targeted to

Figure 3. Characterization and functional assessment of miR-34a binding to the 3'UTR of *ADRA2A*. (A) miR-34a binding to the 3'UTR of *ADRA2A* containing the major allele at rs3750625. Relative luciferase activity (y-axis) indicates miR-34a binding to *ADRA2A* (black bars) in a dose-dependent manner. Transfection of a control miRNA was used to determine background luciferase activity (open bars). Each experiment was repeated 3 times in triplicate; error bars show standard error of the mean. (B) Comparison of miR-34a binding to the 3'UTR of *ADRA2A* with the minor allele (gray line, gray squares) vs the major allele (black line, black circles) at rs3750625. Error bars show standard error of the mean based on 3 experimental replicates. Differences in luciferase activity between the major and minor alleles were significant at all 3 concentrations of miRNA added. $**P < 0.005$ (Mann-Whitney *U* test). (C) Relative location of the miR-34a binding site in the context of the 1.4 kb *ADRA2A* 3'UTR. Mutations that destabilize base pairing (grayed and bolded nucleotides) were made in either the seed binding region or the compensatory binding region of pL-GL3-*ADRA2A*-maj. (D) Relative luciferase activity was measured for miR-34a binding (150 fmol) to the *ADRA2A* 3'UTR with no mutations (WT), with mutations in the seed region (seed mut) or with mutations in the compensatory region (comp mut). Results are the average of 3 experiments performed in triplicate. Error bars show standard deviation.

pain-relevant tissues. Of note, in addition to *ADRA2A*, predicted or experimentally validated targets of miR-34a and/or miR-449 (a member of the same miRNA family) include gene transcripts such as *COMT*, *CRHR1*, *SLC6A1*, *NOTCH1*, and *ADCY5*.^{14,18,40,52} The effect of miR-34a/miR-449 on these transcripts, and/or polymorphisms in these genes affecting miRNA binding, may also influence acute or persistent MSP outcomes after trauma exposure. Therefore, future studies examining more general transcriptional changes in miR-34a-influenced genes following different types of stress and/or pain exposures are warranted.

We focused on rs3750625 in *ADRA2A* because it was predicted to affect a miR-34a binding site by the miRdSNP online database. Interestingly, however, because there is not perfect complementarity between the seed region of *ADRA2A* with the major allele and miR-34a, other miRNA target prediction algorithms such as TargetScan and DIANA MicroT-CDS were not able to predict a binding event at this location. Therefore, our approach enabled us to uncover a miRNA binding event that would not have been discovered using traditional algorithm-based miRNA target identification techniques. Additionally, because of the general importance of seed binding for miRNA-mRNA interactions,³⁵ we hypothesized that we would detect less binding with the major allele than we observed. This observed binding led us to test whether miR-34a bound noncanonically, ie,

via the compensatory region.⁵ Surprisingly, we found that the binding activity was in fact conferred through the seed region, providing evidence that a seed region with a single mismatch can still confer binding. Despite the evident importance of seed region binding in the present study, many nonseed region target sites (with a wobble base pair or bulge in the seed) have been identified and are of high biological relevance.^{10,16,22}

Our data are interesting in that it might shed light on a finding from a paper published in 2004 showing that a genetic association in the 3'UTR of *ADRA2A* is associated with physiological responses to stress.²³ In this manuscript, the authors showed that the minor allele of a polymorphism in a Dral site caused rapid degradation of the mRNA. Upon examination of the *ADRA2A* 3'UTR, there are 4 Dral sites; the functional polymorphism examined in the 2004 manuscript encompasses rs553668 and is located 22 nt upstream of rs3750625. Although this site is immediately outside of the miR-34a binding region, it is still possible that the allele affects miR-34a binding through indirect mechanisms such as RNA folding or protein binding.² Because miRNA was not widely studied until the early part of the millennium, it is likely that the authors of this paper did not consider potential effects of miRNA.

Some limitations should be considered when interpreting the results of this study. First, we focused on a single polymorphism

Figure 4. miR-34a regulates *ADRA2A* expression in neuroblastoma cells. miR-34a was knocked down in the IMR-32 adrenergic neuroblastoma cell line using a sponge construct, pLCE-s34. (A) *ADRA2A* mRNA expression was measured by RT-qPCR of RNA isolated from IMR-32 cells expressing pLCE-s34 (s34). Levels of *ADRA2A* mRNA expression in s34 were calculated relative to IMR-32 cells expressing empty vector (pLCE) or control sponge (sCXCR4). Quantification of mRNA was performed 3 times in duplicate. Error bars show standard deviation. * $P < 0.05$ (Mann-Whitney U test). (B) *ADRA2A* protein levels from cell lysates of sponge (and control) expressing IMR-32 cells were measured by western blotting. Quantification of band intensity was calculated using band intensity measurements from the NIH image analysis program, Image J.

instead of analyzing SNP association with acute pain outcomes using previously established haplotypes.⁵⁴ It is therefore possible that other polymorphisms in LD with rs3750625 such as rs1800763 also serve as functional polymorphisms that work cooperatively with rs3750625-mediating miRNA binding

Figure 5. Comparison of miR-34a and *ADRA2A* mRNA expression levels in Sprague-Dawley rats in response to forced swim stress exposure in the following tissues: adrenal gland (AG), peripheral nerve (PN), and dorsal root ganglion (DRG). RNA was isolated from rat tissues at baseline (no stress exposure) and from rat tissues after exposure to 30 minutes of repeated forced swim stress on 3 consecutive days. RT-qPCR was used for miR-34a and *ADRA2A* mRNA quantification. Fold change was calculated from a minimum of 3 independent experiments performed in duplicate. Error bars represent standard deviations.

changes to affect *ADRA2A* mRNA levels. Second, our genetic association results were not as significant in our large MVC cohort as they were in the smaller sexual assault cohort. This could be due to the relative frequency of the minor allele in European Americans (5% in the MVC cohort) vs in African Americans (~15%). Thus in the sexual assault cohort, which is over one-third African American ($n = 30$, 36%), we might have had more statistical power to detect differences. Third, because miR-34a and *ADRA2A* are predominately expressed in tissues that are not feasible to collect in a live human cohort, we were not able to examine whether levels of these RNA differ in individuals with the major vs minor allele. However, in future studies, it would be interesting to measure plasma catecholamine levels (with the caveat that quantification is difficult and not always reliable^{15,44}) in these individuals. Fourth, animal and cell culture experiments included in this study used a single animal line, Sprague-Dawley rats, and a single cell culture line, IMR-32 neuroblastoma cells. Future studies should evaluate experimental results in additional animal and cell lines. Finally, we did not directly measure the effects of miR-34a on MSP severity following stress exposure in animals. It would be interesting to examine acute hyperalgesia following stress exposure in animals that do not express miR-34a.

Conflict of interest statement

The authors declare no conflict of interest.

Research reported in this publication was supported by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health under Award Number R01AR0563282. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. Funding was also provided by The Mayday Fund and The American Pain Society via a Future Leaders in Pain grant and by the United States Army Medical Research and Materiel Command Combat Casualty Care Research and the Clinical and Rehabilitative Medicine Research programs. The opinions or assertions contained herein are the private views of the author and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense.

Acknowledgements

The authors would like to thank the participants for taking part in this study.

Appendix A. Supplemental Digital Content

Supplemental Digital Content associated with this article can be found online at <http://links.lww.com/PAIN/A354>.

Article history:

Received 22 July 2016

Received in revised form 12 September 2016

Accepted 29 September 2016

Available online 25 October 2016

References

1. U.S. Census Bureau. Statistical abstract of the United States: 2012. Washington, DC: U.S. Census Bureau, 2011.
2. Arnold M, Ellwanger DC, Hartsperger ML, Pfeuffer A, Stümpflen V. Cis-acting polymorphisms affect complex traits through modifications of microRNA regulation pathways. *PLoS One* 2012;7:e36694.

- [3] Bak M, Silahatoglu A, Moller M, Christensen M, Rath MF, Skryabin B, Tommerup N, Kauppinen S. MicroRNA expression in the adult mouse central nervous system. *RNA* 2008;14:432–44.
- [4] Ballina LE, Ulirsch JC, Soward AC, Rossi C, Rotolo S, Linnstaedt SD, Heafner T, Foley KA, Batts J, Collette R, Holbrook D, Zelman S, McLean SA. mu-Opioid receptor gene A118G polymorphism predicts pain recovery after sexual assault. *J Pain* 2013;14:165–71.
- [5] Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009;136:215–33.
- [6] Bernardo BC, Gao XM, Winbanks CE, Boey EJ, Tham YK, Kiriazis H, Gregorevic P, Obad S, Kauppinen S, Du XJ. Therapeutic inhibition of the miR-34 family attenuates pathological cardiac remodeling and improves heart function. *Proc Natl Acad Sci U S A* 2012;109:17615–20.
- [7] Boon RA, Iekushi K, Lechner S, Seeger T, Fischer A, Heydt S, Kaluza D, Tréguer K, Carmona G, Bonauer A. MicroRNA-34a regulates cardiac ageing and function. *Nature* 2013;495:107–10.
- [8] Bortsov AV, Diatchenko L, McLean SA. Complex multilocus effects of catechol-O-methyltransferase haplotypes predict pain and pain interference 6 weeks after motor vehicle collision. *Neuromolecular Med* 2014;16:83–93.
- [9] Bortsov AV, Smith JE, Diatchenko L, Soward AC, Ulirsch JC, Rossi C, Swor RA, Hauda WE, Peak DA, Jones JS, Holbrook D, Rathlev NK, Foley KA, Lee DC, Collette R, Domeier RM, Hendry PL, McLean SA. Polymorphisms in the glucocorticoid receptor co-chaperone FKBP5 predict persistent musculoskeletal pain after traumatic stress exposure. *PAIN* 2013;154:1419–26.
- [10] Brodersen P, Voinnet O. Revisiting the principles of microRNA target recognition and mode of action. *Nat Rev Mol Cell Biol* 2009;10:141–8.
- [11] Brunet A, Weiss DS, Metzler TJ, Best SR, Neylan TC, Rogers C, Fagan J, Marmar CR. The Peritraumatic Distress Inventory: a proposed measure of PTSD criterion A2. *Am J Psychiatry* 2001;158:1480–5.
- [12] Bruno AE, Li L, Kalabus JL, Pan Y, Yu A, Hu Z. miRdSNP: a database of disease-associated SNPs and microRNA target sites on 3'UTRs of human genes. *BMC genomics* 2012;13:44.
- [13] Bryant RA, Moulds ML, Guthrie RM. Acute Stress Disorder Scale: a self-report measure of acute stress disorder. *Psychol Assess* 2000;12:61.
- [14] Capuano M, Iaffaldano L, Tinto N, Montanaro D, Capobianco V, Izzo V, Tucci F, Troncone G, Greco L, Sacchetti L. MicroRNA-449a overexpression, reduced NOTCH1 signals and scarce goblet cells characterize the small intestine of celiac patients. *PLoS One* 2011;6:e29094.
- [15] Christensen NJ. The biochemical assessment of sympathoadrenal activity in man. *Clin Auton Res* 1991;1:167–72.
- [16] Clark PM, Lohr P, Quann K, Brody J, Londin ER, Rigoutsos I. Argonaute CLIP-Seq reveals miRNA targetome diversity across tissue types. *Scientific Rep* 2014;4:5947.
- [17] Concepcion CP, Han YC, Mu P, Bonetti C, Yao E, D'andrea A, Vidigal JA, Maughan WP, Ogradowski P, Ventura A. Intact p53-dependent responses in miR-34-deficient mice. *PLoS Genet* 2012;8:e1002797.
- [18] Dias BG, Goodman JV, Ahluwalia R, Easton AE, Andero R, Ressler KJ. Amygdala-dependent fear memory consolidation via miR-34a and notch signaling. *Neuron* 2014;83:906–18.
- [19] Donello JE, Guan Y, Tian M, Cheevers CV, Alcantara M, Cabrera S, Raja SN, Gil DW. A peripheral adrenoceptor-mediated sympathetic mechanism can transform stress-induced analgesia into hyperalgesia. *Anesthesiology* 2011;114:1403–16.
- [20] Dull T, Zufferey R, Kelly M, Mandel RJ, Nguyen M, Trono D, Naldini L. A third-generation lentivirus vector with a conditional packaging system. *J Virol* 1998;72:8463–71.
- [21] Ebert MS, Neilson JR, Sharp PA. MicroRNA sponges: competitive inhibitors of small RNAs in mammalian cells. *Nat Methods* 2007;4:721–6.
- [22] Elefant N, Altuvia Y, Margalit H. A wide repertoire of miRNA binding sites: prediction and functional implications. *Bioinformatics* 2011;27:3093–101.
- [23] Finley JC Jr, O'Leary M, Wester D, MacKenzie S, Shepard N, Farrow S, Lockette W. A genetic polymorphism of the alpha2-adrenergic receptor increases autonomic responses to stress. *J Appl Physiol* 2004;96:2231–9.
- [24] Forte E, Salinas RE, Chang C, Zhou T, Linnstaedt SD, Gottwein E, Jacobs C, Jima D, Li QJ, Dave SS, Luftig MA. The Epstein-Barr virus (EBV)-induced tumor suppressor microRNA MiR-34a is growth promoting in EBV-infected B cells. *J Virol* 2012;86:6889–98.
- [25] Gold MS, Dastmalchi S, Levine JD. Alpha 2-adrenergic receptor subtypes in rat dorsal root and superior cervical ganglion neurons. *PAIN* 1997;69:179–90.
- [26] Gottwein E, Cullen BR. A human herpesvirus microRNA inhibits p21 expression and attenuates p21-mediated cell cycle arrest. *J Virol* 2010;84:5229–37.
- [27] Haramati S, Navon I, Issler O, Ezra-Nevo G, Gil S, Zwang R, Hornstein E, Chen A. MicroRNA as repressors of stress-induced anxiety: the case of amygdalar miR-34. *J Neurosci* 2011;31:14191–203.
- [28] Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. The human genome browser at UCSC. *Genome Res* 2002;12:996–1006.
- [29] Khasar SG, Burkham J, Dina OA, Brown AS, Bogen O, Alessandri-Haber N, Green PG, Reichling DB, Levine JD. Stress induces a switch of intracellular signaling in sensory neurons in a model of generalized pain. *J Neurosci* 2008;28:5721–30.
- [30] Khasar SG, Dina OA, Green PG, Levine JD. Sound stress-induced long-term enhancement of mechanical hyperalgesia in rats is maintained by sympathoadrenal catecholamines. *J Pain* 2009;10:1073–7.
- [31] Khasar SG, McCarter G, Levine JD. Epinephrine produces a β -adrenergic receptor-mediated mechanical hyperalgesia and in vitro sensitization of rat nociceptors. *J Neurophysiol* 1999;81:1104–12.
- [32] Kohli U, Muszkat M, Sofowora GG, Harris PA, Friedman EA, Dupont WD, Scheinin M, Wood AJ, Stein CM, Kurnik D. Effects of variation in the human alpha2A- and alpha2C-adrenoceptor genes on cognitive tasks and pain perception. *Eur J Pain* 2010;14:154–9.
- [33] Kruger J, Rehmsmeier M. RNAhybrid: microRNA target prediction easy, fast and flexible. *Nucleic Acids Res* 2006;34:W451–454.
- [34] Lebow M, Neufeld-Cohen A, Kuperman Y, Toory M, Gil S, Chen A. Susceptibility to PTSD-like behavior is mediated by corticotropin-releasing factor receptor type 2 levels in the bed nucleus of the stria terminalis. *J Neurosci* 2012;32:6906–16.
- [35] Lewis BP, Burge CB, Bartel DP. Conserved seed pairing, often flanked by adenosines, indicates that thousands of human genes are microRNA targets. *cell* 2005;120:15–20.
- [36] Linnstaedt SD, Gottwein E, Skalsky RL, Luftig MA, Cullen BR. Virally induced cellular microRNA miR-155 plays a key role in B-cell immortalization by Epstein-Barr virus. *J Virol* 2010;84:11670–8.
- [37] McCracken JT, Badashova KK, Posey DJ, Aman MG, Scahill L, Tierney E, Arnold LE, Vitiello B, Whelan F, Chuang SZ, Davies M, Shah B, McDougle CJ, Nurni EL. Positive effects of methylphenidate on hyperactivity are moderated by monoaminergic gene variants in children with autism spectrum disorders. *Pharmacogenomics J* 2014;14:295–302.
- [38] McLean SA, Soward AC, Ballina LE, Rossi C, Rotolo S, Wheeler R, Foley KA, Batts J, Casto T, Collette R, Holbrook D, Goodman E, Rauch SA, Liberzon I. Acute severe pain is a common consequence of sexual assault. *J Pain* 2012;13:736–41.
- [39] Meloto CB, Bortsov A, Bair E, Helgeson E, Ostrom C, Smith S, Dubner R, Slade GD, Fillingim RB, Greenspan JD, Ohrbach R, Maixner W, McLean S, Diatchenko L. Modification of COMT-dependent pain sensitivity by psychological stress and sex. *PAIN* 2016;157:858–67.
- [40] Nemoto T, Mano A, Shibasaki T. miR-449a contributes to glucocorticoid-induced CRF-R1 downregulation in the pituitary during stress. *Mol Endocrinol* 2013;27:1593–602.
- [41] Nishi D, Matsuoka Y, Yonemoto N, Noguchi H, Kim Y, Kanba S. Peritraumatic distress inventory as a predictor of post-traumatic stress disorder after a severe motor vehicle accident. *Psychiatry Clin Neurosci* 2010;64:149–56.
- [42] Norris FH. Epidemiology of trauma: frequency and impact of different potentially traumatic events on different demographic groups. *J Consulting Clin Psychol* 1992;60:409.
- [43] Norton PJ, Asmundson GJ. Amending the fear-avoidance model of chronic pain: what is the role of physiological arousal? *Behav Ther* 2003;34:17–30.
- [44] Peaston RT, Weinkove C. Measurement of catecholamines and their metabolites. *Ann Clin Biochem* 2004;41:17–38.
- [45] Platts-Mills TF, Bortsov AV, Soward A, Swor RA, Jones JS, Lee DC, Peak DA, Domeier RM, Rathlev NK, Hendry PL, McLean SA. Using emergency department-based inception cohorts to determine genetic characteristics associated with long term patient outcomes after motor vehicle collision: methodology of the CRASH study. *BMC Emerg Med* 2011;11:14.
- [46] Quintero L, Moreno M, Avila C, Arcaya J, Maixner W, Suarez-Roca H. Long-lasting delayed hyperalgesia after subchronic swim stress. *Pharmacol Biochem Behav* 2000;67:449–58.
- [47] Rabinovsky ED, Le WD, McManaman JL. Differential effects of neurotrophic factors on neurotransmitter development in the IMR-32 human neuroblastoma cell line. *J Neurosci* 1992;12:171–9.
- [48] Raver-Shapira N, Marciano E, Meiri E, Spector Y, Rosenfeld N, Moskovits N, Bentwich Z, Oren M. Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. *Mol Cell* 2007;26:731–43.
- [49] Resnick HS, Kilpatrick DG, Dansky BS, Saunders BE, Best CL. Prevalence of civilian trauma and posttraumatic stress disorder in

- a representative national sample of women. *J Consult Clin Psychol* 1993; 61:984–91.
- [50] Rubin DH, Althoff RR, Ehli EA, Davies GE, Rettew DC, Crehan ET, Walkup JT, Hudziak JJ. Candidate gene associations with withdrawn behavior. *J Child Psychol Psychiatry* 2013;54:1337–45.
- [51] Sands S, Strong R, Corbitt J, Morilak D. Effects of acute restraint stress on tyrosine hydroxylase mRNA expression in locus coeruleus of Wistar and Wistar-Kyoto rats. *Brain Res Mol Brain Res* 2000;75:1.
- [52] Shenoda BB, Alexander GM, Ajit SK. Hsa-miR-34a mediated repression of corticotrophin releasing hormone receptor 1 regulates pro-opiomelanocortin expression in patients with complex regional pain syndrome. *J Translational Med* 2016;14:1.
- [53] Silber J, Jacobsen A, Ozawa T, Harinath G, Pedraza A, Sander C, Holland EC, Huse JT. miR-34a repression in proneural malignant gliomas upregulates expression of its target PDGFRA and promotes tumorigenesis. *PLoS One* 2012;7:e33844.
- [54] Small KM, Brown KM, Seman CA, Theiss CT, Liggett SB. Complex haplotypes derived from noncoding polymorphisms of the intronless α 2A-adrenergic gene diversify receptor expression. *Proc Natl Acad Sci* 2006;103:5472–7.
- [55] Stone LS, Broberger C, Vulchanova L, Wilcox GL, Hokfelt T, Riedl MS, Elde R. Differential distribution of α 2A and α 2C adrenergic receptor immunoreactivity in the rat spinal cord. *J Neurosci* 1998;18:5928–37.
- [56] Stone LS, Vulchanova L, Riedl MS, Wang J, Williams FG, Wilcox GL, Elde R. Effects of peripheral nerve injury on α -2A and α -2C adrenergic receptor immunoreactivity in the rat spinal cord. *Neuroscience* 1999;93:1399–407.
- [57] Tazawa H, Tsuchiya N, Izumiya M, Nakagama H. Tumor-suppressive miR-34a induces senescence-like growth arrest through modulation of the E2F pathway in human colon cancer cells. *Proc Natl Acad Sci* 2007; 104:15472–7.
- [58] Ulirsch JC, Weaver MA, Bortsov AV, Soward AC, Swor RA, Peak DA, Jones JS, Rathlev NK, Lee DC, Domeier RM, Hendry PL, McLean SA. No man is an island: living in a disadvantaged neighborhood influences chronic pain development after motor vehicle collision. *PAIN* 2014;155:2116–23.
- [59] Valastyan S, Weinberg RA. Assaying microRNA loss-of-function phenotypes in mammalian cells: emerging tools and their potential therapeutic utility. *RNA Biol* 2009;6:541–5.
- [60] von Schack D, Agostino MJ, Murray BS, Li Y, Reddy PS, Chen J, Choe SE, Strassle BW, Li C, Bates B, Zhang L, Hu H, Kotnis S, Bingham B, Liu W, Whiteside GT, Samad TA, Kennedy JD, Ajit SK. Dynamic changes in the microRNA expression profile reveal multiple regulatory mechanisms in the spinal nerve ligation model of neuropathic pain. *PLoS One* 2011;6: e17670.
- [61] West GJ, Uki J, Herschman HR, Seeger RC. Adrenergic, cholinergic, and inactive human neuroblastoma cell lines with the action-potential Na⁺ ionophore. *Cancer Res* 1977;37:1372–6.
- [62] Yang L, Qian Q, Liu L, Li H, Faraone SV, Wang Y. Adrenergic neurotransmitter system transporter and receptor genes associated with atomoxetine response in attention-deficit hyperactivity disorder children. *J Neural Transm* 2013;120:1127–33.