

## Changes in HIV risk behavior and seroincidence among clients presenting for repeat HIV counseling and testing in Moshi, Tanzania

Suzanne P. Fiorillo<sup>a</sup>, Keren Z. Landman<sup>a</sup>, Alison C. Tribble<sup>a</sup>, Antipas Mtalo<sup>b</sup>, Dafrosa K. Itemba<sup>b</sup>, Jan Ostermann<sup>c,d</sup>, Nathan M. Thielman<sup>a,d</sup> and John A. Crump<sup>a,d,e,f,g,\*</sup>

<sup>a</sup>Division of Infectious Diseases and International Health, Department of Medicine, Duke University Medical Center, Durham, NC, United States; <sup>b</sup>Kikundi cha Wanawake Kilimanjaro Kupambana na UKIMWI (KIWAKKUKI Women Against AIDS in Kilimanjaro), Moshi, Tanzania; <sup>c</sup>Center for Health Policy, Duke University, Durham, NC, USA; <sup>d</sup>Duke Global Health Institute, Duke University, Durham, NC, USA; <sup>e</sup>Department of Pathology, Duke University Medical Center, Durham, NC, USA; <sup>f</sup>Kilimanjaro Christian Medical Centre, Moshi, Tanzania; <sup>g</sup>Kilimanjaro Christian Medical College, Tumaini University, Moshi, Tanzania

(Received 9 November 2010; final version received 16 January 2012)

While HIV counseling and testing (HCT) has been considered an HIV preventive measure in Africa, data are limited describing behavior changes following HCT. This study evaluated behavior changes and estimated HIV seroincidence rate among returning HCT clients. Repeat and one-time testing clients receiving HCT services in Moshi, Tanzania were identified. Information about sociodemographic characteristics, HIV-related behaviors and testing reasons were collected, along with HIV serostatus. Six thousand seven hundred and twenty-seven clients presented at least once for HCT; 1235 (18.4%) were HIV seropositive, median age was 29.7 years and 3712 (55.3%) were women. 1382 repeat and 4272 one-time testers were identified. Repeat testers were more likely to be male, older, married, or widowed, and testing because of unfaithful partner or new sexual partner. One-time testers were more likely to be students and testing due to illness. At second test, repeat testers were more likely to report that partners had received HIV testing, not have concurrent partners, not suspect partners have HIV, and have partners who did not have other partners. Clients who intended to change behaviors after the first test were more likely to report having changed behaviors by remaining abstinent (OR 2.58;  $p < 0.0001$ ) or using condoms (OR 2.00;  $p = 0.006$ ) at the second test. HIV seroincidence rate was 1.49 cases/100 person-years (PY). Clients presenting for repeat HCT reported some reduction of risky behavior and improved knowledge of sexual practices and HIV serostatus of their partners. Promoting behavior change through HCT should continue to be a focus of HIV prevention efforts in sub-Saharan Africa.

**Keywords:** Tanzania; HIV; behavior; incidence; diagnosis

### Introduction

HIV Counseling and Testing (HCT) has been promoted as an HIV prevention strategy in sub-Saharan Africa. By focusing on individualized counseling using knowledge of HIV status, HCT aims to motivate people to change behaviors in order to prevent HIV transmission (Luo, 2000). With increased access to antiretroviral therapy (ART) in countries with generalized HIV epidemics, HCT also provides a means to identify HIV infection early (Voluntary HIV-1 Counseling and Testing Efficacy Study Group, 2000) and to facilitate entry into treatment programs (Van de Perre, 2000). However, for treatment programs to be sustainable there must be reduction in HIV incidence (Stover et al., 2006).

Although HCT has been considered a preventive measure against HIV infection in sub-Saharan Africa,

behavior changes following HCT are not completely understood. With the scale-up of access to ART and testing services, more people are taking advantage of HCT with many returning for repeat testing. Studies in developed countries have found that returning HCT clients are often at the highest risk for HIV infection and have not reduced their sexual risk behavior in response to counseling and testing (FERNYAK, PAGE-SHAFFER, KELLOGG, MCFARLAND, & KATZ, 2002; MacKellar et al., 2002; Norton, Elford, Sherr, Miller, & Johnson, 1997; Phillips et al., 1995). Furthermore, there is some evidence that clients may actually be more likely to acquire HIV soon after receiving the results of a seronegative HIV test (MacKellar et al., 2002). Studies in sub-Saharan Africa have shown reduced risky behavior, such as reduction of the number of partners and increases in condom use, are most pronounced

\*Corresponding author. Email: john.crump@duke.edu

Suzanne P. Fiorillo and Keren Z. Landman contributed equally to this work.

**Presented in part:** XVI International AIDS Conference, Toronto, Canada, August 13–18, 2006. Abstract WePe0382.

among testers who are HIV-seropositive or who are in serodiscordant partnerships. However, there has been little evidence to suggest that HCT outside these groups have successfully changed HIV risk behavior (Allen et al., 1992; Arthur et al., 2007; Cremin et al., 2010; Denison, O'Reilly, Schmid, Kennedy, & Sweat, 2008; Kamenga et al., 1991; Sherr et al., 2007, Turner et al., 2009; Weinhardt, Carey, Johnson, & Bickham, 1999). Several studies have examined behavioral trends among seronegative testers and some have found increased risk behavior over time in this population (Cremin et al., 2010; Matovu et al., 2007; Sherr et al., 2007)

The purpose of this study was to evaluate the sociodemographic characteristics, HIV risk behaviors and reasons for testing in a cohort of men and women presenting at a HCT center in Moshi, Tanzania. Since a large proportion of this cohort returned for recommended repeat testing (The United Republic of Tanzania Ministry of Health National AIDS Control Programme, 2005) we evaluated differences between these clients and clients who tested only once. We also studied returning HCT clients who previously tested HIV-seronegative and had reported history of sexual activity to evaluate changes in sexual risk behavior. HIV seroincidence rate was determined among repeat testers.

## Methods

### *Study location*

Participants were recruited at a freestanding HCT center operated by Kikundi cha Wanawake Kilimanjaro Kupambana na UKIMWI (KIWAKKUKI; Women Against AIDS in Kilimanjaro). KIWAKKUKI, founded in 1990, is a women-led organization providing home-based care, counseling, education, and HCT services in the Kilimanjaro Region. Initially, KIWAKKUKI was the main provider of HCT services, although other providers began testing services before the study ended. Clients were initially charged 1000 Tanzanian shillings (US \$0.95; 2003 exchange rate), but testing became free in May 2004 (Thielman et al., 2006). This was followed in September 2004 by expanding access to free ART provided through the Tanzanian government (Shorter et al., 2009). KIWAKKUKI saw an average of 13 clients each weekday (Thielman et al., 2006). Data collection was between November 2003 and January 2008.

### *HIV counseling and testing procedures*

Consecutive clients 18 years and older presenting for HCT were invited to participate. Although the refusal

rate was not consistently reported during the study period when measured it was always <5%. Presenting clients received confidential pretest counseling, including risk assessment and reduction planning with a trained, Tanzanian, Kiswahili-speaking counselor. Counseling and testing was done according to the Tanzanian Ministry of Health National AIDS Control Programme guidelines (The United Republic of Tanzania Ministry of Health National AIDS Control Programme, 2005). After informed consent was obtained, counselors administered a structured questionnaire. The questionnaire obtained data on sociodemographic characteristics, reasons for testing, HIV risk behavior, HIV testing history, and planned behavior changes after testing (Thielman et al., 2006). Client response was recorded on paper by the counselor.

After pretest counseling and questionnaire completion, a blood sample was obtained for HIV testing. HIV antibody testing was performed on whole blood using Capillus (Trinity Biotech, Bray, Wicklow, Ireland) and Determine (Abbott Laboratories, Abbott Park, IL, USA) rapid HIV antibody tests. If test results were discordant, a blood sample was tested with ELISA (Vironostika Uni-Form II plus O Ab, BioMerieux, Durham, NC, USA). If the ELISA was seronegative, no additional testing was done. If the ELISA was positive, a Western blot (Genetic Systems HIV-1 Western Blot kit, Bio-Rad, Hercules, CA, USA) was done to confirm the result (Mayhood et al., 2008). Clients received HIV results in approximately 30 minutes. Clients who tested seropositive were referred to care and treatment centers and encouraged to have sexual partners and children tested. If seronegative, clients received post-test counseling focusing on HIV prevention. Clients were encouraged to return for repeat testing three and six months after the initial test according to national guidelines.

### *Data collection*

We identified "repeat testers" as those who were seen at KIWAKKUKI at least twice during the study period and tested HIV seronegative at their first test. If repeat testers tested more than two times, data from their first and second tests were used. "One-time testers" were identified as those who presented at KIWAKKUKI once during the study period and reported not having previously tested at KIWAKKUKI or any other testing center.

Estimation of HIV seroincidence was determined among repeat testers. An HIV seroconverter was defined as a repeat tester who had a seronegative initial HIV test and a seropositive HIV test at the second visit. The period of observation was the

interval between tests. Seroincidence is reported as number of infections per 100 person-years (PY).

### Statistical analysis

Paper questionnaire data and HIV results were entered using Epi Info 2002 or Epi Info 3.3 (Center for Disease Control and Prevention, Atlanta, GA, USA) or Teleform 9.0 (Cardiff, Visa, CA, USA). Data were validated by randomly sampling 10% of the questionnaires, with an acceptable error rate of <1 error per 5 forms. During the study, the questionnaire was modified five times to improve data quality. Consequently, some behavior variables were not collected for all clients. Actual missing data due to client non-response was <0.5% for each variable.

For determining differences between repeat testers and one-time testers, data on sociodemographic information, reasons for HIV testing and HIV risk behaviors from the first test of repeat testers were compared to the same data from one-time testers. HIV risk behaviors changes and planned behavior changes were compared between first and second HCT encounters among repeat testers. Repeat testing clients' intended changes in risk behaviors reported at the first test were compared to risk behaviors reported at the second test. A stratified analysis using Mantel-Haenszel chi-square test (Mantel and Haenszel, 1959) was undertaken in order to adjust for potential confounding. The *p*-values were based on two-tailed test results and a *p*-value ≤ 0.05 was used to define statistical significance. All analyses were performed using SAS, version 9.2 (SAS Institute, Cary, NC, USA).

### Research ethics

Ethical approval was granted by Kilimanjaro Christian Medical Centre Research Ethics Committee, Tanzania National Institutes for Medical National Research Ethics Coordinating Committee, and an institutional review board of Duke University. All participants provided informed consent and were given the written consent document in Kiswahili.

## Results

### Characteristics

During the study period (4 years; 2 months), 6727 clients presented one or more times at KUWAKKUKI for a total of 8682 HCT encounters. Among clients, 5345 (79.5%) had one HCT encounter and 1382 (20.5%) had two or more encounters. The median number of encounters was 1 (range 1–6). At the first HCT, 1235 (18.4%) clients were HIV seropositive.

Women were significantly more likely to test HIV seropositive than men (OR 3.15; 95% CI 2.74, 3.63). The median age was 29.7 (range 18.0–87.3) years, 3712 (55.3%) were women and 3108 (47.2%) lived in urban Moshi (population 144,336) whereas the rest lived in rural villages (population 1,236,713) (United Republic of Tanzania National Bureau of Statistics, 2002) in the Kilimanjaro Region. Among clients, 1382 were identified as “repeat testers” and 4272 were identified as “one-time testers” (Table 1). Among repeat testers, 1296 (93.8%) reported at least one lifetime sexual partner.

### Differences between repeat testers and one-time testers

Differences in sociodemographic characteristics, HIV risk behaviors and reasons for testing were evaluated between repeat testers and one-time testers. Repeat testers at their first test were more likely to be male, older, married, testing because of suspicion of an unfaithful partner, or having a new sexual partner (*p* < 0.01 for each variable). One-time testers were more likely to be students, widowed, testing due to illness, having a sexual partner who had died or having multiple partners (*p* < 0.02 for each variable). HIV risk behaviors and reasons for testing were adjusted for possible confounding by sex and age (Table 2).

Table 1. Baseline characteristics of all clients presenting at KIWAKKUKI Centre, Moshi, Tanzania, 2003–2008, *n* = 6727.

|                               | <i>n</i> (%)     |
|-------------------------------|------------------|
| Gender                        |                  |
| Men                           | 3000 (44.7)      |
| Women                         | 3712 (55.3)      |
| Age median (range) years      | 29.7 (18.0–87.3) |
| Residence                     |                  |
| Urban                         | 3108 (47.2)      |
| Rural                         | 3482 (52.8)      |
| HIV seroprevalence            |                  |
| Total                         | 1235 (18.4)      |
| Men                           | 292 (9.7)        |
| Women                         | 942 (25.4)       |
| Number of HCT encounters      |                  |
| 1 test only                   | 5345 (79.5)      |
| 2 tests                       | 833 (12.4)       |
| 3 tests                       | 530 (7.9)        |
| 4 tests                       | 16 (0.2)         |
| 5 tests                       | 1 (<0.1)         |
| 6 tests                       | 2 (<0.1)         |
| One-time testers <sup>a</sup> | 4272 (63.5)      |
| Repeat testers                | 1382 (20.5)      |

Missing data (*n*): 15 (gender); 18 (age); 137 (residence).

<sup>a</sup>One-time testers were HCT-naïve and received HCT only once during the study period.

Table 2. Client characteristics, reasons for testing and HIV risk behaviors for repeat testers and one-time testers at KIWAKKUKI Centre, Moshi, Tanzania, 2003–2008.

|   | Repeat testers ( <i>n</i> = 1382)<br><i>n</i> (%) | One time testers ( <i>n</i> = 4272)<br><i>n</i> (%) | <i>p</i> -Value |
|---|---|---|-----------------|
| <i>Sociodemographic characteristics</i> |   |   |                 |
| <b>Gender</b>                           |   |   |                 |
| Men                                     | 645 (46.7)  | 1820 (42.7)   | 0.01            |
| Women                                   | 737 (53.3)  | 2441 (57.3)   |                 |
| <b>Age</b>                              |   |   |                 |
| > 30 years                              | 722 (52.2)  | 2031 (47.7)   | 0.003           |
| ≤ 30 years                              | 660 (47.8)  | 2230 (52.3)   |                 |
| <b>Residence</b>                        |   |   |                 |
| Urban                                   | 620 (45.5)  | 1959 (46.9)   | 0.34            |
| Rural                                   | 744 (54.5)  | 2215 (53.1)   |                 |
| <b>Occupation</b>                       |   |   |                 |
| Business                                | 381 (27.6)  | 1208 (28.3)   | 0.61            |
| Farming                                 | 337 (24.4)  | 896 (21.0)  | 0.01            |
| Salaried worker                         | 188 (13.6)  | 563 (13.2)  | 0.69            |
| Skilled worker                          | 136 (9.8)   | 391 (9.2)   | 0.44            |
| Unskilled worker                        | 111 (8.0)   | 283 (6.6)   | 0.07            |
| Student                                 | 92 (6.7)  | 380 (8.9)   | 0.01            |
| Other                                   | 137 (9.9)   | 551 (12.9)  | 0.003           |
| <b>Education</b>                        |   |   |                 |
| Secondary or higher                     | 420 (30.5)  | 1369 (32.2)   | 0.24            |
| Primary                                 | 959 (69.5)  | 2888 (67.8)   |                 |
| <b>Marital status</b>                   |   |   |                 |
| Single                                  | 660 (47.8)  | 2118 (49.6)   | 0.24            |
| Divorced/separated                      | 131 (9.5)   | 393 (9.2)   | 0.76            |
| Cohabiting                              | 99 (7.2)  | 382 (8.9)   | 0.04            |
| Married                                 | 376 (27.2)  | 931 (21.8)  | <0.0001         |
| Widowed                                 | 116 (8.4)   | 448 (10.5)  | 0.02            |
| <b>Religion</b>                         |   |   |                 |
| Catholic                                | 578 (41.8)  | 1628 (38.1)   | 0.01            |
| Muslim                                  | 237 (17.2)  | 902 (21.1)  | 0.001           |
| Protestant                              | 511 (37.0)  | 1555 (36.4)   | 0.70            |
| Other                                   | 56 (4.0)  | 175 (4.1)   | 0.94            |
| <i>Reasons for testing<sup>a</sup></i>  |   |   |                 |
| Illness                                 | 149 (10.8)  | 937 (22.0)  | <0.0001         |
| Suspect unfaithful sexual partner       | 555 (40.2)  | 1514 (35.5)   | 0.0006          |
| Sexual partner died                     | 114 (8.3)   | 466 (10.9)  | 0.002           |
| New sexual partner                      | 85 (21.3)   | 245 (15.1)  | 0.003           |
| Multiple sexual partners                | 286 (20.7)  | 980 (23.0)  | 0.01            |
| Premarriage                             | 292 (21.1)  | 868 (20.4)  | 0.26            |
| Preconception                           | 55 (4.0)  | 167 (3.9)   | 0.79            |
| <i>HIV Risk Behaviors<sup>a</sup></i>   |   |   |                 |
| Partner died                            | 160 (12.6)  | 571 (14.7)  | 0.02            |
| Partner(s) tested for HIV               | 227 (20.4)  | 607 (17.7)  | 0.06            |
| Suspect partner(s) have HIV             | 109 (17.3)  | 349 (15.7)  | 0.49            |
| Having concurrent sexual partners       | 166 (14.6)  | 471 (13.8)  | 0.70            |
| Condom use past month                   | 66 (14.5)   | 216 (13.8)  | 0.52            |
| Had partner with other partners         | 367 (32.5)  | 1096 (32.1)   | 0.76            |

Missing data (*n*): 11 (gender); 11 (age); 116 (residence); 64 occupation; 18 (education); 18 (marital status); 12 (religion); 5 (illness); 12 (died); 13 (unfaithful partner); 18 (new partner); 14 (multiple partners); 13 (premarriage); 18 (preconception).

<sup>a</sup>All variables adjusted for sex and age.

**HIV risk behavior changes between first and second tests for repeat testers**

Compared to their first test, repeat testers at their second test were more likely to report having partners who tested for HIV, not have had concurrent sexual partners, not suspect partners are HIV-infected, not have had partners who are known to have other partners and to have used condoms in the past month ( $p < 0.04$  for each variable). There was no difference in abstinence during the past year between tests ( $p = 0.16$ ) (Table 3).

Before learning the results of their HIV test, all clients reported what behaviors they planned to change in the event of either a seropositive or a seronegative HIV test result. We compared intended behavior changes at the first HCT with behaviors reported by repeat clients at their second HCT. There were no differences in plans to change behavior between the first and second HCT for repeat testers (Table 3). However, planned intentions to change behavior at the first test were compared to the specific behavior at the second test. Clients who planned to remain abstinent after the first test if receiving a seronegative test were more likely to have remained abstinent by the second test compared to clients who did not plan to remain abstinent (OR 2.58;  $p < 0.0001$ ). Clients who planned to use condoms after the first test if receiving a seronegative test were more likely to have used condoms by the second test compared to clients who did not plan to use condoms (OR 2.00;  $p = 0.003$ ) (Table 4).

**HIV seroincidence rate**

All repeat testers were HIV seronegative at their first test and seven repeat testers had seroconverted by their second test. The median time between the first and second tests was 94 (range 24–1920) days. Repeat testers contributed 468.43 PY of follow-up between their first and second tests. Thus, the HIV seroincidence rate among repeat testers was estimated at 1.49 infections per 100 PY (95% CI; 0.39, 2.60).

**Discussion**

This study provides evidence that initially HIV-seronegative clients presenting for repeat HCT demonstrated some reduction in risky behavior but were also more knowledgeable about their partner's risk behaviors at the second test. Although partner's risk behaviors are not in the control of the client, knowing the partner's behaviors may influence the client's perception of their risk and intention to change risky behavior. By the second HCT, repeat testers were more likely to report that partners were tested for HIV and less likely to have concurrent sexual partners. They were also more likely to not suspect a partner has HIV and know that their partner does not have other partners. There were no differences in planned behavior change between the first and second tests for abstinence, using condoms and reducing the number of partners. However, clients planning to be abstinent or use condoms at the first test were more likely to report that they

Table 3. Changes in HIV risk behavior from the first and second HIV tests among previously sexually active repeat HCT clients in Moshi, Tanzania, 2003–2008.

|  | First test<br><i>n</i> (%) | Second test<br><i>n</i> (%) | <i>p</i> -Value <sup>a</sup> |
|--|----------------------------|-----------------------------|------------------------------|
| <i>Behavior change/partner knowledge</i>       |                            |                             |                              |
| Partner(s) have tested for HIV                 | 210 (20.7)                 | 332 (32.7)                  | <0.0001                      |
| Do not have concurrent sexual partners         | 792 (83.3)                 | 853 (89.7)                  | <0.0001                      |
| Do not suspect partner(s) have HIV             | 510 (82.7)                 | 556 (90.1)                  | 0.0001                       |
| Have partner(s) who do not have other partners | 622 (66.0)                 | 686 (72.7)                  | 0.001                        |
| Have used condom in past month                 | 56 (15.3)                  | 77 (21.0)                   | 0.04                         |
| Abstinent past year                            | 181 (31.3)                 | 203 (35.1)                  | 0.16                         |
| <i>Planned behavior change</i>                 |                            |                             |                              |
| Will remain abstinent                          |                            |                             |                              |
| If test seropositive                           | 826 (64.0)                 | 868 (67.2)                  | 0.07                         |
| If test seronegative                           | 574 (44.5)                 | 583 (45.2)                  | 0.66                         |
| Will use condoms                               |                            |                             |                              |
| If test seropositive                           | 249 (19.3)                 | 242 (18.8)                  | 0.63                         |
| If test seronegative                           | 263 (20.4)                 | 262 (20.3)                  | 0.88                         |
| Will reduce partners                           |                            |                             |                              |
| If test seropositive                           | 298 (23.1)                 | 325 (25.2)                  | 0.21                         |
| If test seronegative                           | 295 (22.9)                 | 330 (25.6)                  | 0.11                         |

<sup>a</sup>Adjusted for sex and age.

Table 4. Intention to reduce risky behavior at first test and actual behavior change at the second test among previous sexually active repeat HCT clients, 2003–2008.

|  | Odds ratio <sup>a</sup> | 95% CI     | <i>p</i> -Value |
|--|-------------------------|------------|-----------------|
| Planned abstinence after 1st test            |                         |            |                 |
| Had not remained abstinent prior to 2nd test | 1.00 (ref)              |            |                 |
| Remained abstinent prior to 2nd test         | 2.58                    | 1.89, 3.53 | <0.0001         |
| Planned condom use after 1st test            |                         |            |                 |
| Had not used condoms prior to 2nd test       | 1.00 (ref)              |            |                 |
| Used condoms prior to 2nd test               | 2.00                    | 1.28, 3.14 | 0.003           |

Note: OR: Odds ratio, 95% CI: 95% confidence interval.

<sup>a</sup>Adjusted for sex and age.

adhered to this behavior by the second test then those who did not plan to change behavior. Together, this provides some evidence that clients who returned for recommended follow-up HCT were more likely to have changed their own behavior and were more knowledgeable about their partners risk behaviors, both which could reduce risk for HIV infection.

Previously we have described characteristics of one-time testers (Shorter et al., 2009). However, significant differences in sociodemographic characteristics and reasons for testing were observed between one-time testers and repeat testing clients. Repeat testers were more likely to be older, male, married, testing because of new sexual partner, or suspected an unfaithful partner. One-time testers were more likely to be widowed or testing because of an illness. This suggests that repeat testers may test more often after any perceived possibility of HIV exposure whereas one-time testers are more likely to test only after a singular life event, such as being widowed, triggers concern they may be HIV infected. Consistent with our previous work (Chu et al., 2005; Landman et al., 2008; Shorter et al., 2009), we also found that women were less likely to test repeatedly and had higher HIV seroprevalence than men. It was suggested that women use HCT as a point of entry for seeking care and treatment and are more likely to present for HCT when they have an illness or are symptomatic (Shorter et al., 2009). Women may be more likely to seek HCT long after a specific exposure when they begin to experience an illness or symptoms whereas men are more likely to seek HCT sooner and more frequently after any potential exposure (Shorter et al., 2009).

We did not find differences in behavioral risk factors between repeat and one-time testers. Previous studies (Fernyak et al., 2002; MacKellar et al., 2002; Norton et al., 1997) have found that repeat testers reported more risky behavior than first time testers, although these studies were done in developed countries in different study populations. A study in

Uganda (Matovu et al., 2007) also found that HIV-seronegative repeat testers were less likely to reduce their sexual risk behaviors following repeated HCT. However, HCT populations, regardless of repeat or one-time testing, have reported higher risk behaviors than in the general population (Chu et al., 2005).

Although we found differences between repeat and one-time testers, we are unable to ascertain whether the same behavior changes observed in repeat testers would have also been observed in one-time testers if we had been able to measure their behavior at a later time. These groups were similar in terms of HIV risks behaviors but it is unknown whether one-time testers also changed their behavior after receiving HCT. In this study, follow-up HCT was recommended to all clients. However, only 20% sought repeat testing. It may be suggested that clients who follow recommendations to return for testing are those who also follow counseling advice to change behavior.

Even with some reduction in risky behavior evident in repeat testers, HIV seroincidence in this cohort was 1.49 cases per 100 PY of follow-up, remaining relatively high compared to similar cohorts. HIV seroincidence among HCT cohorts, reported as cases per 100 PY of follow-up, was 1.21 in Harare, Zimbabwe (Corbett et al., 2007), 1.4 in Rakai, Uganda (Matovu et al., 2007), 1.3 in Nairobi, Kenya (Oyugi et al., 2009), and 0.69 and 1.04 among men and women, respectively, in Chiang Mai, Thailand (Kawichai et al., 2004). The high HIV seroincidence in this group suggests further efforts in prevention are still needed. However, HIV seroprevalence in this HCT cohort was 18.4%, more than twice as high as the overall HIV seroprevalence in Tanzania of 8.8% in 2005 (UNAIDS, 2004). Therefore it is likely that seroincidence rates in our study population reflect a group at higher than average risk for HIV infection that should not be extrapolated to the general population.

There are several limitations in this study. The cohort was comprised of people who self-selected for

testing likely due to their perception of higher personal risk and the results may not be generalized to the entire population. This study also relied on self-reporting of information to counselors, which could result in inaccurate estimation of behaviors due to social desirability bias. This may be more evident when clients returned for repeat testing as they were already familiar with counseling procedures and knew the socially acceptable response to questionnaires. It is also possible that some HCT clients were misclassified as one-time testers as they may have received further HCT at another testing site or after the study period ended. Finally, since one-time testers did not return for repeat testing, we do not know if they changed their behavior after receiving initial HCT.

In conclusion, we found that clients presenting for repeat HCT reported reducing risky behavior and improved knowledge of the sexual practices and HIV serostatus of their partners. It was promising to see that clients who planned to change their behavior after the first test were more likely to report that they adhered to this change at the second test. Unfortunately, clients who did not plan to change their behavior after the first test, did not report behavior change at the second test. Therefore, a goal of HCT should be to provide counseling to help all clients make their own decisions to successfully change their behavior, specifically targeting clients who may not feel behavior change is necessary. Women represent an important target group for HCT efforts and continued reinforcement is needed to engage them in HIV education before they are sexually active and their risk increases. The high HIV seroincidence rate in this and other cohorts of repeat HCT clients in sub-Saharan Africa, despite repeatedly receiving HIV education, suggests a need to aggressively tailor education and prevention interventions for this particular high risk group. Finally, the impact of HCT on HIV prevention among HIV-seronegative clients in sub-Saharan Africa requires further study and the effect of repeat testing beyond the initial repeat test needs to be explored as recommendations for regular retesting are rolled out (Waters et al., 2011). The continued assessment of HIV knowledge and promotion of behavior change is essential.

### Acknowledgements

This study was funded in part by Roche Laboratories and by an International Studies on AIDS Associated Co-infections (ISAAC) award, a United States National Institutes of Health (NIH) funded program (U01 AI062563). Authors received support from NIH awards ISAAC (NMT, JAC); AIDS International Training and

Research Program D43 PA-03-018 (NMT, JAC); the Duke Clinical Trials Unit and Clinical Research Sites U01 AI069484 (SPF, NMT, JAC), the Duke Center for AIDS Research P30 AI 64518 (NMT, JAC); the Center for HIV/AIDS Vaccine Immunology U01 AI067854 (JAC); and the US Department of State Fulbright Program (NMT); and the Hubert-Yeargan Center for Global Health at Duke University (ACT). The authors thank the study participants and the staff and volunteers of the KIWAKKUKI AIDS Information Centre, especially the many HCT counselors Rose Mosille, Awaichi Malle, Caroline Sululu, Dayness Alexander, Alice Msuya, Rosalia Nyaky, Eunice Mbanda, Catherine Puka, Fudasia Kishe, Eliakesia Shangali, Praxed Moshi, Chrisanta Shayo, Anna Mchaki, and laboratory technician Epimack Ndanu. We acknowledge the Hubert-Yeargan Center for Global Health at Duke University for critical infrastructure support for the Kilimanjaro Christian Medical Centre-Duke University Collaboration.

### References

- Allen, S., Tice, J., Van de Perre, P., Serufilira, A., Hudes, E., Nsengumuremyi, F., ... Hulley, S. (1992). Effect of serotesting with counseling on condom use and seroconversion among HIV discordant couples in Africa. *BMJ*, *304*, 1605–1609.
- Arthur, G., Nduba, V., Forsythe, S., Mutemi, R., Odhiambo, J., & Gilks, C. (2007). Behaviour change in clients of health centre-based voluntary HIV counseling and testing services in Kenya. *Sexually Transmitted Infections*, *83*(7), 541–546.
- Chu, H.Y., Crump, J.A., Ostermann, J., Oenga, R.B., Itemba, D.K., Mgonja, A., ... Thielman, N.M. (2005). Sociodemographic and clinical characteristics of clients presenting for HIV voluntary counseling and testing in Moshi, Tanzania. *International Journal of STD & AIDS*, *16*, 691–696.
- Corbett, E.L., Makamure, B., Cheung, Y.B., Dauya, E., Matambo, R., Bandason, T., ... Hayes, R.J. (2007). HIV incidence during a cluster-randomized trial of two strategies providing voluntary counseling and testing at the workplace, Zimbabwe. *AIDS*, *19*(4), 483–489.
- Cremin, I., Nyamukapa, C., Sherr, L., Hallett, T.B., Chawira, G., Cauchernex, S., ... Gregson, S. (2010). Patterns of self-reported behaviour change associated with receiving voluntary counseling and testing in a longitudinal study from Manicaland, Zimbabwe. *AIDS and Behavior*, *14*, 708–715.
- Denison, J.A., O'Reilly, K.R., Schmid, G.P., Kennedy, C.E., & Sweat, M.D. (2008). HIV voluntary counseling and testing and behavioral risk reduction in developing countries: A meta-analysis, 1990–2005. *AIDS and Behavior*, *12*, 363–373.
- Fernyak, S.E., Page-Shafer, K., Kellogg, T.A., McFarland, W., & Katz, M.H. (2002). Risk behaviors and HIV incidence among repeat testers at publicly funded HIV testing sites in San Francisco. *Journal of Acquired Immune Deficiency Syndromes*, *31*, 63–70.

- Kamenga, M., Ryder, R.W., Jingu, M., Mbuyi, N., Mbu, L., Behets, F., ... Heyward, W.L. (1991). Evidence of marked sexual behavior change associated with low HIV-1 seroconversion in 149 married couples in discordant HIV-1 serostatus: Experience at an HIV counseling center in Zaire. *AIDS*, 5(1), 61–67.
- Kawichai, S., Beyrer, C., Khamboonruang, C., Celentano, D.D., Natpratan, C., Rungruenthanakit, K., & Nelson, K.E. (2004). HIV incidence and risk behaviours after voluntary HIV counseling and testing (HCT) among adults aged 19–35 years living in peri-urban communities around Chiang Mai city in northern Thailand, 1999. *AIDS Care*, 16(1), 21–35.
- Landman, K.Z., Ostermann, J., Crump, J.A., Mgonjas, A., Mayhood, M.K., Itemba, D.K., ... Thielman, N.M. (2008). Gender differences in the risk of HIV infection among persons reporting abstinence, monogamy and multiple sexual partners in northern Tanzania. *PLoS one*, 3(8), 3075.
- Luo, C. (2000). Achievable standard of care in low-resource settings. *Annals of the New York Academy of Sciences*, 918, 179–187.
- MacKellar, D.A., Valleroy, L.A., Secura, G.M., Bartholow, B.N., McFarland, W., Shehan, D., ... & Janssen, R.S. (2002). Repeat HIV testing, risk behaviors, and HIV seroconversion among young men who have sex with men. *Journal of Acquired Immune Deficiency Syndromes*, 29, 76–85.
- Mantel, N., & Haenszel, W. (1959). Statistical aspects of the analysis of data from retrospective studies of disease. *Journal of the National Cancer Institute*, 22(4), 719–748.
- Matovu, J.K.B., Gray, R.H., Kiwanuka, N., Kigozi, G., Wabwire-Mangen, F., Nalugoda, F., ... Wawer, M.J. (2007). Repeat voluntary HIV counseling and testing (HCT), sexual risk behavior and HIV incidence in Rakai, Uganda. *AIDS and Behavior*, 11, 71–78.
- Mayhood, M.K., Afwamba, I.A., Odhiambo, C.O., Ndanu, E., Thielman, N.M., Morrissey, A.B., ... Crump, J.A. (2008). Validation, performance under field conditions, and cost-effectiveness of Capillus HIV-1/HIV-2 and Determine HIV-1/2 rapid HIV antibody assays using sequential and parallel testing algorithms in Tanzania. *Journal of Clinical Microbiology*, 46, 3946–3951.
- Norton, J., Elford, J., Sherr, L., Miller, R., & Johnson, M.A. (1997). Repeat HIV testers at a London same-day testing clinic. *AIDS*, 11(6), 773–781.
- Oyugi, J.O., Oyugi, F.J., Otieno, C.A., Jaoko, W., Bwayo, J.J., & Anzala, O. (2009). Serologic testing algorithm for recent HIV seroconversion in estimating incidence of HIV-1 among adults visiting a HCT centre in a Kenyan tertiary health institution. *East African Medical Journal*, 86(5), 212–218.
- Phillips, K.A., Paul, J., Kegeles, S., Stall, R., Hoff, C., & Coates, T.J. (1995). Predictors of repeat HIV testing among gay and bisexual men. *AIDS*, 9, 769–775.
- Sherr, L., Lopman, B., Kakowa, M., Dube, S., Chawira, G., Nyamukapa, C., ... Gregson, S. (2007). Voluntary counseling and testing: Uptake, impact on sexual behavior, and HIV incidence in a rural Zimbabwean cohort. *AIDS*, 21(7), 851–860.
- Shorter, M.M., Ostermann, J., Crump, J.A., Tribble, A.C., Itemba, D.K., Mgonja, A., ... Thielman, N.M. (2009). Characteristics of HIV voluntary counseling and testing clients before and during care and treatment scale-up in Moshi, Tanzania. *Journal of Acquired Immune Deficiency Syndromes*, 52(5), 648–654.
- Stover, J., Bertozzi, S., Gutierrez, J.P., Walker, N., Stanek, K.A., Greener, R., ... Ghys, P.D. (2006). The global impact of scaling up HIV/AIDS prevention programs in low- and middle-income countries. *Science*, 311(5766), 1474–1476.
- The United Republic of Tanzania Ministry of Health National AIDS Control Programme. (2005). *National Guidelines for Voluntary Counseling and Testing, 2005*. Dar es Salaam, Tanzania: Ministry of Health and Social Welfare.
- The Voluntary HIV-1 Counseling and Testing Efficacy Study Group. (2000). Efficacy of voluntary HIV-1 counseling and testing in individual and couple in Kenya, Tanzania and Trinidad: A randomized trial. *Lancet*, 356, 103–112.
- Thielman, N.M., Chu, H.Y., Ostermann, J., Itemba, D.K., Mgonja, A., Mtwewe, S., ... Crump, J.A. (2006). Cost-effectiveness of free HIV voluntary counseling and testing through a community-based AIDS service organization in northern Tanzania. *American Journal of Public Health*, 96, 114–119.
- Turner, A.N., Miller, W.C., Padian, N.S., Kaufman, J.S., Behets, F.M., Chipato, T., ... Morrison, C.S. (2009). Unprotected sex following HIV testing among women in Uganda and Zimbabwe: short- and long-term comparison with pre-test behaviour. *International Journal of Epidemiology*, 38(4), 997–1007.
- UNAIDS. (2004). *Epidemiological fact sheets*. Geneva, Switzerland: United Republic of Tanzania.
- United Republic of Tanzania National Bureau of Statistics. (2002). *Population and Housing Census 2002*. Dar es Salaam, Tanzania.
- Van de Perre, P. (2000). HIV voluntary counseling and testing in community health services. *Lancet*, 356, 86–87.
- Waters, R.C., Ostermann, J., Reeves, T.D., Masnick, M.F., Thielman, N.M., Bartlett, J.A., & Crump, J.A. (2011). A cost-effectiveness analysis of alternative HIV retesting strategies in Sub-Saharan Africa. *Journal of Acquired Immune Deficiency Syndromes*, 56, 443–452.
- Weinhardt, L.S., Carey, M.P., Johnson, B.T., & Bickham, N.L. (1999). Effects of HIV counseling and testing on sexual risk behavior: A meta-analytic review of published research, 1985–1997. *American Journal of Public Health*, 89(9), 1397–1405.