

Epigenetic Regulation of *Aicda* Transcription in B Cells

by

Srikanth Kodali

Department of Biomedical Engineering
Duke University

Date: _____

Approved:

William Reichert, Supervisor

Shyam Unniraman

Charles Gersbach

Thesis submitted in partial fulfillment of
the requirements for the degree of Master of Science in the Department of
Biomedical Engineering in the Graduate School
of Duke University

2012

ABSTRACT

Epigenetic Regulation of *Aicda* Transcription in B Cells

by

Srikanth Kodali

Department of Biomedical Engineering
Duke University

Date: _____

Approved:

William Reichert, Supervisor

Shyam Unniraman

Charles Gersbach

An abstract of a thesis submitted in partial
fulfillment of the requirements for the degree
of Master of Science in the Department of
Biomedical Engineering in the Graduate School
of Duke University

2012

Copyright by
Srikanth Kodali
2012

Abstract

Activation-induced cytidine deaminase (AID), encoded by the *Aicda* gene, is indispensable for secondary antibody diversification through somatic hypermutation (SHM) and class switch recombination (CSR). It is expressed predominantly in germinal center (GC) B cells, where it deaminates cytosine to uracil in the DNA of immunoglobulin (Ig) genes, triggering mutagenesis (SHM) or deletional recombination events (CSR). However, when misregulated, AID can also mutate non-Ig genes, including proto-oncogenes, thereby contributing to genomic instability and cancer. Due to its potentially deleterious effects, AID expression must be tightly controlled. At the transcriptional level, *Aicda* is regulated by four highly conserved *cis*-regulatory regions (Regions 1-4). Region 1 contains the promoter, and is responsible for basal transcription of *Aicda*. Region 2 contains both enhancer and silencer elements, but functions as a negative regulatory region, restricting AID expression to antigen-activated B cells. Region 3 is reported to be essential for normal AID expression, but its role is otherwise unclear. Region 4, which contains two STAT6 sites essential for its function, provides enhancer activity in response to cytokine stimulation and is responsible for the high levels of AID expression found in GC B cells. Epigenetic mechanisms could add another layer of control to the regulation of *Aicda*; however, little research has been done in this regard. In this study, I investigated the role of DNA methylation in the *cis*-regulation of *Aicda* transcription. Through bisulfite sequencing of mouse splenic B cell DNA, I

demonstrate that Region 4 is highly methylated in naïve B cells, but becomes demethylated in activated B cells. This is in contrast to Region 2, which I found to be constitutively unmethylated, and Region 1, which only becomes partially demethylated. Using quantitative methylation-specific PCR (qMSP), I show that loss of methylation in Region 4 correlates positively with cell division number, consistent with a passive mechanism of demethylation. I also show that demethylation in Region 4 cannot be achieved by cell proliferation alone, and that it depends upon induction of AID expression (via IL-4 stimulation). However, bisulfite sequencing of CH12F3-2 cell DNA shows that demethylation by itself is insufficient to activate AID expression. Taken together, the data suggests that IL-4 induces STAT6 to bind to its cognate sites in Region 4 and inhibit DNMT1, likely through chromatin reorganization, triggering passive demethylation during the induction of *Aicda* transcription.

Contents

Abstract.....	iv
List of Tables.....	viii
List of Figures.....	ix
Acknowledgements.....	x
1. Introduction.....	1
1.1 The Humoral Immune Response.....	1
1.2 Mechanisms of Antibody Diversification.....	2
1.2.1 V(D)J Recombination.....	3
1.2.2 Somatic Hypermutation.....	4
1.2.3 Class Switch Recombination.....	5
1.2.4 Activation-induced Cytidine Deaminase.....	7
1.3 <i>Cis</i> -regulation of <i>Aicda</i> Expression.....	9
1.3.1 Region 1.....	9
1.3.2 Region 2.....	10
1.3.3 Region 3.....	11
1.3.4 Region 4.....	11
1.4 DNA Methylation.....	15
1.4.1 DNA Methyltransferases.....	15
1.4.2 DNA Demethylation.....	16
1.4.3 Bisulfite-based DNA Methylation Analysis.....	17
1.4.4 Non-CpG Island Methylation and AID Expression.....	19

2. Materials and Methods.....	21
2.1 Mice and Cell Lines.....	21
2.1.1 Primary B Cells.....	21
2.1.2 CH12F3-2 Cells.....	21
2.2 Flow Cytometry.....	22
2.3 DNA Preparation.....	22
2.4 Bisulfite Genomic Sequencing.....	22
2.4.1 Primers and PCR Conditions.....	22
2.4.2 Cloning and Sequencing.....	25
2.5 Quantitative Methylation-specific PCR.....	26
2.5.1 Methylation-specific PCR.....	26
3. Results.....	28
3.1 Regions 1 and 4 Become Demethylated in Activated B cells.....	28
3.1 Demethylation in Region 4 Correlates with Cell Division.....	30
4. Discussion.....	35
References.....	41

List of Tables

Table 2.1: PCR primers used for bisulfite sequencing.....	24
Table 2.2: Bisulfite PCR setup.....	25
Table 2.3: Thermal cycling conditions.....	25
Table 2.4: Primers used for qMSP.....	27
Table 2.5: qMSP setup.....	27

List of Figures

Figure 1.1 Structure of a typical antibody molecule.....	3
Figure 1.2 Mechanisms of primary and secondary antibody diversification.....	7
Figure 1.3 Organization of the mouse <i>Aicda</i> locus.....	14
Figure 2.1 <i>Aicda</i> region 1, 2, and 4 fragments covered by bisulfite sequencing.....	24
Figure 3.1 Methylation status of <i>Aicda</i>	29
Figure 3.2 Division-linked demethylation of Region 4.....	32
Figure 3.3 Average methylation at each CpG site probed in <i>Aicda</i> regions 1, 2, and 4 in unstimulated CH12F3-2 cells.....	34
Figure 4.1 IL-4-induced AID expression via STAT6-mediated demethylation.....	39

Acknowledgements

My great gratitude and respect go to my advisor, Shyam Unniraman, for taking me on as a graduate student and providing outstanding advice and support throughout my time in the lab. I would like to express my appreciation and thanks to my committee members, William Reichert and Charles Gersbach, for serving on my committee and providing constructive criticism and feedback. I would like to thank the members of the Unniraman lab for all their help and advice: Rahul Arya for mentoring me when I first joined the lab, Anurodh Agrawal for helping me set up the qMSP assay, and Zach Carico for isolating, culturing, and FACS-sorting all of the mouse splenic B cells used in this study. Finally, I would like to express my deepest thanks to my family: my father Rao, my mother Veni, and my sister Santhi, for their love and encouragement throughout the years.

1. Introduction

1.1 *The Humoral Immune Response*

The hallmark characteristics of vertebrate adaptive immunity are the ability to discriminate between self and non-self antigens, the ability to mount an exquisitely antigen-specific immune response, and the ability to retain immunological memory of an antigenic encounter (Bartl *et al.*, 2003; Willermain *et al.*, 2012). In the jawed vertebrates (gnathostomes), adaptive immunity is the responsibility of the subset of leukocytes known as lymphocytes. There are two types of lymphocytes, T cells and B cells, both of which recognize and bind antigens through surface antigen recognition receptors (Bartl *et al.*, 2003). B cells are the cellular mediators of the humoral arm of adaptive immunity. The humoral response to antigenic challenge relies upon the recruitment of antigen-specific B cell clones, each defined by the expression of a distinct antigen receptor (BCR), from a pool of mature B cells (Bartl *et al.*, 2003).

Developing B cells in the bone marrow assemble a receptor gene from a diverse pool of gene segments in order to express functional BCRs, a process known as V(D)J recombination (Xu *et al.*, 2010). These immature B cells migrate to the periphery and complete maturation. Upon encountering cognate antigen and receiving CD40L costimulation from T helper (T_h) cells, naïve B cells become activated and migrate to secondary lymphoid tissues, where they undergo rapid proliferation and differentiation in regions called germinal centers (GCs) (Xu *et al.*, 2010). In the GC, B cells modify their receptor genes through somatic hypermutation (SHM) and class switch recombination

(CSR) to generate highly-specific and effective BCRs (antibodies). The DNA editing enzyme activation-induced cytidine deaminase (AID), expressed predominantly in GC B cells, plays an essential role in CSR and SHM (LeBien and Tedder, 2008; Muramatsu *et al.*, 1999; Xu *et al.*, 2010). B cells that have successfully generated highly specific BCRs downregulate AID expression and undergo terminal differentiation into either antibody-secreting plasma cells or memory B cells (Xu *et al.*, 2010).

1.2 Mechanisms of Antibody Diversification

Membrane-bound immunoglobulin (Ig) constitutes the antigen-binding subunit of the BCR (LeBien and Tedder, 2008). A typical immunoglobulin molecule, or antibody, consists of four polypeptide chains: two identical, high molecular weight heavy (IgH) chains, and two identical, low molecular weight light (IgL) chains (Barstad *et al.*, 1974; Masat *et al.*, 1994). Each heavy chain and each light chain has a variable region followed by a constant region (Barstad *et al.*, 1974). The constant regions of the heavy chains form the Ig constant (C) region, which determines the class (isotype) of the antibody. The variable regions of each heavy and light chain pair form the Ig variable (V) regions, which serve as the antigen binding pockets (Masat *et al.*, 1994). (See Figure 1.1 for an example of antibody structure.)




Figure 1.1 Structure of a typical antibody molecule. V_L and C_L correspond to the variable and constant regions of the antibody light chain, respectively. V_H and C_H indicate the variable and constant regions of the antibody heavy chain, respectively. The variable regions of each light and heavy chain pair form the antibody variable (V) region, which establishes the antigen recognition site. The constant region of the heavy chains form the antibody constant (C) region, which determines the isotype, and therefore the effector functions, of the antibody.

The vast spectrum of potential antigens dictates that there should be a similarly diverse repertoire of antibodies in order to respond promptly to antigenic challenge. Consequently, B cells undergo multiple mechanisms of primary and secondary antibody diversification, including V(D)J recombination, SHM, and CSR. Following is an overview of these processes (see Figure 1.2).

1.2.1 V(D)J Recombination

B cells undergo germline rearrangement of their Ig loci during early development in the bone marrow (pro- to pre-B cell stages), a process known as V(D)J

recombination. The IgH variable region is encoded at loci composed of numerous variable (V_H), diversity (D_H), and joining (J_H) gene segments; the IgL variable region is encoded by V_L and J_L gene segments. At the IgH loci, V(D)J recombination brings together a V_H , a D_H , and a J_H gene segment at random to form a contiguous VDJ exon, which encodes the IgH variable region (see Figure 1.2). Subsequent rearrangement at the IgL loci results in assembly of a VJ exon, which encodes the IgL variable region.

Together, these exons code for the V region of the immunoglobulin. In addition to the combinatorial diversity resulting from the random recombination of gene segments, V(D)J recombination introduces further receptor gene variation through imprecise fusion of the V, D, and J gene segments. This results in random addition or deletion of nucleotides at the segment junctions, a phenomenon known as junctional diversity.

1.2.2 Somatic Hypermutation

SHM results in the rapid accumulation of point mutations in the V region of GC B cells (Rajewsky *et al.*, 1987). The mutation rate due to SHM is a million times greater than the background mutation rate (Xu *et al.*, 2007). Purely by chance, some of these mutations improve the affinity of the BCR to the initial antigen. Multiple rounds of SHM, positive selection of high-affinity mutants, and clonal expansion of selected cells results in the formation of B cell clones producing high affinity antibodies.

AID triggers SHM by deaminating cytosines to uracil in the Ig V region.

Transcription of the V region prior to SHM is crucial, since it allows AID access to short stretches of single-stranded DNA (ssDNA), its preferred substrate, in the V region

transcription bubbles (Roy and Lieber, 2009). If the uracils are not discovered and removed prior to DNA replication, they are mistaken for thymines by the replication machinery, resulting in transition mutations ($G \rightarrow A$) (Casali *et al.*, 2006). On the other hand, if the uracils are excised by uracil DNA glycosylase (UNG), the resulting abasic sites block the replication fork, and translesion synthesis (TLS) polymerases incorporate random nucleotides opposite these abasic sites ($G \rightarrow A$, $G \rightarrow T$, $G \rightarrow C$) (Casali *et al.*, 2006).

1.2.3 Class Switch Recombination

CSR of the IgH locus takes place concomitantly with SHM in GC B cells. The IgH locus contains multiple constant (C_H) region gene segments, and different segments encode heavy chain constant regions of different isotypes. For each C_H region (except for C_{δ}), there is a corresponding repetitive GC-rich switch (S) region that lies just upstream. CSR exchanges the initially expressed C_H region (C_{μ} in naïve B cells) for one of the downstream C_H regions, thus altering the isotype of the expressed immunoglobulin (see Figure 1.2).

In antigen-activated B cells, T_h cell-secreted cytokines (e.g. IL-4, TGF- β) induce germline transcription from intronic promoters located immediately upstream of each S region. The extent to which a particular S region is transcribed is dependent upon the composition of the cytokine milieu. AID converts cytosines to uracil on the S region ssDNA of the transcription bubbles to initiate CSR. The resulting U:G mismatches are repaired by the BER pathway to generate single-stranded breaks (SSBs). SSBs on

opposite strands are converted into double-stranded breaks (DSBs), either spontaneously if close by, or by mismatch repair (MMR) if far apart (Shrader *et al.*, 2007). Formation of DSBs in S_μ region and a downstream S region results in looping out and deletion of the intervening DNA, and end-joining of the two DSBs to complete CSR.

CSR has been shown to correlate with cell division number (Hasbold *et al.*, 1998; Hodkin *et al.*, 1996; Rush *et al.*, 2005). CD40L+IL-4-treated mouse B cells were tracked with CFSE (a fluorescent dye used to trace cells across multiple cell divisions), and shown to undergo switching to IgG1 and IgE after three and five cell divisions, respectively, irrespective of time or CD40L dose (Hasbold *et al.*, 1998). qRT-PCR for AID mRNA in division-sorted, class switch-stimulated mouse B cells has shown that AID transcript levels increase with successive cell divisions, coincident with the frequency of CSR, providing molecular mechanistic insight into the cell cycle dependence of CSR (Rush *et al.*, 2005).




Figure 1.2 Mechanisms of primary and secondary antibody diversification. During the pre- and pro-B cell stages, V(D)J recombination brings together a V, a D, and a J segment from a pool of V, D, and J segments to form a contiguous variable region exon. In germinal center B cells, somatic hypermutation introduces numerous point mutations in the variable region to improve antigen binding specificity. Concomitantly, class switch recombination exchanges the current constant region for a downstream one, looping out and deleting the intervening DNA, altering the effector functions of the antibodies produced. (Adapted from Kinoshita and Honjo, 2008.)

1.2.4 Activation-Induced Cytidine Deaminase

Activation-induced cytidine deaminase (AID), encoded by the *Aicda* gene, is a 24 kDa, 198 amino acid member of the APOBEC family of RNA/DNA editing enzymes (Muramatsu *et al.*, 1999). It was first identified through subtractive hybridization of

cDNAs between class switch-stimulated and unstimulated CH12F3-2 (a murine B cell lymphoma cell line) cells as an essential factor for CSR (Muramatsu *et al.*, 1999).

AID was initially postulated to be an RNA editing enzyme, based on its sequence homology with APOBEC1, the catalytic subunit of the multiprotein holoenzyme that edits apolipoprotein B (apoB) mRNA. APOBEC1 deaminates a cytidine to uridine at nucleotide position 6666 of the apoB mRNA, creating a premature stop codon that results in the synthesis of a truncated form of apoB (Dance *et al.*, 2000). According to the RNA editing hypothesis, AID modifies a pre-mRNA to generate a new mRNA encoding an endonuclease that nicks the V region to initiate SHM and nicks the S region to initiate CSR. However, there is no direct evidence for the RNA editing hypothesis, because no pre-mRNA substrates for AID have been identified yet.

The DNA editing hypothesis proposes that AID directly targets DNA for mutagenesis. In support of this, ectopically expressed AID has been shown to induce mutations in the genomes of humans, mice, Chinese hamsters, yeast, and *Escherichia coli*; it is improbable that AID would edit the same pre-mRNA across these diverse species to generate a DNA mutator (Martin and Scharff, 2002; Mayorov *et al.*, 2005; Petersen-Mahrt *et al.*, 2002; Yoshikawa *et al.*, 2002). Current evidence suggests that AID preferentially deaminates cytidine residues to uridine on the single-stranded DNA (ssDNA) of transcription bubbles (Larijani and Martin, 2007; Odegard and Schatz, 2006).

AID is expressed primarily in GC B cells, where it targets the Ig genes for mutagenesis and recombination, triggering SHM and CSR, respectively. Although it is

indispensable for secondary antibody diversification, AID is promiscuous, and can mutate other genes, including oncogenes, if misregulated (Komeno *et al.*, 2010; Kotani *et al.*, 2004; Pasqualucci *et al.*, 2004; Robbiani *et al.*, 2009). Indeed, a GC phenotype and dysregulated expression of AID is a feature of many B cell lymphomas (Hardianti *et al.*, 2004; Komeno *et al.*, 2010; Kotani *et al.*, 2004; Pasqualucci *et al.*, 2004). Thus AID expression must be subject to multiple layers of regulation, including at the transcriptional level.

1.3 Cis-regulation of AID Expression

The mechanism of transcriptional regulation of AID has been the subject of several previous studies. Yadav *et al.* (2005) have identified four *cis*-regulatory non-coding regions (Regions 1-4) of the mouse *Aicda* locus that are evolutionarily conserved between humans and mice (see Figure 1.3).

1.3.1 Region 1

Region 1, which is located just upstream of the transcription start site (TSS), functions as a promoter, but is not lymphoid-specific. Region 1 contains two Sp binding sites, which the general transcription factors Sp1 and Sp3 both compete for; Sp1 binding activates RNA polymerase II (pol II)-dependent transcription from the promoter, while Sp3 binding serves to repress it. Region 1 also contains one binding site each for the transcription factors STAT6, NF- κ B, and HoxC4, which are all positive regulators of *Aicda* transcription. Tran *et al.* (2010) have performed experiments in which they transiently transfected CH12F3-2 cells with luciferase expression plasmids containing

various fragments of Regions 1-4, in order to dissect the *cis*-regulation of *Aicda* transcription. Through such transient transfection assays, they have found that CIT (anti-CD40L+TGF- β +IL-4) stimulation of Region 1 does not elevate transcription beyond the basal level (four- to eightfold compared to the promoter-less plasmid). They have also identified a 100 base pair (bp) fragment just upstream of the TSS, containing only the Sp and HoxC4-Oct sites, as the minimal promoter required for basal activity.

1.3.2 Region 2

Region 2 is an intronic regulatory region, located in the first intron of the *Aicda* locus. It contains binding sites for positive regulators, including Pax5, NF- κ B, and E47, as well as negative modulators, including c-Myb and E2f.

Pax5, which is indispensable for maintaining B cell identity, is upregulated in GC B cells, which undergo SHM and CSR, but is repressed during plasma cell differentiation; thus Pax5 expression is coincident with AID expression during the GC reaction (Xu *et al.*, 2007). Moreover, enforced overexpression of Pax5 in Ba/F3 cells, a mouse pro-B cell line, has been shown to induce endogenous AID expression, implying that Pax5 directly regulates *Aicda* transcription (Nagaoka *et al.*, 2010; Xu *et al.*, 2007).

E47, a member of the E-protein family of transcription factors, is upregulated concomitantly with AID soon after B cell activation (Sayegh *et al.*, 2003). Ectopic expression of the E-protein antagonist Id3 in activated B cells hinders E47 binding to Region 2 and impairs *Aicda* transcription; conversely, overexpression of E47 in activated B cells elevates expression of AID (Sayegh *et al.*, 2003).

In contrast, E2f and c-Myb are ubiquitous repressor proteins which inhibit transcriptional activation in many cell types (Nagaoka *et al.*, 2010). Indeed, the addition of Region 2 to a luciferase reporter plasmid carrying the minimal *Aicda* promoter resulted in a twofold reduction in transcriptional activity in transfected CH12F3-2 cells (Tran *et al.*, 2010). Like Region 1, Region 2 is also unresponsive to CIT stimulation.

According to the proposed model, the silencers E2f and c-Myb act independently to repress *Aicda* transcription and prevent inappropriate AID expression, both in non-lymphoid cells and in unactivated B cells. Binding of the B cell-specific enhancers Pax5 and E47 can only partially derepress *Aicda* transcription, and so Region 2 functions as a negative regulatory region overall, in contrast to Region 1 (Tran *et al.*, 2010).

1.3.3 Region 3

Region 3 is located far downstream of the fifth (last) *Aicda* exon, and is reportedly required for physiological expression of AID, based on experiments in transgenic mice carrying a bacterial artificial chromosome (BAC) bearing mouse *Aicda* (Crouch *et al.*, 2007). Beyond that, its role remains unclear at present

1.3.4 Region 4

Region 4, which is located far upstream of Region 1, enhances transcription in response to cytokine stimulation, unlike Regions 1 and 2; thus it is the region responsible for overcoming the repression of the Region 2 silencer elements and upregulating the expression of AID in activated B cells, beyond the basal level induced by Region 1 (Tran *et al.*, 2010). This implies that Region 4 has binding sites for transcriptional activators

that are responsive to IL-4, CD40L, or TGF- β (Tran *et al.*, 2010). Indeed it contains a number of positive regulatory *cis*-elements, namely three C/EBP sites, two NF- κ B sites, two STAT6 sites, and a Smad3/4 site (Tran *et al.*, 2010). NF- κ B is an ubiquitous transcriptional factor that induces AID expression in B cells in response to CD40L costimulation.

The Smad transcription factor family members Smad3 and Smad4 are central to the intracellular TGF- β signaling pathway. Upon TGF- β -induced activation, TGF- β receptors, which possess serine/threonine kinase activity, phosphorylate Smad3, which then forms complexes with Smad4 (ten Dijke and Hill, 2004). These Smad3/Smad4 complexes are then shuttled to the nucleus, where they regulate transcription of TGF- β -responsive genes (ten Dijke and Hill, 2004). Mutation of the Smad3/4 binding site impairs the basal activity of Region 4, as well as its enhancer activity in response to TGF- β (Tran *et al.*, 2010).

STAT6 plays a crucial role in mediating IL-4-dependent immune responses. IL-4 is a critical T_h cell-secreted cytokine that stimulates activated B cells to proliferate, and induces class switching to the IgG1 and IgE isotypes (Takeda *et al.*, 1996). Upon binding to its receptor, IL-4 induces activation of Janus kinase (JAK) family members, which in turn activate STAT6 through tyrosine phosphorylation (Takeda *et al.*, 1996). Using the human B cell line BL2, Dedeoglu *et al.* (2004) have demonstrated that STAT6 binds to a cognate sequence in Region 4 (corresponding to the second STAT6 site in mouse Region 4) following IL-4 stimulation. Moreover, they have shown that IL-4 fails to induce *Aicda*

mRNA synthesis in *Stat6*^{-/-} mouse B cells. *In vitro* mutagenesis experiments demonstrate that mutation of one or both of the STAT6 sites in Region 4 severely curtails transcriptional activation by Region 4, whether induced by IL-4 alone or by CIT (Tran *et al.*, 2010). Taken together, these observations suggest that STAT6 is required for IL-4-dependent *Aicda* transcription, and that the STAT6 sites in Region 4 serve as vital enhancer elements for AID expression.

The C/EBP proteins are widely-expressed transcription factors that have been reported to function together with other transcription factors, including STAT6, to promote gene expression (Nagaoka *et al.*, 2010; Wurster *et al.*, 2000). Indeed, mutations introduced into one or both of the two C/EBP sites upstream of the Smad3/4 site abolish Region 4 responsiveness to both TGF- β alone or CIT (Tran *et al.*, 2010), demonstrating that the C/EBP sites are required for the enhancer activity of Region 4.




Figure 1.3. Organization of the mouse *Aicda* locus. Arrow marks transcriptional start site (TSS). (Adapted from Tran *et al.*, 2010.)

1.4 DNA Methylation

Epigenetic marks are heritable modifications of DNA that influence gene

expression without altering the nucleotide sequence. Many higher organisms possess the ability to epigenetically remodel their genomes through DNA methylation -- the covalent addition of a methyl group to the 5-carbon position of the cytosine ring to yield 5-methylcytosine. In mammals, methylation of cytosine residues occurs predominantly in a CpG (cytosine followed by a guanine) dinucleotide context, and is typically associated with a repressed chromatin structure and transcriptional silencing. Conversely, removal of the methyl marks induces an open chromatin state that promotes transcriptional activation.

Cytosines in a CpG context exhibit significantly higher mutability than cytosines in any other dinucleotide context. This is due to the fact that 5-methylcytosine spontaneously deaminates to thymine, which, if not corrected, results in transition mutations (Fritz and Papavasiliou, 2010). Consequently, the CpG dinucleotide is severely underrepresented in the mammalian genome. However, there are small fractions of the genome, called CpG islands, where CpGs are found in clusters (Illingworth and Bird, 2009). Although it is estimated that 70-80% of all CpGs are methylated in adult mammalian cells, most CpG islands are largely unmethylated (Feltus *et al.*, 2003; Fritz and Papavasiliou, 2010). CpG islands are frequently located in gene regulatory regions, and methylation of such islands can generate differential gene expression patterns, for example between different tissue types or stages of development (Fritz and Papavasiliou, 2010).

1.4.1 DNA Methyltransferases

A class of enzymes called DNA methyltransferases (DNMTs) is responsible for the establishment and maintenance of DNA methylation patterns by catalyzing the transfer of a methyl group from S-adenosylmethionine (SAM) to the fifth carbon of cytosine (Dhe-Paganon *et al.*, 2011). In mice and humans, the three active methyltransferases that have been identified are DNMT1, DNMT3A, and DNMT3B.

DNMT3A and DNMT3B are present at abundant levels in embryonic stem cells (ESCs), but are significantly repressed in postnatal somatic tissues (Okano *et al.*, 1999). Following the global erasure of DNA methylation in primordial germ cells and the pre-implantation embryo, DNMT3A and DNMT3B act synergistically to re-establish the genome-wide pattern of methylation during gametogenesis and embryogenesis (Okano *et al.*, 1999). Previous studies have shown that deficiency in both *Dnmt3a* and *Dnmt3b* perturbs this *de novo* methylation, and results in embryonic lethality (Okano *et al.*, 1999; Dhe-Paganon *et al.*, 2011).

DNMT1 is the most ubiquitous and abundant of the three mammalian methyltransferases (Dhe-Paganon *et al.*, 2011). During the G1 and G2 phases of the cell cycle, it is spread throughout the nucleus at low levels. However, during S phase, it is upregulated and accumulates at DNA replication foci. In contrast to DNMT3A and DNMT3B, DNMT1 has a five to thirtyfold preference for completing the methylation of hemimethylated DNA (only one strand is methylated) over catalyzing the methylation of unmethylated DNA (Okano *et al.*, 1999; Yoder *et al.*, 1997). Furthermore, inactivation of *Dnmt1* results in gradual and extensive loss of methylation, but does not disrupt *de*

novo methylation activity (Dhe-Paganon *et al.*, 2011; Lei *et al.*, 1996). Taken together, this evidence shows that DNMT1 is primarily a maintenance methyltransferase; it copies methylation marks from the template strand onto the newly synthesized strand during DNA replication, preserving existing methylation patterns across multiple rounds of the cell cycle.

1.4.2 DNA Demethylation

The removal of DNA methylation can take place via two distinct mechanisms, passive or active demethylation. Passive demethylation occurs as a result of cell division. During DNA replication, the newly synthesized strand is formed using unmodified bases. Thus, if DNMT1 is absent or blocked from accessing replication foci, over the course of multiple cell divisions there will be gradual dilution of DNA methylation marks. In contrast, active demethylation is replication-independent, and would require a demethylase to either remove 5-methylcytosine directly (generating an abasic site that is repaired through BER), remove the 5-methyl group from 5-methylcytosine, or initiate the conversion of 5-methylcytosine into another base that is subsequently converted to cytosine (Chen and Riggs, 2011). In plants, 5-methylcytosine can be removed directly by DME/ROS1 family member DNA glycosylases, but no corresponding 5-methylcytosine-specific DNA glycosylase has been identified in mammals (Chen and Riggs, 2011). However, there is evidence to suggest that active demethylation may proceed through an oxidative mechanism involving 5-hydroxymethylcytosine as an intermediate (Zhu, 2009). In this process, 5-methylcytosine

would be enzymatically oxidized to 5-hydroxymethylcytosine, which would in turn be converted to cytosine via BER (Zhu, 2009). Furthermore, since DNMT1 does not recognize 5-hydroxymethylcytosine well, conversion of 5-methylcytosine to 5-hydroxymethylcytosine may also trigger passive demethylation (Zhu, 2009). In support of this hypothesis, the ten-eleven translocation (TET) enzymes have been shown to be able to mediate oxidation of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian cells (Booth *et al.*, 2012; Zhu, 2009).

1.4.3 Bisulfite-based DNA Methylation Analysis

Many current methods for ascertaining the presence of 5-methylcytosine in target DNA sequences rely upon bisulfite-induced modification of genomic DNA. Such methods exploit the fact that sodium bisulfite efficiently deaminates unmethylated cytosine into uracil, but reacts poorly with 5-methylcytosine, leaving it unchanged (Hashimoto *et al.*, 2007; Hayatsu, 2008; Herman *et al.*, 1996). Upon PCR amplification, uracil and 5-methylcytosine are replaced in the resultant amplicon by thymine and cytosine, respectively (Hashimoto *et al.*, 2007; Hayatsu, 2008; Herman *et al.*, 1996). Bisulfite sequencing and methylation-specific PCR (MSP) are two principal techniques that have been developed for analysis of bisulfite-converted DNA. In bisulfite sequencing, primers are directed at non-CpG sequences book-ending the region of interest, in order to amplify DNA independently of methylation status (Hashimoto *et al.*, 2007). The resulting PCR product can then be cloned into a vector and sequenced to yield a target DNA methylation profile at single-base resolution (Hashimoto *et al.*, 2007;

Hayatsu, 2008). In contrast, primers for MSP are designed to overlap CpG sites in the target sequence, and thus selectively amplify methylated DNA (Herman *et al.*, 1996). MSP can be used to provide a semi-quantitative assessment of the overall methylation level of the region of interest when the ratio of product generated by methylation-specific primers versus methylation-insensitive primers is determined (Hashimoto *et al.*, 2007). The use of quantitative PCR (qPCR) rather than endpoint PCR can greatly enhance the sensitivity of this assay (Hashimoto *et al.*, 2007).

1.4.4 Non-CpG Island Methylation and AID Expression

Aicda is a gene whose expression must be tightly regulated, and the balance between enhancer elements and repressor elements in its multiple *cis*-regulatory regions serves as one layer of transcriptional control. However, given the significance of *cis*-regulatory regions, and the ability of DNA methylation to modulate transcription, methylation of *cis*-regulatory regions themselves may provide an additional layer of transcriptional control. Although there are no CpG islands located in Regions 1-4, even in CpG-poor gene regulatory elements, DNA methylation may still be an important mechanism of transcriptional regulation. Many transcription factors have been shown to bind inefficiently to DNA sequences that include, or are adjacent to, methylated CpGs. Thus CpG-containing *cis*-elements, such as the first Sp site in the *Aicda* minimal promoter, are potential candidates for epigenetic regulation. Moreover, although 45% of human gene promoters lack a CpG island, many have still been reported to have tissue-specific methylation patterns (Han *et al.*, 2011). Methylation of the non-CpG island

human β -globin, human γ -globin, and rat α -actin gene promoters resulted in transcriptional repression, albeit only in cells where these promoters have weak activity (Boyes and Bird, 1992). On the other hand, methylation of the CpG-poor human *LAMB3* and *RUNX3* promoters has been shown to directly cause transcriptional silencing even in cells that express these genes efficiently (Han *et al.*, 2011). No previous studies have investigated whether DNA methylation influences *Aicda* transcription. Therefore, I aim to elucidate the role of DNA methylation in regulating the expression of AID in B cells, in the context of its *cis*-regulatory network.

2. Materials and Methods

2.1 Mice and Cell Lines

2.1.1 Primary B Cells

Naïve splenic B cells were isolated from 13-16 wk-old female mice of the C57BL/6 (B6) inbred strain using the EasySep Mouse B Cell Enrichment Kit (Stem Cell Technologies). Cells were grown in RPMI 1640 medium (Sigma-Aldrich) supplemented with 10% (vol/vol) fetal bovine serum (FBS), 20 mM HEPES, 100 μ M non-essential amino acids, 0.1% (vol/vol) β -mercaptoethanol, 1 mM sodium pyruvate, and 1 u/mL penicillin/streptomycin/L-glutamine (PGS). Cells were either harvested at day 0 with no stimulus, or cultured for three days in the presence of LPS (10 μ g/mL) and IL-4 (10 ng/mL), anti-CD40 (1 μ g/mL) and IL-4 (10 ng/mL), or anti-CD40 (1 μ g/mL) alone. For experiments investigating the relationship between demethylation of *Aicda* and cell division, anti-CD40+IL-4-activated cells were labeled with carboxyfluorescein succinimidyl ester (CFSE) prior to culture.

2.1.2 CH12F3-2 Cells

CH12F3-2 cells were seeded at a density of 5×10^4 cells/mL in RPMI 1640 medium supplemented with 10% (vol/vol) FBS, 1 u/mL PGS, 0.1% (vol/vol) β -mercaptoethanol, and 10mM HEPES, and were harvested after two days of culture. Cells were either grown with no stimulus, or were induced to switch to IgA with a combination of IL-4 (2.5 ng/mL), TGF- β (0.15 ng/mL), and anti-CD40 (100 ng/mL).

2.2 Flow Cytometry

CFSE-labeled anti-CD40+IL-4-activated B cells were harvested at day 3 and separated by division number for the first four divisions, using a MoFlo cell sorter (Beckman Coulter). Flow cytometric data was analyzed using FlowJo software (TreeStar).

2.3 DNA Preparation

Briefly, genomic DNA was isolated from CH12F3-2 and splenic B cells using the Wizard SV Genomic Purification System (Promega), concentrated by ethanol precipitation, and quantified using an Infinite 200 PRO NanoQuant absorbance reader (Tecan). The DNA (1 µg per sample) was subjected to bisulfite modification using the EpiTect Bisulfite Kit (QIAGEN), according to the manufacturer's protocol. The bisulfite-treated DNA was then eluted in 40 µL of buffer EB (QIAGEN, 10 mM Tris-Cl, pH 8.5).

2.4 Bisulfite Genomic Sequencing

2.4.1 Primers and PCR Conditions

Bisulfite sequencing of *Aicda* regions 1, 2, and 4 was performed in order to identify differentially methylated *cis*-regulatory regions of the *Aicda* locus in B cells. Bisulfite PCR primers that amplify fragments of Regions 1, 2, and 4 were designed using the online program Bisearch (<http://bisearch.enzim.hu/>) and purchased from Sigma-Aldrich (St. Louis, MO). Analysis with MethPrimer software (<http://www.urogene.org/methprimer/index.html>) did not reveal any CpG islands in Regions 1, 2, or 4, so primers were simply selected to include as many CpG dinucleotides in the product as possible, with emphasis placed on CpGs in or adjacent to

transcription factor binding sites (see Figure 2.1). Furthermore, all primers were constrained to avoid any CpGs within their sequences in order to amplify DNA independently of methylation status. One primer set (AID_BiS(F), AID_BiS(R)) covers most of the minimal promoter element of Region 1 (-241 to -30 relative to transcription start site (TSS)), and probes five CpGs, including a CpG at the center of a Sp site. The second primer set (AID_2(2)_BiS_F, AID_2(2)BiS_R) spans part of Region 2 (+1621 to +1867 relative to TSS), and probes four CpGs, which includes one at the 3' end of an NF-κB site and one near the 3' end of an E-box. The third primer set (AID_4(1)_BiS_F, AID_4(1)_BiS_R) covers Region 4 (-8228 to -7996 relative to TSS), and probes seven CpGs, but none in or adjacent to transcription factor binding sites. (See Table 2.1 for primer information.)

Sp
Pax5
E-box
NF-κB
E2f
Cp2/c-Myb
Smad3/4
C/EBP

Aicda region 1 (-241 to -30)

TGGTACCTGGGCTGGCTTTTTAGAGGAACAGCCTCGAAGGAAGTTGGACATTAAGCATGAGCAGAAGTCC
CCCCCGCCCCCAATCATTTAATCCGTGTGGCTCTGCCACCACAGCCCCCGCCCATCTTTACTGGACCCAAC
CCAGGAGGCAGATGTTGGATACCTGGTGGTAGTGATGCTGTCTGGGGGAGGAGCCCAAGAGCAAGCT

Aicda region 2 (+1621 to +1867)

TGCATTGGCTGGGGAAGCAGCTAAAGTTCTGCCCAAGGAGCAGCTGTCTCAGCAGCTGGCGATCTACAG
GTTTGGACACCACCTAGCAAAGTCCTCCTTACCGGGAAGGACATCCCGAGGAGAGGGAGCTGGAAATAGG
CTCCTAGCTAGAGTTGAGGGGAGTGCTGGATGGAGGTGCCAGTCCACAGGTCAGGACTGTGCAGCTCCTC
CCACCGTGGCTGGAATCTTAAAATAGAAACAGTC

Aicda region 4 (-8228 to -7996)

TGTGCTGGCTCTGAGACACAGGAATGCAGCCCCCCCCCCCCCCCCCGAGGCTTTCCAGAGACCAGCCAGCCC
CTCATGCTGGCCATAACGAAACCTCACCTGTGTGTCTCATAGGGAAAGCGAGGTGGTGAAGTCCAGATGTC
CAAACTCAGCTCAGCCAGGCGACTCGATTTCCAAGTCACAGTATTTTTGGTTAAAAACAATGTGACCA

AGTTGGCTTGTGGTTTT**TCA**

Figure 2.1. *Aicda* region 1, 2, and 4 fragments covered by bisulfite sequencing. Transcription factor binding sites are highlighted. For each region, CpG sites (underlined and in blue font) are numbered in ascending order, starting from 1, relative to the TSS in the 5' to 3' direction.

Table 2.1: PCR primers used for bisulfite sequencing

Primer Name	Sequence (5' to 3')	T _{Annealing}	Product Size
AID_BiS(F)	TGGTATTGGGTTGGTTTTT	52.1 °C	212 bp
AID_BiS(R)	AACTTACTCTTATAAACTCC		
AID_2(2)_BiS_F	TGTATTGTTGGGGAAGTAGTTAAGG	58.5 °C	247 bp
AID_2(2)_BiS_R	AACTATTTCTATTTTAAAATTCCAACCAC		
AID_4(1)_BiS_F	TGTGTTGGTTTTGAGATATAGGAAT	58.5 °C	233 bp
AID_4(1)_BiS_R	TAAAAAACCACAAACCAACTTAATC		

Gradient PCR was used to optimize annealing temperatures for each primer pair, in order to avoid nonspecific amplification and to achieve maximum product yield. Each PCR was set up using the GoTaq Hot Start Polymerase kit (Promega) according to manufacturer's recommendations (see Table 2.2). All PCRs were performed in a Mastercycler ep Gradient S thermocycler (Eppendorf) under the following conditions: initial 2 min incubation at 95 °C to activate the DNA polymerase, 40 cycles of denaturation at 95 °C for 30 s, annealing at optimal annealing temperature for 30 s, and extension at 72 °C for 30 s, followed by a final 5 min extension at 72 °C (see Table 2.3). If not used immediately, all PCRs were stored at -20 °C.

Table 2.2: Bisulfite PCR setup

Component	Final Volume	Final Concentration
5X Green GoTaq Flexi Buffer	5 µL	1X
MgCl ₂ Solution, 25 mM	3 µL	3 mM
KAPA dNTP Mix, 10 mM each	0.5 µL	0.2 mM each dNTP
Forward primer, 10 µM	0.25 µL	0.1 µM
Reverse primer, 10 µM	0.25 µL	0.1 µM
GoTaq Hot Start Polymerase (5 u/µL)	0.25 µL	1.25 u
Template DNA, 25 ng/µL	1 µL	25 ng

Nuclease-free water to 25 μ L	14.75 μ L	N/A
-----------------------------------	---------------	-----

Table 2.3: Thermal cycling conditions

Step	Temperature	Time	Number of Cycles
Initial incubation	95 °C	2 min	1
Denaturation	95°C	30 s	40
Annealing	X °C*	30 s	
Extension	72 °C	30 s	
Final extension	72 °C	5 min	1
Soak	4 °C	Indefinite	1

*Optimum annealing temperature for the primer pair.

2.4.2 Cloning and Sequencing

PCR products were electrophoresed on 2% agarose-TAE gels containing 0.5 μ g/mL ethidium bromide, for 25 min at 5 V/cm. A 100 bp ladder (New England Biolabs) was used as a marker of molecular size. The resulting DNA bands were visualized with a UV transilluminator. Desired bands were excised using clean razor blades and were gel-purified using the Wizard SV Gel and PCR Clean-Up System according to the manufacturer's instructions (Promega). Each purified PCR product was ligated into the blue-white selection vector pCR 2.1-TOPO using the TOPO TA Cloning Kit as per the manufacturer's instructions (Invitrogen). Briefly, ligation was performed by incubating a mixture of PCR product (4 μ L), pCR 2.1-TOPO vector (1 μ L), and salt solution (1 μ L) at room temperature for 20 min. NEB5- α competent cells (New England Biolabs) were transformed with 3 μ L of ligation reaction, grown in 100 μ L of SOC medium for 1 h at 37 °C, combined with 60 μ L of X-Gal and 8 μ L of IPTG, and subjected to blue-white screening on Luria-Bertani (LB) agar plates containing 50 μ g/mL ampicillin (selective for pCR 2.1-TOPO). After 14 h of selection, successful recombinants (white colonies) were

harvested using sterile pipette tips and sent to Eton Bioscience (Durham, NC) for direct sequencing using the M13F(-21) universal sequencing primer (5'-TGTAACGACGGCCAGT-3'). Sequencing traces were analyzed using FinchTV software (Geospiza). The statistical significance of the methylation change at each CpG site was established using a two-tailed Student's T-test (significance level $\alpha = 0.05$). Three independent sequencing replicates (each from a different mouse) were obtained.

2.5 Quantitative Methylation-specific PCR

2.5.1 Methylation-specific PCR

Quantitative methylation-specific PCR (qMSP) was performed to determine the relationship between demethylation of Region 4 and cell division. A methylation-specific primer pair (AID4.1_MF, AID4.1M_R) that amplifies the most differentially methylated part of the previously-sequenced Region 4 fragment (-8156 to -8042 relative to TSS) was designed using MethPrimer and purchased from Sigma-Aldrich. Both forward and reverse primers were designed to include 2-3 CpGs within their sequence and to preferentially amplify methylated DNA (see Table 2.4). As done previously, gradient PCR was used to optimize the annealing temperature of the primer pair. All qMSPs were performed in triplicate in a 96-well plate (Eppendorf), in a 10 μ L volume containing 1X SYBR FAST qPCR Master Mix (Kapa Biosystems), 25 ng of bisulfite-modified DNA, and forward and reverse methylation-specific primers at a concentration of 0.25 μ M each (see Table 2.5). In order to control for the amount of input DNA in each qMSP, qPCRs using the methylation-independent primer pair for Region 4

(AID_4(1)_BiS_F, AID_4(1)_BiS_R) were run in parallel under otherwise identical conditions. All qPCRs were carried out in a Mastercycler ep realplex thermocycler (Eppendorf) according to the SYBR FAST qPCR protocol (Kapa Biosystems).

Realplex software was used for data acquisition. Data from all qPCR experiments (three independent replicates) were analyzed using the comparative Ct method (Schmittgen and Livak, 2008). All levels of methylation are reported relative to a previously-sequenced naïve B cell sample, known to be hypermethylated at Region 4.

Table 2.4: PCR primers used for qMSP

Primer Name	Sequence	T _{Annealing}	Product Size
AID4.1M_F	TCGTTAGTTTTTTATGTTGGTTATAAC	58 °C	115 bp
AID4.1M_R	AAAATCGAATCGCCTACGAC		

Table 2.5: qMSP setup

Component	Final Volume	Final Concentration
2X SYBR Green Master Mix	5 µL	1X
Forward primer, 10 µM	0.25 µL	0.25 µM
Reverse primer, 10 µM	0.25 µL	0.25 µM
Template DNA	X µL	25 ng
Nuclease-free water to 10 µL	Y µL	N/A

3. Results

3.1 Regions 1 and 4 Become Demethylated in Activated B Cells

AID expression is absent in naïve B cells, but is present at high levels in antigen-activated B cells during the GC reaction. Given this, I sought to identify *Aicda* cis-regulatory regions which are differentially methylated between unactivated and activated B cells. Thus, I performed bisulfite sequencing of portions of Regions 1 (promoter region), 2 (negative intronic regulatory region), and 4 (enhancer region) (see Figure 1.3) in genomic DNA isolated from naïve and LPS+IL-4-activated mouse splenic B cells. Region 2 is almost completely unmethylated in both unactivated and activated cells (see Figures 3.1b and 3.1e). The decrease in methylation at CpG site 1 (see Figure 2.1), the E-box proximal CpG, is not statistically significant ($P = 0.26$) (see Figure 3.1e). Region 1 is substantially methylated in unactivated cells, but becomes somewhat demethylated in activated cells (see Figures 3.1a and 3.1d). Of particular interest is CpG site 4 (see Figure 2.1), which is located within the first Sp binding site and undergoes a significant ($P = 0.011912$) decrease in methylation of 40.34 percentage points following activation (79.17% to 38.83%) (see Figure 3.1d). Region 4 exhibits the most striking change in methylation status, going from hypermethylated in unactivated cells to largely unmethylated in activated cells (see Figures 3.1c and 3.1f). Given the magnitude of this methylation change, I devoted most of my subsequent efforts toward understanding the mechanism of demethylation in Region 4. As discussed previously, Region 4 is indispensable for AID induction in response to cytokine stimulation (Tran *et*

al., 2010).




Figure 3.1. Methylation status of *Aicda*. (a-c) Lollipop diagrams showing the CpG methylation pattern of *Aicda* regulatory regions 1, 2, and 4 in B6 mouse B cells. Rows represent individual clones from bisulfite sequencing. For a given clone, each circle indicates the methylation status at a single CpG site. Black circles and white circles represent methylated and unmethylated CpGs, respectively. Unstimulated B cell sequences are on top, day 3 LPS+IL-4-activated B cell sequences are on the bottom. Data from B cells prepared from three different mice are shown. (d-f) Average methylation at each CpG site probed in *Aicda* regions 1, 2, and 4 in unactivated (black bars) and day 3 LPS+IL-4-activated (white bars) B cells.

3.2 Demethylation in Region 4 Correlates with Cell Division

It has been shown that CSR is cell division-dependent, and that *Aicda* transcription increases with successive cell divisions, coincident with the frequency of class switching (Rush *et al.*, 2005). Given this, I sought to address two specific questions: (i) Is demethylation of *Aicda* due to cell division alone, or does it coincide with the induction of *Aicda* expression? (ii) If the latter is true, then what is the relationship between cell division and demethylation of *Aicda*?

For this experiment, I made use of the fact that, in mouse B cells, anti-CD40, both alone or in conjunction with IL-4, stimulates proliferation, while anti-CD40+IL-4, but not anti-CD40 alone, induces robust *Aicda* transcription (Dedeoglu *et al.*, 2003; Nomura *et al.*, 1995). Naïve mouse splenic B cells were harvested at day 0 with no stimulus, or cultured for three days in the presence of anti-CD40, anti-CD40+IL-4, or LPS+IL-4. In order to analyze the relationship between AID expression and cell division, some of the anti-CD40+IL-4-activated B cells were labeled with the fluorescent dye CFSE prior to culture. CFSE irreversibly binds the amine groups of intracellular proteins via its succinimidyl ester moiety, and thus persists in cells for extended periods of time. Cell division results in halving of CFSE fluorescence intensity, and hence flow cytometry can be used to track division progression (Jedema *et al.*, 2004). All activated cells were harvested at day 3, and the labeled cells were then FACS-sorted into populations of uniform division number across the first four divisions, as measured by CFSE dilution. However, the relatively low numbers of cells obtained for each division posed a challenge for bisulfite sequencing. To overcome this problem, I used a quantitative methylation-specific PCR

(qMSP) approach to detect the methylation status of Region 4. Specifically, I amplified a 115 bp fragment of Region 4, using a forward primer that overlaps CpG sites 2 and 3, and a reverse primer that overlaps CpG sites 5, 6, and 7, with both primers designed to preferentially amplify methylated DNA. Naïve (~85% methylated at Region 4) and day 3 LPS+IL-4-activated (~0% methylated at Region 4) B cell DNA, whose methylation status had previously been determined by bisulfite sequencing, were used as positive and negative controls, respectively, for the specificity of the qMSP primers.

As in day 3 LPS+IL-4-activated cells, I found Region 4 to be substantially demethylated in day 3 anti-CD40+IL-4-activated cells. However, Region 4 was hypermethylated in cells cultured in the presence of anti-CD40 alone, much like in cells that were prepared without stimulation (day 0) (see Figure 3.2a). This finding was semi-quantitatively corroborated by bisulfite sequencing of Region 4 in genomic DNA prepared from the day 3 anti-CD40 and anti-CD40+IL-4 samples (see Figure 3.2c). Based on the data, demethylation of Region 4 correlates with the induction of AID gene expression.

Methylation analysis of the division-sorted B cells demonstrates that Region 4 loses methylation gradually over successive cell divisions following IL-4 stimulation (see Figure 3.2b). This is consistent with a passive, rather than active, mechanism of demethylation. In principle, downregulation of DNMT1, the maintenance DNA methyltransferase, could be responsible for the demethylation of Region 4. However, DNMT1 is highly expressed in GC B cells, and so this cannot be the case (Shaknovich *et*

al., 2011). Instead, taken together, the data suggests that DNMT1 is blocked from binding Region 4 in activated B cells due to the effects of IL-4.




Figure 3.2 Division-linked demethylation of Region 4. (a, b) The data is presented as a fold change compared to the *Aicda* region 4 methylation level in a previously sequenced unactivated (day 0) B cell DNA sample. (c) Average methylation at each CpG site probed in Region 4 in day 3 anti-CD40+IL-4-activated (green bars) and day 3 anti-CD40-activated (blue bars) B cells.

I next sought to determine whether the DNA methylation patterns of the *Aicda* loci interrogated in the mouse splenic B cells could be recapitulated in CH12F3-2 cells (a mouse B lymphoma cell line used as a B cell model). Similar to primary B cells, CH12F3-2 cells can be induced to undergo CSR upon stimulation with CIT (anti-CD40, IL-4, and TGF- β). Furthermore, basal AID expression in unstimulated CH12F3-2 cells is negligible

(Tran *et al.*, 2010). Taken together, these observations suggest that regulation of *Aicda* in CH12F3-2 cells is a good approximation of physiological reality (Tran *et al.*, 2010). Thus, I carried out bisulfite sequencing of Regions 1, 2, and 4 in genomic DNA isolated from unstimulated CH12F3-2 cells. As expected, Region 2 was unmethylated, but surprisingly, I found Regions 1 and 4 to be mostly unmethylated as well (see Figure 3.3), despite the absence of extrinsic cytokine signaling, in stark contrast to what I had observed in naïve splenic B cells (see Figure 3.2). Since AID expression is barely detectable in CH12F3-2 cells without CIT stimulation, it appears that demethylation of Region 4 is necessary, but not sufficient for induction of AID expression.




Figure 3.3 Average methylation at each CpG site probed in *Aicda* regions 1, 2, and 4 in unstimulated CH12F3-2 cells.

4. Discussion

Methylation in *cis*-regulatory regions may play an important role in controlling gene expression, even in the absence of CpG islands. In Region 1, a CpG (CpG site 4) was identified that exhibited a significant decrease in methylation following activation. Previous studies regarding the effect of Sp site methylation upon Sp1 binding are conflicting. Some investigators have reported that both Sp1 binding and transcriptional activation are unaffected by methylation at the Sp site (Harrington *et al.*, 1987; Holler *et al.*, 1988). However, other studies show that methylation of a CpCpG trinucleotide within a Sp site can hinder Sp1 binding (Clark *et al.*, 1998; Mancini *et al.*, 1999). Specifically, Clark *et al.* (1998) show that methylation of both cytosines decreases Sp1 binding by 95%, while methylation of the CpG cytosine alone only reduces Sp1 binding by 20-25%. CpG site 4 is found in a CpCpG trinucleotide context, but the outer cytosine is constitutively unmethylated in our analyses. However, given the decrease in methylation at CpG site 4 upon activation, it is conceivable that Sp1 binding in Region 1, and hence basal transcription from the *Aicda* promoter, may be partly sensitive to DNA methylation.

Region 2 is almost completely unmethylated in both unactivated and activated B cells. This is unsurprising since it functions overall as a negative regulator of AID expression. Given the strength of the repressor elements in Region 2, DNA methylation is probably not required for silencing of *Aicda* transcription in unactivated B cells.

Demethylation of Region 4 of the *Aicda* locus in activated B cells appears to take

place as a passive process during cell division. It also appears that IL-4 signaling is necessary for efficient demethylation of Region 4, while CD40 ligation (which is simply a B cell mitogen) by itself is not sufficient. Furthermore, as indicated by the CH12F3-2 data, it appears that AID expression cannot be induced by demethylation of the *Aicda* locus alone, and that it requires extrinsic cytokine stimulation as well. The two STAT6 binding sites in Region 4 are known to be crucial for the induction of AID expression in B cells under conditions of IL-4 stimulation (Tran *et al.*, 2010).

STAT6 plays a pivotal role in mediating IL-4-dependent gene expression, and is known to be critical for lineage commitment in several immune cell types (Ishii *et al.*, 2009; Kaplan *et al.*, 1996; Schroder *et al.*, 2002). In recent years, STAT6 has been demonstrated to transduce extrinsic cytokine signals into epigenetic changes in cell type-specific genes (Ishii *et al.*, 2009; Kim *et al.*, 2007; Wei *et al.*, 2010). For example, the Th2 locus control region (LCR), which regulates expression of the Th2 cytokine genes (IL-4, IL-5, and IL-13), contains a hypersensitive site that undergoes STAT6-induced demethylation in T cells stimulated by IL-4 (Kim *et al.*, 2007).

We postulate that STAT6 binds its cognate sequences in Region 4 in response to IL-4 signaling, and triggers passive demethylation during the induction of AID gene expression (see Figure 4.1). Such passive demethylation would require exclusion of DNMT1, the maintenance DNA methyltransferase, from Region 4. Recruitment of DNMT1 could potentially be inhibited by histone modifications induced by STAT6 binding. It has been reported that methylation of histone H3 lysine 9 (H3K9me)

establishes binding sites for heterochromatin protein 1 (HP1) family proteins, which directly interact with DNMT1 to mediate DNA methylation (Paul *et al.*, 2010; Smallwood *et al.*, 2007). Similarly, trimethylation of histone H3 lysine 27 (H3K27me3) is recognized by enhancer of zeste homolog 2 (EZH2), a polycomb group protein which recruits DNMT1 (Paul *et al.*, 2010; Viré, *et al.*, 2006). In CD4⁺ T cells, STAT6 has been reported to decrease H3K27 trimethylation of its target genes (Wei *et al.*, 2010). In IL-4-stimulated macrophages, STAT6 induces upregulation of H3K27 demethylase Jumonji domain containing 3 (JMJD3), a histone demethylase which removes the repressive H3K27 methylation marks from specific inflammatory genes (Ishii *et al.*, 2009). Thus, STAT6 could mediate IL-4-dependent histone demethylation of Region 4, resulting in a chromatin environment that fails to recruit DNMT1 for maintenance methylation.

DNA demethylation in Region 4 would likely be accompanied by the formation of an active chromatin configuration that is accessible to crucial trans-activating factors. At least in the case of NF- κ B, it appears that a closed chromatin architecture inhibits binding of its cognate sequence (Rau *et al.*, 2012). Still, whether NF- κ B, C/EBP, and/or Smad3/4 require Region 4 to be demethylated in order to bind remains unclear. However, a chromatin immunoprecipitation (ChIP) sequencing approach should allow this question to be addressed directly in future.

Ultimately, developing a comprehensive understanding of how *Aicda* is regulated will allow us to better understand pathways of lymphomagenesis, given that many B cell lymphomas express high levels of AID (Hardianti *et al.*, 2004; Komeno *et al.*,

2010; Kotani *et al.*, 2004; Pasqualucci *et al.*, 2004). In addition to advancing medical knowledge, understanding *Aicda* regulation in greater detail may help us to harness AID for directed evolution of proteins with desirable properties. AID has been used to artificially improve the affinity of antibodies through SHM, but given its promiscuity, it can be applied even more broadly (Chen *et al.*, 2012). Indeed, AID has been used to engineer diverse functional libraries of non-antibody proteins, such as novel fluorescent proteins and more powerful anti-apoptosis proteins (Majors, *et al.*, 2011).




Figure 4.1 IL-4-induced AID expression via STAT6-mediated demethylation. (a) In naïve B cells, Region 4 is hypermethylated, and DNMT1 maintains this methylation pattern across multiple rounds of cell division. As a result, Region 4 adopts a closed chromatin configuration, preventing the binding of trans-activating factors. Thus, the enhancer activity of Region 4 is silenced, and *Aicda* transcription is repressed. (b) IL-4 induces STAT6 to bind its cognate sequences in Region 4. STAT6 inhibits DNMT1 from accessing hemimethylated CpGs during replication (possibly

by antagonizing repressive chromatin marks that recruit DNMT1), resulting in passive demethylation. Demethylation is accompanied by formation of an open chromatin architecture, allowing trans-activating factors such as NF- κ B to bind Region 4 and induce *Aicda* transcription.

References

- Barstad, P., V. Farnsworth, M. Weigert, M. Cohn, and L. Hood, 1974. Mouse immunoglobulin heavy chains are coded by multiple germ line variable region genes. *Proceedings of the National Academy of Sciences of the United States of America* 71:4096-4100.
- Bartl S., M. Baish, I. L. Weissman, and M. Diaz, 2003. Did the molecules of adaptive immunity evolve from the innate immune system? *Integrative and Comparative Biology* 43:338-346.
- Booth, M. J., M. R. Branco, G. Ficz, D. Oxley, F. Krueger, W. Reik, and S. Balasubramanian, 2012. Quantitative sequencing of 5-methylcytosine and 5-hydroxymethylcytosine at single-base resolution. *Science* 336:934-937.
- Boyes J. and A. P. Bird, 1992. Repression of genes by DNA methylation depends on CpG density and promoter strength: evidence for involvement of a methyl-CpG binding protein. *The EMBO Journal* 11:327-333.
- Casali P., Z. Pal, Z. Xu, and H. Zan, 2006. DNA repair in antibody somatic hypermutation. *Trends in Immunology* 27:313-321.
- Chen, S., J. Qui, C. Chen, C. Liu, Y. Liu, L. An, J. Jia, J. Tang, L. Wu, and H. Hang, 2012. Affinity maturation of anti-TNF-alpha scFv with somatic hypermutation in non-B cells. *Protein & Cell* 3:460-469.
- Chen, Z. X and A. D. Riggs, 2011. DNA methylation and demethylation in mammals. *The Journal of Biological Chemistry* 286:18347-18353.
- Clark, S. J., J Harrison, and P. L. Molloy, 1997. Sp1 binding is inhibited by mCpmCpG methylation. *Gene* 195:67-71.
- Crouch, E. E., Z. Li, M. Takizawa, S. Fichtner-Feigl, P. Gourzi, C. Montano, L. Feigenbaum, P. Wilson, S. Janz, F. N. Papavasiliou, and R. Casellas, 2007. Regulation of AID expression in the immune response. *Journal of Experimental Medicine* 204:1145-1156.
- Dance, G. S. C., M. P. Sowden, Y. Yang, and H. C. Smith, 2000. APOBEC-1 dependent cytidine to uridine editing of apolipoprotein B RNA in yeast. *Nucleic Acids Research* 28:424-429.
- Dedeoglu, F., B. Horwitz, J. Chaudhari, F. W. Alt, and R. S. Geha, 2004. Induction

of activation-induced cytidine deaminase gene expression by IL-4 and CD40 ligation is dependent on STAT6 and NF- κ B. *International Immunology* 16:395-404.

Dhe-Paganon, S., F. Syeda, and L. Park, 2011. DNA methyl transferase 1: regulatory mechanisms and implications in health and disease. *International Journal of Biochemistry and Molecular Biology* 2:58-66.

Feltus, F. A., E. K. Lee, J. F. Costello, C. Plass, and P. M. Vertino, 2003. Predicting aberrant CpG island methylation. *Proceedings of the National Academy of Sciences of the United States of America* 100:12253-12258.

Fritz, F. L. and F.N. Papavasiliou, 2010. Cytidine deaminases: AIDing DNA methylation? *Genes & Development* 24:2107-2114.

Han, H., C. C. Cortez, X. Yang, P. W. Nichols, P. A. Jones, and G. Liang, 2011. DNA methylation directly silences genes with non-CpG island promoters and establishes a nucleosome occupied promoter. *Human Molecular Genetics* 20:4299-4310.

Hardianti, M. S., E. Tatsumi, M. Syampurnawati, K. Furuta, K. Saigo, S. Kawano, S. Kumagai, F. Nakamura, and Y. Matsuo, 2004. Expression of activation-induced cytidine deaminase (AID) in Burkitt lymphoma cells: rare AID-negative cell lines with the unmutated rearranged VH gene. *Leukemia & Lymphoma* 45:155-160.

Hasbold, J., A. B. Lyons, M. R. Kehry, and P. D. Hodgkin, 1998. Cell division number regulates IgG1 and IgE switching of B cells following stimulation by CD40 ligand and IL-4. *European Journal of Immunology* 28:1040-1051.

Hashimoto, K., S. Kokubun, E. Itoi, and H. Roach, 2007. Improved quantification of DNA methylation using methylation-sensitive restriction enzymes and real-time PCR. *Epigenetics* 2:86-95.

Hayatsu, H., 2008. Discovery of bisulfite-mediated cytosine conversion to uracil, the key reaction for DNA methylation analysis -- a personal account. *Proceedings of the Japan Academy Series B, Physical and Biological Sciences* 84:321-330.

Herman, J. G., J. R. Graff, S. Myohanen, B. D. Nelkin, and S. B. Baylin, 1996. Methylation-specific PCR: a novel PCR assay for methylation status of CpG islands. *Proceedings of the National Academy of Sciences of the United States of America* 93:9821-9826.

Illingworth, R. S. and A. P. Bird, 2009. CpG islands--'a rough guide'. *FEBS Letters* 583:1713-1720.

Ishii, M., H. Wen, C. A. Corsa, T. Liu, A. L. Coelho, R. M. Allen, W. F. Carson, K. A. Cavassani, X. Li, N. W. Lukacs, C. M. Hogaboam, Y. Dou, and S. L. Kunkel, 2009. Epigenetic regulation of the alternatively activated macrophage phenotype. *Blood* 114:3244-3254.

Jedema, I., N. M. van der Werff, R. M. Barge, R. Willemze, and J. H. Falkenberg, 2004. New CFSE-based assay to determine susceptibility to lysis by cytotoxic T cells of leukemic precursor cells within a heterogeneous target cell population. *Blood* 103:2677-2682.

Kaplan, M. H., U. Schindler, S. T. Smiley, and M. J. Grusby, 1996. Stat6 is required for mediating responses to IL-4 and for development of Th2 cells. *Immunity* 4:313-319.

Kim, S. T., P. E. Fields, and R. A. Flavell, 2007. Demethylation of a specific hypersensitive site in the Th2 locus control region. *Proceedings of the National Academy of Sciences of the United States of America* 104:17052-17057.

Kinoshita, K. and T. Honjo, 2001. Linking class-switch recombination with somatic hypermutation. *Nature Reviews Molecular Cell Biology* 2:493-503.

Komeno, Y., J. Kitaura, N. Watanabe-Okochi, N. Kato, T. Oki, F. Nakahara, Y. Harada, H. Harada, R. Shinkura, H. Nagaoka, Y. Hayashi, T. Honjo, and T. Kitamura, 2010. AID-induced T-lymphoma or B-leukemia /lymphoma in a mouse BMT model. *Leukemia* 24:1018-1024.

Kotani, A., N. Kakazu, T. Tsuruyama, I. Okazaki, M. Muramatsu, K. Kinoshita, H. Nagaoka, D. Yabe, and T. Honjo, 2007. Activation-induced cytidine deaminase (AID) promotes B cell lymphomagenesis in Emu-cmyc transgenic mice. *Proceedings of the National Academy of Sciences of the United States of America* 104:1616-1620.

Larijani, M., A. P. Petrov, O. Kolenchenko, M. Berru, S. N. Krylov, and A. Martin, 2007. AID associates with single-stranded DNA with high affinity and a long complex half-life in a sequence-independent manner. *Molecular and Cell Biology* 27:20-30.

LeBien, T. W. and T. Tedder, 2008. B lymphocytes: how they develop and function. *Blood* 112:1570-1580.

Lee-Theilen, M. and J. Chaudhari, 2010. Walking the AID tightrope. *Nature Immunology* 11:107-109.

Lei, H., S. P. Oh, M. Okanao, R. Juttermann, K. A. Goss, R. Jaenisch, and E. Li,

1996. De novo DNA cytosine methyltransferase activities in mouse embryonic stem cells. *Development* 122:3195-3205.

Majors, B. S., G. G. Chiang, N. Pederson, and M. J. Betenbaugh, 2011. Directed evolution of mammalian anti-apoptosis proteins by somatic hypermutation. *Protein Engineering, Design, and Selection* 25.

Mancini, D., S. M. Singh, T. K. Archer, and D. I. Rodenhiser, 1999. Site-specific DNA methylation in the neurofibromatosis (NF1) promoter interferes with binding of CREB and SP1 transcription factors. *Oncogene* 18:4108-4119.

Martin A. and M. D. Scharff, 2002. Somatic hypermutation of the AID transgene in B and non-B cells. *Proceedings of the National Academy of Sciences of the United States of America* 99:12304-12308.

Masat, L., M. Wabl, and J. P. Johnson, 1994. A simpler sort of antibody. *Proceedings of the National Academy of Sciences of the United States of America* 91:893-896.

Mayorov, V., I. B. Rogozin, L. R. Adkison, C. Frahm, T. A. Kunkel, and Y. I. Pavlov, 2005. Expression of human AID in yeast induces mutations in context similar to the context of somatic hypermutation at G-C pairs in immunoglobulin genes. *BMC Immunology* 5:10.

Muramatsu, M., V. S. Sanakaranand, S. Anant, M. Sugai, K. Kinoshita, N. O. Davidson, and T. Honjo, 1999. Specific expression of activation-induced cytidine deaminase (AID), a novel member of the RNA-editing deaminase family in germinal center B cells. *The Journal of Biological Chemistry* 274:18470-18476.

Nagaoka H., T. T. Tran, M. Kobayashi, M. Aida, and T. Honjo, 2010. Preventing AID, a physiological mutator, from deleterious activation: regulation of the genomic instability that is associated with antibody diversity. *International Immunology* 22:227-235.

Nomura J., S. Inui, T. Yamasaki, S. Kataoka, K. Maeda, K. Nakanishi, and N. Sakaguchi, 1995. Anti-CD40 monoclonal antibody induces the proliferation of murine B cells as B-cell mitogen through a distinct pathway from receptors for antigens or lipopolysaccharide. *Immunology Letters* 45:195-203.

Odegard, V. H. and D. G. Schatz, 2006. Targeting of somatic hypermutation. *Nature Reviews Immunology* 6:573-583.

Okano, M., D. W. Bell, D. A. Haber, and E. Li, 1999. DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* 99:247-257.

Pasqualucci, L., R. Guglielmino, J. Houldsworth, J. Mohr, S. Aoufouchi, R. Polakiewicz, R. S. K. Chaganti, and R. Dalla-Favera, 2004. Expression of the AID protein in normal and neoplastic B cells. *Blood* 104:3318-3325.

Paul, T.A., J. Bies, D. Small, and L. Wolff, 2010. Signatures of polycomb repression and reduced H3K4 trimethylation are associated with *p15INK4b* DNA methylation in AML. *Blood* 115:3098-3108.

Petersen-Mahrt, S. K., R. S. Harris, and M. S. Neuberger, 2002. AID mutates *E. coli* suggesting a DNA deamination mechanism for antibody diversification. *Nature* 418:99-103.

Rau, T. T., A. Rogler, M. Frischauf, A. Jung, P. C. Konturek, A. Dimmler, G. Faller, B. Sehnert, W. El-Rifai, A. Hartmann, R. E. Voll, and R. Schneider-Stock, 2012. Methylation-dependent activation of CDX1 through NF- κ B: a link from inflammation to intestinal metaplasia in the human stomach. *The American Journal of Pathology* [epub ahead of print].

Rajewsky K., I. Forster, and A. Cumano, 1987. Evolutionary and somatic selection of the antibody repertoire in the mouse. *Science* 238:1088-1094.

Robbiani, D. F., S. Bunting, N. Feldhahn, A. Bothmer, J. Camps, S. Deroubaix, K. M. McBride, I. A. Klein, G. Stone, T. R. Eisenreich, T. Ried, A. Nussenzweig, and M. C. Nussenzweig, 2009. *Molecular Cell* 36:631-641.

Roy, D. and M. Lieber, 2009. G-clustering is important for the initiation of transcription-induced R loops in vitro, whereas high G density without clustering is sufficient thereafter. *Molecular Cell Biology* 29:3124-3133.

Rush, J. S., M. Liu, V. H. Odegard, S. Unniraman, and D. G. Schatz, 2005. Expression of activation-induced cytidine deaminase is regulated by cell division, providing a mechanistic basis for division-linked class switch recombination. *Proceedings of the National Academy of Sciences of the United States of America* 102:13242-13247.

Sayegh, C. E., M. W. Quong, Y. Agata, and C. Murre, 2003. E-proteins directly regulate expression of activation-induced cytidine deaminase in mature B cells. *Nature Immunology* 4:586-593.

Schmittgen, T. D. and K. J. Livak, 2008. Analyzing real-time PCR data by the comparative CT method. *Nature Protocols* 3:1101-1108.

Schroder, A. J., P. Pavlidis, A. Arimura, D. Capece, P. B. Rothman, 2002. Cutting edge: STAT6 serves as a positive and negative regulator of gene expression in IL-4-stimulated B lymphocytes. *The Journal of Immunology* 168:996-1000.

Shaknovich, R., L. Cerchiatti, L. Tsikitas, M. Kormaksson, S. De, M. E. Figueroa, G. Ballon, S. N. Yang, N. Weinhold, M. Reimers, T. Clozel, K. Luttrup, T. J. Ekstrom, J. Frank, A. Vasanthakumar, L. A. Godley, F. Michor, O. Elemento, and A. Melnick, 2011. DNA methyltransferase 1 and DNA methylation patterning contribute to germinal center B-cell differentiation. *Blood* 118:3559-3569.

Shrader, C. E., J. E. J. Guikema, E. K. Linehan, E. Selsing, and J. Stavnezer, 2007. Activation-induced cytidine deaminase-dependent DNA breaks in class switch recombination occur during G1 phase of the cell cycle and depend upon mismatch repair. *The Journal of Immunology* 179:6064-6071.

Smallwood, A. P., Esteve, S. Pradhan, and M. Carey, 2007. Functional cooperation between HP1 and DNMT1 mediates gene silencing. *Genes & Development* 21:1169-1178.

Takeda, K., T. Tanaka, W. Shi, M. Matsumoto, M. Minami, S. Kashiwamura, K. Nakanishi, N. Yoshida, T. Kishimoto, and S. Akira, 1996. Essential role of Stat6 in IL-4 signalling. *Nature* 380:627-630.

ten Dijke, P. and C. S. Hill, 2004. New insights into TGF-beta-Smad signalling. *Trends in Biochemical Sciences* 29:265-273.

Tran, T. H., M. Nakata, K. Suzuki, N. A. Begum, R. Shinkura, S. Fagarasan, T. Honjo, and H. Nagaoka, 2010. B cell-specific and stimulation-responsive enhancers derepress Aicda by overcoming the effects of silencers. *Nature Immunology* 11:148-154.

Vire, E., C. Brenner, R. Deplus, L. Blanchon, M. Fraga, C. Didelot, L. Morey, A. Van Eynde, D. Bernard, J. M. Vanderwinden, E. M. Bollen, M. Esteller, L. Di Croce, Y. de Launoit, and F. Fuks, 2006. The polycomb group protein EZH2 directly controls DNA methylation. *Nature* 439:871-874.

Wei, L., G. Vahedi, H. Sun, W. T. Watford, H. Takatori, H. L. Ramos, H. Takahashi, J. Liang, G. Gutierrez-Cruz, C. Zang, W. Peng, J. J. O'Shea, and Y. Kanno, 2010. Discrete roles of STAT6 and STAT4 transcription factors in tuning epigenetic modifications and transcription during T helper cell differentiation. *Immunity* 32:840-

851.

Willermain, F., J. T. Rosenbaum, B. Bodaghi, H. L. Rosenzweig, S. Childers, T. Behrend, G. Wildner, and A. D. Dick, 2012. Interplay between innate and adaptive immunity in the development of non-infectious uveitis. *Progress in Retinal and Eye Research* 31:182-94.

Wurster, A. L., T. Takanaka, and M. J. Grusby, 2000. The biology of Stat4 and Stat6. *Oncogene* 19:2577-2584.

Xu, Z., E. J. Pone, A. Al-Qahtani, S. Park, H. Zan, and P. Casali, 2007. Regulation of *aicda* expression and AID activity: relevance to somatic hypermutation and class switch DNA recombination. *Critical Reviews in Immunology* 27:367-397.

Yadav, A., A. Olaru, M. Saltis, A. Setren, J. Cerny, and F. Livak, 2005. Identification of a ubiquitously active promoter of the murine activation-induced cytidine deaminase (AICDA) gene. 43:529-541.

Yoder, J. A., N. S. Soman, G. L. Verdine, and T. H. Bestor, 1997. New 5' regions of the murine and human genes for DNA (cytosine-5)-methyltransferase. *The Journal of Biological Chemistry* 271:31092-31097.

Yoshikawa, K., I. M. Okazaki, T. Eto, K. Kinoshita, M. Muramatsu, H. Nagaoka, and T. Honjo, 2002. AID enzyme-induced hypermutation in an actively transcribed gene in fibroblasts. *Science* 296:2033-2036.

Zhu, J. K, 2009. Active demethylation mediated by DNA glycosylases. *Annual Review of Genetics* 43:143-166.