Free Radical Biology and Medicine 129 (2018) 611

Contents lists available at ScienceDirect

Free Radical Biology and Medicine

journal homepage: www.elsevier.com/locate/freeradbiomed

Erratum

Erratum to "Manganese porphyrin redox state in endothelial cells: Resonance Raman studies and implications for antioxidant protection towards peroxynitrite" [Free Radic. Biol. Med. 126 (2018) 379–392]☆

Sebastián Carballal[a,b,1], Valeria Valez[a,b,1], Damián Alvarez-Paggi[d,2], Artak Tovmasyan[e], Ines Batinic-Haberle[e], Gerardo Ferrer-Sueta[b,c], Daniel H. Murgida[d], Rafael Radi[a,b,*]

[a] Departamento de Bioquímica, Facultad de Medicina, Universidad de la República, Montevideo, Uruguay
[b] Center for Free Radical and Biomedical Research, Universidad de la República, Montevideo, Uruguay
[c] Laboratorio de Fisicoquímica Biológica, Facultad de Ciencias, Universidad de la República, Montevideo, Uruguay
[d] Departamento de Química Inorgánica, Analítica y Química Física and INQUIMAE-CONICET, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Ciudad Universitaria, Pab. 2, piso 1, C1428EHA Buenos Aires, Argentina
[e] Department of Radiation Oncology, Duke University Medical Center, Durham, NC 27710, USA

The publisher regrets in Fig. 4, the legend does not correspond to that Figure, as it was incorrectly duplicated from the legend of Fig. 2 during the production process.



Fig. 4. RR spectra of MnP incubated with BAEC. Confluent BAEC were incubated with Mn(III)TnHex-2-PyP^{5+} (5 μM) for 2 h. After treatment, cells were extensively washed with PBS, repeating three cycles of cell centrifugation at 1000 *g* for 5 min, pellet was resuspended in PBS and RR spectra were acquired. (a) Control: BAEC in PBS; and (b) preincubated with Mn(III)TnHex-2-PyP^{5+}. Measurements were performed at 457.9 nm excitation, 5 mW laser power at sample and accumulation times of 480 s (8 × 60 s).

The correct legend of Fig. 4 is as follows.
The publisher would like to apologise for any inconvenience caused.

DOI of original article: https://doi.org/10.1016/j.freeradbiomed.2018.08.023

☆ The term peroxynitrite is used to refer to the sum of peroxynitrite anion (ONOO−) and peroxynitrous acid (ONOOH, pKa ~ 6.8). IUPAC recommended names are oxoperoxonitrate (1−) and hydrogen oxoperoxonitrate, respectively.

* Corresponding author at: Departamento de Bioquímica, Facultad de Medicina, Avda. Gral. Flores 2125, 11800 Montevideo, Uruguay.
[1] These authors contributed equally.
[2] Present Address: Instituto de Investigaciones Bioquímicas de Buenos Aires, FIL-CONICET.
E-mail address: rradi@fmed.edu.uy (R. Radi).

https://doi.org/10.1016/j.freeradbiomed.2018.10.001

Available online 11 October 2018