

Published in final edited form as:

*Biochem Biophys Res Commun.* 2010 July 30; 398(3): 585–587. doi:10.1016/j.bbrc.2010.06.125.

## IDH1 mutation identified in human melanoma

Giselle Y. Lopez<sup>a</sup>, Zachary J. Reitman<sup>a</sup>, David Solomon<sup>b</sup>, Todd Waldman<sup>b</sup>, Darell D. Bigner<sup>a</sup>, Roger E. McLendon<sup>a</sup>, Yardena Samuels<sup>c</sup>, and Hai Yan<sup>a,\*</sup>

Giselle Y. Lopez: giselle.lopez@duke.edu; Zachary J. Reitman: zachary.reitman@duke.edu; David Solomon: das67@georgetown.edu; Todd Waldman: waldmant@georgetown.edu; Darell D. Bigner: bigne001@mc.duke.edu; Roger E. McLendon: roger.mclendon@duke.edu; Yardena Samuels: samuelsy@mail.nih.gov; Hai Yan: yan00002@mc.duke.edu

<sup>a</sup> Preston Robert Tisch Brain Tumor Center, The Pediatric Brain Tumor Foundation, and The Department of Pathology, Duke University Medical Center, DUMC 3156, Durham, NC 27710, USA

<sup>b</sup> Department of Oncology, Lombardi Comprehensive Cancer Center, Georgetown University School of Medicine, Washington, District of Columbia 20057, USA

<sup>c</sup> Cancer Genetics Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892-8000, USA

### Abstract

*Isocitrate dehydrogenase 1 (IDH1)* and *isocitrate dehydrogenase 2 (IDH2)* are enzymes which convert isocitrate to  $\alpha$ -ketoglutarate while reducing nicotinamide adenine dinucleotide phosphate (NADP<sup>+</sup> to NADPH). *IDH1/2* were recently identified as mutated in a large percentage of progressive gliomas. These mutations occur at *IDH1*<sup>R132</sup> or the homologous *IDH2*<sup>R172</sup>.

Melanomas share some genetic features with *IDH1/2*-mutated gliomas, such as frequent *TP53* mutation. We sought to test whether melanoma is associated with *IDH1/2* mutations. 78 human melanoma samples were analyzed for *IDH1*<sup>R132</sup> and *IDH2*<sup>R172</sup> mutation status. A somatic, heterozygous *IDH1* c.C394T (p.R132C) mutation was identified in one human melanoma metastasis to the lung. Having identified this mutation in one metastasis, we sought to test the hypothesis that certain selective pressures in the brain environment may specifically favor the cell growth or survival of tumor cells with mutations in *IDH1/2*, regardless of primary tumor site. To address this, we analyzed *IDH1*<sup>R132</sup> and *IDH2*<sup>R172</sup> mutation status 53 metastatic brain tumors, including 9 melanoma metastases. Results revealed no mutations in any samples. This lack of mutations would suggest that mutations in *IDH1*<sup>R132</sup> or *IDH2*<sup>R172</sup> may be necessary for the formation of tumors in a cell-lineage dependent manner, with a particularly strong selective pressure for mutations in progressive gliomas; this also suggests the lack of a particular selective pressure for growth in brain tissue in general. Studies on the cell-lineages of tumors with *IDH1/2* mutations may help clarify the role of these mutations in the development of brain tumors.

### Keywords

isocitrate dehydrogenase 1; isocitrate dehydrogenase 2; melanoma; brain tumors metastases

\*Corresponding author: Hai Yan, DUMC3156, 199B-MSRBI, Research Drive Durham, NC, 27710, USA, (919) 668-7850 (office), (919) 684-8756 (fax), Yan00002@mc.duke.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## Introduction

*Isocitrate dehydrogenase 1 (IDH1)* and *isocitrate dehydrogenase 2 (IDH2)*, which convert isocitrate to  $\alpha$ -ketoglutarate while reducing nicotinamide adenine dinucleotide phosphate (NADP<sup>+</sup> to NADPH), were identified as mutated in a large percentage of progressive gliomas and acute myeloid leukemias [1,2,3,4,5]. These mutations occur at the R132 residue in *IDH1* or the homologous R172 residue in *IDH2*. Melanomas share some genetic features with *IDH1/2*-mutated gliomas, such as frequent *TP53* mutation [6]. Previously, no *IDH1*<sup>R132</sup> mutations were found in a group of 23 melanomas [7], and to our knowledge, melanoma has not been analyzed for *IDH2*<sup>R172</sup> mutations. We sought to test whether melanoma is associated with *IDH1/2* mutations by analyzing a larger group of samples. In addition, given the finding of high *IDH1/2* mutation frequency (80%) in progressive gliomas, we sought to test the hypothesis that certain selective pressures in the brain environment may favor cell growth/survival of tumor cells with mutations in *IDH1/2*. Therefore, to address whether mutation of *IDH1/2* might be required for the development of brain metastases from non-primary central nervous system (CNS) tumors, we sequenced *IDH1*<sup>R132</sup> and *IDH2*<sup>R172</sup> in 53 metastatic brain tumors, including melanomas.

## Materials and Methods

*IDH1/2* gene mutation status was first analyzed in a panel of cell lines derived from human non-CNS metastatic melanoma tumor resections, paired with pheresis-collected peripheral blood mononuclear cells from 78 patients enrolled in Institutional Review Board-approved clinical trials at the Surgery Branch of the National Cancer Institute at the United States National Institutes of Health. Pathology-confirmed melanoma cell lines were derived from mechanically or enzymatically dispersed tumor cells, was subsequently analyzed in a panel of 53 tumor resections from metastases to the brain. Genomic DNA was isolated from frozen tumor tissue samples obtained from the Tissue Bank at the Preston Robert Tisch Brain Tumor Center at Duke University. In both cases, genomic DNA was isolated using the DNeasy Blood & Tissue kit (Qiagen, Valencia, CA, USA). PCR and sequencing primers were designed using Primer 3 ([http://www-genome.wi.mit.edu/cgi-bin/primer/primer3\\_www.cgi](http://www-genome.wi.mit.edu/cgi-bin/primer/primer3_www.cgi)) and synthesized by Invitrogen (Carlsbad, CA, USA) and Integrated DNA Technologies (Coralville, IA, USA). PCR primers were designed to amplify the selected *IDH1* and *IDH2* exons. PCR products were 300–500 bp in length. PCR primers were designed to amplify the region surrounding the commonly mutated codons in both *IDH1* and *IDH2*. In one melanoma sample, all coding exons of *TP53*, *CDKN2A*, and *CDKN2B*, as well as hotspot exons of *BRAF* and *NRas*, were sequenced using these methods.

## Results and Discussion

Of 78 human melanoma samples analyzed for *IDH1*<sup>R132</sup> and *IDH2*<sup>R172</sup> mutation status, a somatic, heterozygous *IDH1* c.C394T (p.R132C) mutation was identified in tissue derived from a melanoma lung metastasis from a 53 year old female (Figure 1). No *IDH2*<sup>R172</sup> mutations were detected in any of these samples. In the *IDH1*-mutated sample, we sequenced regions of *BRAF*, *NRas*, *TP53*, and *CDKN2A/CDKN2B* to identify any changes in genes that are commonly altered in melanoma. We found a *BRAF* c.T1799A (p.V600E) mutation, while *NRas*, *TP53*, and *CDKN2A/CDKN2B* were unaltered in this sample.

*IDH1/2* mutations are frequent in progressive gliomas, but are very rare in other cancers besides acute myelogenous leukemia. To date, mutations in *IDH1* have been identified only in one B-cell acute lymphoblastic leukemia [8], two prostate cancers [8], and one colorectal cancer [9]. Noting the presence of mutations in this small fraction of other cancers, we

postulated two hypotheses. In the first hypothesis, mutations in *IDH1/2* may be cell-lineage dependent, and thus only occur in the cell-of-origin for a specific subset of tumors, including progressive gliomas. In the second hypothesis, a tissue-dependent selective pressure for growth in brain tissue may select for frequent mutations in *IDH1/2*. If the second hypothesis is correct, we should be able to identify the presence of relatively frequent mutations in *IDH1/2* in non-primary CNS metastases to the brain. Therefore, we analyzed *IDH1<sup>R132</sup>* and *IDH2<sup>R172</sup>* mutation status in a set of 9 melanoma metastases to the brain and an additional panel of 44 metastases of other tumor types to the brain, including lung, breast, renal, uterine, ovarian, esophageal, and urothelial cancer metastases. Results revealed no mutations regardless of tumor type, consistent with a previous small-scale study looking at colon cancer metastases to the brain [10].

Here, we present the first identification of a mutation in *IDH1* or *IDH2* in melanoma. These results have important clinical implications regarding the role of mutations in *IDH1/2* in the development of tumors. Mutations in *IDH1* have now been identified in gliomas, leukemias, prostate cancer, colorectal cancers, and melanomas, suggesting the possibility of a commonly altered pathway that may prove advantageous to the formation of tumors in all these cell types. The lack of mutations in a panel of non-primary CNS metastases to the brain would suggest that mutations in *IDH1<sup>R132</sup>* or *IDH2<sup>R712</sup>* are not necessarily required specifically for growth in brain tissue. The low frequency of mutations in *IDH1<sup>R132</sup>* and *IDH2<sup>R172</sup>* in primary glioblastomas, which grow extremely aggressively in brain tissue, further supports this. This leads to the conclusion that mutations in these genes may be necessary for formation of tumors in a cell-lineage dependent manner, with a particularly strong selective pressure for mutations in progressive gliomas and acute myelogenous leukemias. Studies focusing on the cell-lineages from which tumors with *IDH1* or *IDH2* mutations develop may help to elucidate the role of these mutations in cancer pathogenesis.

#### Research Highlights

- An *IDH1<sup>R132C</sup>* mutation was identified in a human melanoma metastasis to the lung.
- *IDH1/2* mutations were not identified in non-CNS tumor metastases to the brain.

## Acknowledgments

We thank Kristin E. Yates for technical assistance. This work is supported by the Intramural Research Programs of the National Human Genome Research Institute and the National Cancer Institute, National Institutes of Health, USA.


## Abbreviations

<b>CNS</b>	central nervous system
<b>IDH1</b>	isocitrate dehydrogenase 1
<b>IDH2</b>	isocitrate dehydrogenase 2

## References

1. Yan H, Parsons DW, Jin G, McLendon R, Rasheed BA, Yuan W, Kos I, Batinic-Haberle I, Jones S, Riggins G, Friedman HS, Friedman AH, Reardon DA, Herndon JE, Kinzler KW, Velculescu VE, Vogelstein B, Bigner D. *IDH1* and *IDH2* Mutations in Gliomas. *N Engl J Med* 2009;360:765–773. [PubMed: 19228619]

2. Gross S, Cairns RA, Minden MD, Driggers EM, Bittinger MA, Jang HG, Sasaki M, Jin S, Schenkein DP, Su SM, Dang L, Fantin VR, Mak TW. Cancer-associated metabolite 2-hydroxyglutarate accumulates in acute myelogenous leukemia with isocitrate dehydrogenase 1 and 2 mutations. *J Exp Med* 2010;207:339–344. [PubMed: 20142433]
3. Green A, Beer P. Somatic mutations of IDH1 and IDH2 in the leukemic transformation of myeloproliferative neoplasms. *N Engl J Med* 2010;362:369–370. [PubMed: 20107228]
4. Ward PS, Patel J, Wise DR, Abdel-Wahab O, Bennett BD, Collier HA, Cross JR, Fantin VR, Hedvat CV, Perl AE, Rabinowitz JD, Carroll M, Su SM, Sharp KA, Levine RL, Thompson CB. The common feature of leukemia-associated IDH1 and IDH2 mutations is a neomorphic enzyme activity converting alpha-ketoglutarate to 2-hydroxyglutarate. *Cancer Cell* 2010;17:225–234. [PubMed: 20171147]
5. Mardis ER, Ding L, Dooling DJ, Larson DE, McLellan MD, Chen K, Koboldt DC, Fulton RS, Delehaunty KD, McGrath SD, Fulton LA, Locke DP, Magrini VJ, Abbott RM, Vickery TL, Reed JS, Robinson JS, Wylie T, Smith SM, Carmichael L, Eldred JM, Harris CC, Walker J, Peck JB, Du F, Dukes AF, Sanderson GE, Brummett AM, Clark E, McMichael JF, Meyer RJ, Schindler JK, Pohl CS, Wallis JW, Shi X, Lin L, Schmidt H, Tang Y, Haipek C, Wiechert ME, Ivy JV, Kalicki J, Elliott G, Ries RE, Payton JE, Westervelt P, Tomasson MH, Watson MA, Baty J, Heath S, Shannon WD, Nagarajan R, Link DC, Walter MJ, Graubert TA, DiPersio JF, Wilson RK, Ley TJ. Recurring mutations found by sequencing an acute myeloid leukemia genome. *N Engl J Med* 2009;361:1058–1066. [PubMed: 19657110]
6. Ibrahim N, Haluska FG. Molecular pathogenesis of cutaneous melanocytic neoplasms. *Annu Rev Pathol* 2009;4:551–579. [PubMed: 19400696]
7. Bleeker FE, Lamba S, Leenstra S, Troost D, Hulsebos T, Vandertop WP, Frattini M, Molinari F, Knowles M, Cerrato A, Rodolfo M, Scarpa A, Felicioni L, Buttitta F, Malatesta S, Marchetti A, Bardelli A. IDH1 mutations at residue p.R132 (IDH1(R132)) occur frequently in high-grade gliomas but not in other solid tumors. *Hum Mutat* 2009;30:7–11. [PubMed: 19117336]
8. Kang MR, Kim MS, Oh JE, Kim YR, Song SY, Seo SI, Lee JY, Yoo NJ, Lee SH. Mutational analysis of IDH1 codon 132 in glioblastomas and other common cancers. *Int J Cancer* 2009;125:353–355. [PubMed: 19378339]
9. Sjoblom T, Jones S, Wood LD, Parsons DW, Lin J, Barber TD, Mandelker D, Leary RJ, Ptak J, Silliman N, Szabo S, Buckhaults P, Farrell C, Meeh P, Markowitz SD, Willis J, Dawson D, Willson JK, Gazdar AF, Hartigan J, Wu L, Liu C, Parmigiani G, Park BH, Bachman KE, Papadopoulos N, Vogelstein B, Kinzler KW, Velculescu VE. The consensus coding sequences of human breast and colorectal cancers. *Science* 2006;314:268–274. [PubMed: 16959974]
10. Holdhoff M, Parsons DW, Diaz LA Jr. Mutations of IDH1 and IDH2 are not detected in brain metastases of colorectal cancer. *J Neurooncol* 2009;94:297. [PubMed: 19350208]



**Figure 1.** Sequencing of melanoma metastases identified IDH1 R132C mutation in a melanoma cell line derived from a melanoma lung metastasis.