



Review

Review of contaminant levels and effects in shorebirds: Knowledge gaps and conservation priorities

Yanju Ma^{1,2}, Chi-Yeung Choi, Alex Thomas, Luke Gibson*

School of Environmental Science & Engineering, Southern University of Science and Technology, Shenzhen, Guangdong 518055, China



ARTICLE INFO

Edited by: Professor Bing Yan

Keywords:

Waders
Flyway
Pollutants
POPs
Heavy metals
Conservation

ABSTRACT

Environmental pollution has emerged as a major threat to bird populations. Many shorebird populations are declining, although contamination has been documented in some shorebirds, evidence of negative impacts is sparse and this important topic remains understudied. To guide future research and develop effective conservation strategies, we carried out a comprehensive review of environmental pollutants and their consequences on shorebirds. In total, we found 93 relevant articles which examined pollutant contamination in ~37% (79 of 215) of all shorebird species, mostly from the Charadriidae and Scolopacidae families. Studies were geographically biased: the majority were conducted in American flyways, while only 1 was found from Australasia and few were conducted in Asian flyways. The main geographic gap for research includes East Africa, South Asia and Siberian Arctic. The most well-documented pollutants included mercury (Hg, 37 studies), cadmium (33), and lead (Pb, 28); less well studied pollutants were barium (1), calcium (1), strontium (1), dicofols (1), and other newly emerging contaminants, such as plastic debris/microplastics (4) and antibiotics resistance (2). Several pollutants have caused considerable concerns in shorebirds, including embryotoxicity caused by PCBs at non-optimum temperature (laboratory experiments); reduced reproduction performance linked to maternal Hg and paternal Pb (field evidence); and reduced refueling and flight performance related to oil contamination (both field and laboratory evidence). Our results confirm that an in-depth understanding of the local, regional and global factors that influence population trends of shorebirds in light of increasing pollution threats is essential for accurate and effective management and conservation strategies.

1. Introduction

Shorebirds (Order Charadriiformes) form an essential component in wetland ecosystems worldwide. Many populations have recently experienced steep declines, possibly due to anthropogenic impacts such as climate change (Van Gils et al., 2016) and habitat loss/modification (Ma et al., 2014). As one of the largest avian groups, shorebirds could be vulnerable to environmental pollution (Melville et al., 2016; Tang et al., 2015), likely due to their life-history traits (e.g., foraging behavior and migratory ecology) for the whole life cycle. During non-breeding seasons, they may stop or overwinter at various wetlands, including estuarine mudflats, coastal tidal flats, and freshwater ecosystems (e.g., rivers and lakes). Some of these habitats are close to human settlements and receive constant inputs of anthropogenic toxins via industrial and

agricultural activities (Chapagain et al., 2009; C. Li et al., 2022; Zhang et al., 2009). Even in remote northern regions, which serve as breeding grounds for many migratory shorebird species, volatile or semi-volatile persistent organic pollutants (POPs, UNEP, 2021) and mercury (Hg) have been recorded at concerning concentrations via long-range atmospheric transportation (Dietz et al., 2013; Wania and Mackay, 1993).

Environmental pollutants have been studied through investigations of various endpoints/consequences in wild free-living and captive avian subjects. In general, their detrimental impacts on birds have been widely publicized, with impacts including depressed immune system (e.g., Kenow et al., 2007; Spalding et al., 2000; Vallverdú-Coll et al., 2019), embryotoxicity (King et al., 2021), decreased reproductive success (Fernie et al., 2009), altered parental care behavior (Hartman et al., 2019; Pedersen and Saether, 1999), behavioral deficits (e.g.,

* Corresponding author.

E-mail address: biodiversity@sustech.edu.cn (L. Gibson).¹ Current affiliation: Key Laboratory of Ecology of Rare and Endangered Species and Environmental Protection, Ministry of Education, Guangxi Normal University, Guilin, China.² Current affiliation: Guangxi Key Laboratory of Rare and Endangered Animal Ecology, College of Life Science, Guangxi Normal University, Guilin, China.

locomotion, food-begging, and individual recognition, Burger and Gochfeld, 2010), impaired migratory performance (Flahr et al., 2015; Ma et al., 2018b; Maggini et al., 2017b), and even reduced survival (Grade et al., 2018; Ma et al., 2018a). Environmental pollutants could pose a great threat to the health of shorebirds throughout their dynamic life cycle, leading to further population declines. However, contamination of natural or synthetic pollutants and associated impacts on shorebirds have not been fully reviewed. Here, we conducted an exhaustive literature review to: (1) summarize the type of pollutants and shorebird species that have been studied; (2) assess the impacts of these key contaminants on shorebird species and populations; and (3) identify research priorities that can help ensure the effective conservation of this vulnerable taxonomic group.

2. Material and methods

To construct a database of published studies on environmental contaminants in shorebirds, we retrieved related papers on the Web of Science (WoS) on May 18, 2022, by using the following search item: topic = ("pollut*" or "heavy metal" or "trace metal" or Pb or lead or Cd or cadmium or Zn or zinc or Mn or manganese or Cu or copper or Tl or thallium or Cr or chromium or Co or cobalt or Ni or nickel or arsenic or Hg or mercury or "POPs" or PCBs or PAHs or DDTs or microplastics or plastic debris or antibiotics resistance) AND topic = shorebird or wader). Although we attempted to systematically compile all relevant studies within the scope of the review, we acknowledged that there may exist other studies that we were unable to locate.

3. Results and discussion

Our search produced 578 results. After manually reviewing these studies, we excluded irrelevant studies related to other wildlife or humans and eventually included 93 relevant studies in this review. The earliest research on environmental pollutants and impacts on shorebirds date back to at least the 1970 s – there was one study documented that several waterbird species using rice fields were poisoned by intaking aldrin-treated seeds or through aldrin-related contamination accumulated in the food chain along Texas Gulf Coast (Flickinger and King, 1972). Two direct poisoning events related to shorebirds were documented in the 1980 s and the 1990 s. Specifically, Bull et al. (1983) reported a poisoning event that caused death and sickness in about 2400 birds, including Dunlin *Calidris alpina* and Redshank *Tringa tetanus*, as a result of regular lead discharge on the Mersey Estuary, UK. Dunlins and Killdeers *Charadrius vociferous* in the USA were poisoned by toxins related to rodent control (Warnock and Schwarzbach, 1995), indicating shorebird species likely to be susceptible as non-target species for pest control. The number of studies increased from 5 studies in the 1980 s to 17 cases in the 1990 s and 18 cases in the 2000 s. The study cases jumped to 44 in the 2010 s, with those published in the 2010 s twice the number of those published in the 2000 s. Studied contaminants in shorebirds were mainly classified into two major classes: well-known POPs (organics) and trace elements (elements), but others involving recent research related to oil spills, antibiotics and microplastics/ plastic debris were also covered in recent years (See Fig. 1). The summary of studied pollutants, shorebird species, conservation status, population trend, life stages, critical sites along flyways as well as other key information are discussed below and Table S1.

3.1. Studied pollutants and species

The contaminant abbreviations and their frequency (indicated by number of studies) are summarized in Table 1.

Overall, thirty-seven pollutants have been studied at least once, including both essential elements such as Mn, Cu, Zn and Se as well as toxic elements such as Hg and Pb. For trace elements, the most studied element is Hg followed by Cd, Pb and Se, while the least studied are on




Fig. 1. Trends in the number of published articles on environmental pollution in shorebirds between the 1970 s and 2022 on a decadal basis.

Table 1

Contaminant abbreviations used and the number of studies performed for each contaminant. E: trace elements, O: organic pollutants.

Pollutant	Abbreviations	# of cases	Category
Mercury	Hg	37	E
Cadmium	Cd	33	E
Lead	Pb	28	E
Selenium	Se	25	E
Copper	Cu	20	E
Chromium	Cr	18	E
Zinc	Zn	18	E
Arsenic	As	17	E
Polychlorinated biphenyls	PCBs	17	O
Manganese	Mn	14	E
Dichlorodiphenyltrichloroethane and its metabolites	DDTs	14	O
Nickel	Ni	12	E
Oil contamination related (Polycyclic aromatic hydrocarbon included)	NA (PAH included)	12	Other
Cobalt	Co	11	E
Iron	Fe	10	E
Hexachlorocyclohexane	HCHs	9	O
Hexachlorobenzene	HCB	8	O
Vanadium	V	6	E
Chlordanes	CHLs	6	E
Silver	Ag	5	E
Hexachlorocyclopentadienes	HCCPDs	5	O
Molybdenum	Mo	4	E
Polybrominated diphenyl ethers	PBDEs	4	O
Heptachlor epoxide	NA	4	O
Plastic debris/micro plastic	NA	4	Other
Beryllium	Be	3	E
Magnesium	Mg	3	E
Aluminum	Al	2	E
Boron	B	2	E
Antimony	Sb	2	E
Thallium	Tl	2	E
Polychlorinated dibenzo-p-dioxins and dibenzofurans	PCDD/Fs	2	O
Antibiotics resistance	ABR	2	Other
Barium	Ba	1	E
Calcium	Ca	1	E
Strontium	Sr	1	E
Dicofols	NA	1	O

Ba, Ca, and Sr. In terms of industrial products, the polyhalogenated compounds PCBs and oil contamination (e.g., oil spills and PAH) are the most studied, followed by DDTs, HCHs, HCB and CHLs, while PCDD/Fs and dicofols are being the least studied. Some studies focus on only one

pollutant, including 12 cases for Hg (e.g., Ackerman et al., 2007), 3 for Cd (Ferns and Anderson, 1994; McFarland et al., 2002; Stock et al., 2007), and 2 for Pb (Bull et al., 1983; Ferns and Anderson, 1997). There are four studies (Braune and Noble, 2009; Lindberg et al., 1985; Pratte et al., 2020; Saalfeld et al., 2016) examined both elemental and organic pollution. In addition, to gain enough sample weight, carcass, muscle, adipose tissue and egg are generally used for organic related analysis, while feather and internal tissues (e.g., liver and kidney) are more often served as selected tissues/organs for elemental analysis (Table S1). For biomonitoring in the future, it is encouraged to screen multiple pollutants in the same samples to gain as much baseline information as possible, since such information is limited in shorebird groups.

Of the roughly 215 shorebird species known worldwide (Colwell, 2010), only 79 species have been subjected to pollutant-related investigations. Among the studied species, Great Knot *Calidris tenuirostris* is classified as Endangered (EN), while other 12 species were listed as Near Threatened (NT) on the IUCN Red List, mostly with decreasing population trends (Table S1, IUCN, 2022). The studied species mainly belong to the Charadriidae and Scolopacidae families, while fewer belong to Recurvirostridae (4 species) and Haematopodidae (5 oystercatcher species). Still, 34 species were studied only once for the determination of pollutant exposure. The most studied species were Dunlin (20 studies), followed by Semipalmated Sandpiper *Calidris pusilla* (17 studies), Sanderling *Calidris alba* (15 studies), Kentish Plover *Charadrius alexandrinus* (13 studies), Western Sandpiper *Calidris mauri* (12 studies), and Red Knot *Calidris alba* (11 studies). Specifically, the Western Sandpiper, one of the most common, long-distance migrating shorebirds in the Western Hemisphere, has been used as a model species for laboratory studies related to oil spill effects (Bursian et al., 2017; Maggini et al., 2017a, b; Perez et al., 2017). In field studies, Oystercatchers (Black Oystercatcher *Haematopus bachmani* and Eurasian Oystercatcher *Haematopus ostralegus*) are often used as studied subjects (Goede and Wolterbeek, 1994; Schwemmer et al., 2015; Stock et al., 2007; Vermeer and Castilla, 1991) for investigating the residue patterns of Cd and Se in European and South American continents. The Kentish Plover, a species with a wide distribution within the Palearctic realm (Argüelles-Ticó et al., 2016), has also been studied to understand how the molt patterns could help in the biomonitoring of trace elements (Picone et al., 2019).

Notably, several studies examined organochlorines and heavy metals (e.g., Hg and Cd) in both shorebird and waterfowl species (order Anseriformes, ducks, geese and swans) to assess the potential risk of consumption of waterbirds from a public health perspective (Hiller and Barclay, 2011; Tsuji et al., 2008; Vermeer and Castilla, 1991). For example, in the Western James Bay region, northern Ontario, Canada, the levels of PCBs and DDTs in Hudsonian Godwit *Limosa haemastica* and Marbled Godwit *Limosa fedoa* were within acceptable range, but Hudsonian Whimbrels *Numenius phaeopus* sampled from Chile and American Woodcock *Scolopax minor* harvested in Connecticut of the USA showed high concentrations of Cd and Pb residues, respectively, that may cause public health concerns. Besides DDTs, most shorebirds accumulated greater concentrations of PCBs (240–12000 ng/g, fat weight) than the concentrations of PCBs in other waterfowls (180–830 ng/g, fat weight) in Asia (Kunisue et al., 2003). Except for Black-necked Stilts *Himantopus mexicanus* from the historical mining site of San Francisco Bay contained muscle total Hg concentrations of 1.30–8.76 mg/kg, dry weight (Eagles-Smith et al., 2009), the total Hg concentrations were generally less than 1 mg/kg in both wetland associated shorebirds and waterfowls (Burger et al., 2014; Fernández et al., 2019; Lucia et al., 2014; Pandiyan et al., 2020; Zamani-Ahmadmohammoodi et al., 2010).

3.2. Flyways and important sites

The 93 relevant studies span over 25 countries, with most research conducted in the Atlantic Americas (AAF), Pacific Americas (PAF), and Mississippi Americas (MAF) Flyways along the North American continent (the USA and Canada), followed by the East Atlantic Flyway (EAF)

in Europe. In comparison, the Asia-Pacific region is much less studied; only 14 studies were carried out along the East Asian-Australasian Flyway (EAAF), the most threatened but least understood flyway (Yong et al., 2015), while other Asian related flyways were also understudied (Fig. 2; Table S1). Despite being a major shorebird wintering region, there was only one pollutant related study conducted in Australia and it was on antibiotics resistance (Smith et al., 2022).

About 25 studies were conducted on breeding grounds including Alaska (e.g., Andres, 1999; Murphy et al., 1997; Saalfeld et al., 2016), northern Canada (Hargreaves et al., 2011, 2010), the Wadden Sea in northern Europe (Goede and Wolterbeek, 1994; Lindberg et al., 1985) and the Cavallino-Treporti Peninsula of Southern Europe (Picone et al., 2019). Another 26 studies were conducted in wintering sites including Venezuela (Pratte et al., 2020), Suriname (Burger et al., 2018), Pampas of South America (Scherer et al., 2015), Chile (Navedo et al., 2021), Mexico (Fernández et al., 2019) and China (Su et al., 2020). Finally, a further 25 studies were conducted in stopover/staging sites such as the Gulf of Mexico (Bianchini and Morrissey, 2018a), the Baltic Sea (e.g., Blomqvist et al., 1987), and the Yellow Sea (e.g., Zheng et al., 2018). For some studies, we could not easily define the location due to unclear sampling information. A few other studies are laboratory-based research (e.g., Bianchini and Morrissey, 2018b; Lunny et al., 2020; Maggini et al., 2017a).

Several North American-based studies have investigated the profiles of various long-distance transport contaminants in eggs, blood, and feathers, and their correlations with reproduction success in Alaska and Nunavut. Specifically, elevated ecological risks to Hg (Perkins et al., 2016) and chlorinated contaminants such as PCBs and DDTs (Braune and Noble, 2009; Pratte et al., 2020) have been recorded. Particularly, in the Canadian Arctic, shorebirds were exposed to elevated Hg in blood and eggs, while the body condition of Red Phalarope *Phalaropus fulicarius* declined in relation to As (Hargreaves et al., 2011). In Alaska, elevated Sr in eggs had been observed in several species, especially the Black Oystercatcher (Saalfeld et al., 2016). Strontium has a similar structure to Ca, potentially affecting eggshell thickness, bone growth, and hatching success (Mora et al., 2007). Yet, the potential sources and impacts of Sr on Alaskan shorebirds remain unclear.

Research conducted in Europe has mainly focused on the Wadden Sea: the transportation of Pb, Hg, and Cd along food webs of Black-tailed Godwit *Limosa limosa* (Roodbergen et al., 2008); Se in eggs and parental blood, and its role in antioxidation in Eurasian Oystercatcher (Goede, 1993; Goede and Wolterbeek, 1994); Hg accumulation and potential drivers among shorebird chicks (Tavares et al., 2009, 2004); and the profiles of PCBs, DDTs, HCHs, and PBDEs in Eurasian Oystercatcher




Fig. 2. A geographic summary of shorebird pollutant studies identified in this review. The locations represent where shorebirds were captured and sampled. For those studies that did not report specific GPS coordinates, the coordinates of the general location described were used. Map layers were provided by Esri, HERE, Garmin, © OpenStreetMap contributors, and the GIS user community.

(Lambeck et al., 1991; Schwemmer et al., 2015). In the Iberian Peninsula, Pb, Zn, and Cu were found in feathers of Kentish Plover at concentrations above the established thresholds for toxicological effects in some populations (Vidal et al., 2021). Notably, there was only one study involving research conducted in the remote Asian breeding grounds in Russia and Japan, which suggested that PCBs and DDTs were the most common organochlorine pollutants recorded (Kunisue et al., 2003).

Exposures of various environmental pollutants were commonly detected at remote breeding grounds. However, it is hard to find sufficient evidence that this contamination has adversely affected the breeding success of resident avifauna. Yet, recent laboratory research indicated that the interaction between climate change and environmental toxins might have pronounced impacts on birds living at northern breeding grounds. By using a controlled egg injection, an incubation study on Killdeer investigated the cumulative effects of exposure to PCBs at suboptimal incubation temperature and found alterations in heartbeat onset, incubation length, post-hatch growth rate, and higher post-hatch mortality during the embryonic and early post-hatch periods (Lunny et al., 2020). This is particularly concerning since pollution is not the only ecological stressor for the Arctic breeding shorebirds under climate change.

Research on contamination during the non-breeding stage has been conducted at several critical stopover sites including Delaware Bay and San Francisco Bay in North America; Bohai Bay in East Asia and the Wadden Sea in Europe. At the stopover/staging sites of Delaware Bay in the AAF, a series of studies conducted mainly by Burger and colleagues focused on trace elements (e.g., Hg, Pb, Cd, Co, As and Se) in multiple tissues of migratory Semipalmated Sandpipers, Sanderlings and Red Knots. The research findings included the correlation between body weight and element burden, the ratio between Se and Hg (Burger et al., 2014), annual and interspecific variation (Tsipoura et al., 2017), and accumulation among food webs (Burger et al., 2019b), particularly related to horseshoe crab eggs (Burger et al., 2017), the key dietary component for stopover shorebirds and impacted feeding behavior caused by oil pollution (Burger, 1997). In general, the levels of elements in blood were below any threshold, except Se which was higher than the sublethal effect levels for birds (Burger et al., 2019a).

In South America, studies conducted at wintering sites mostly focused on the exposure levels of trace elements. Shorebirds foraging at beaches near copper mine tailings were recorded with high Cd residues in Chile (Vermeer and Castilla, 1991). Blood samples in Semipalmated Sandpipers wintering in Brazil indicated that Se was a concern and worth further investigation. In Europe, PCBs were investigated in Eurasian Oystercatchers that died from starvation during severe winter weather in the Wadden Sea, but it was doubtful if PCBs contributed to this winter mortality (Lambeck et al., 1991).

3.3. Influencing factors

The identified factors driving interspecific variation of bioaccumulation impacts include: differences in a trophic niche (Lucia et al., 2014); feeding ecology (Levengood et al., 2014), and exposure timing (Ackerman et al., 2007). In the same species, gender and age were commonly investigated as biological factors for variation in trace element concentrations among tissues. In general, males tend to carry a higher body burden of heavy metals than females, while adults accumulate more than juveniles. For instance, Hui (1998) and Stock et al. (2007) both reported that males of Willet *Catoptrophorus semipalmatus* and Eurasian Oystercatcher had higher levels of Cd than females, and male plovers had higher levels of Fe, Hg, and Mn than females. In adult Dunlin, males contained higher concentrations of Pb than females; while adults had significantly higher Pb than juveniles (Ferns and Anderson, 1997). Even in remote breeding grounds, similar patterns were found (Perkins et al., 2016). In contrast, adult female Kentish Plovers inhabiting the Iberian Peninsula showed significantly higher levels of Mn in feathers (Vidal et al., 2021) despite female Kentish Plovers could excrete

pollutants (e.g., 20–40% of a load of POPs) into eggs (Wang et al., 2019).

The understanding of seasonal variation in contamination is essential as many migrants have a large spatial range during their dynamic annual life histories. Pratte et al. (2020) compared contaminant exposure (organochlorines and toxic trace elements) of four Arctic-breeding shorebird species collected during breeding, migration, and wintering seasons in Canada and found the levels of chlorinated pesticides and Hg were consistently the highest on the breeding grounds. Perkins et al. (2016) also pointed out that some breeding individuals in Alaska had sufficiently high concentrations of pollutants which could contribute to potentially adverse effects, while the exposure levels in staging individuals were lower in the same location.

Although shorebirds occupy a relatively low trophic position, accumulation of toxins existed even in species that spent only a few months at wintering or breeding grounds, but bioaccumulation varies between toxins in soil or invertebrates (Ferns and Anderson, 1994; Roodbergen et al., 2008). Hargreaves et al. (2011) showed that Hg, Se, Cd, Cu and Zn bioconcentrated from soil to invertebrates while Hg and Se were observed to be biomagnified from invertebrates to shorebird blood. Thus, shorebird conservationists may put more effort to environmental contaminants that have the ability to bioaccumulate, such as Hg and Se. Other potential factors such as body size and diet composition/preference that were thought to influence bioaccumulation in other avian groups have not been well investigated.

3.4. Case studies - key contaminants

For pollutants i.e., oil pollution, mercury, lead, cadmium, and persistent organochlorines, are of great concern whose impacts have been studied in depth and in a quantitative manner, we summarize the latest research findings below.

3.4.1. Oil contamination

Oil spill events are regarded as an ongoing tragedy (e.g., Bruederle and Hodler, 2019) in many regions globally. Compared to the acute mortality in rafting seabirds, shorebirds exposed to crude oil mainly via large high-profile spills (Deepwater Horizon, Exxon Valdez, and Anitra). Shorebirds are likely facing threats through: (1) external oil on feathers; (2) ingestion through preening oil from oiled feathers; (3) foraging on prey contaminated by oil; (4) disturbance by clean-up activities. Although an increased risk to survival in shorebirds due to oil contamination seems likely, it is difficult to find support that shorebird demography was negatively influenced by oil spill events (Gibson et al., 2017).

To better understand the negative impacts caused by the Deepwater Horizon oil spill, a variety of laboratory studies were conducted to evaluate food intake, and physiological and flight performance using wild-caught Western Sandpipers and Sanderlings. Specifically, oiled feathers caused impaired takeoff flight, potentially increasing predation risk in the wild (Maggini et al., 2017b). In addition, increased flight energy expenditures was observed in wind tunnel facilities, likely caused delays at stopover/staging sites en route to breeding grounds (Maggini et al., 2017b). In field studies, shorebirds were observed to be interrupted by oil spill clean-up personnel and vehicles; as well as to spend more time cleaning their oiled feathers (Burger, 1997; Henkel et al., 2014). When ingested, the toxic constituents from oil spill, such as PAHs, are believed to be linked to delayed departure at a heavily oil polluted staging site in the Gulf of Mexico (Bianchini and Morrissey, 2018a; Bianchini et al., 2021). Besides, when exposed to oil for short periods, individuals were found to have lower body temperatures and reduced body mass (Maggini et al., 2017a).

In seabirds, negative impacts in reproduction from the consumption of petroleum contaminated prey were found, including decreased parental care and reduced weight gain in chicks (Miller et al., 1978). In shorebirds, only one study exists, which investigated the impacts of oil ingestion on chick mass gain of the Black Oystercatcher chicks at Prince

William Sound, Alaska (Andres, 1999). The chicks exhibited a slower growth rate prior to fledgling, but overall fledge success was not reduced. Also, in the same area, the abundance and distribution were compared before and after the Exxon Valdez oil spill (Irons et al., 2000; Murphy et al., 1997). These results showed variable responses among species with the population densities of Black Oystercatchers being negatively affected by the oil spill, particularly in the early years following the oil spill.

3.4.2. Mercury

Shorebirds are thought to be sensitive to Hg due to their foraging preference for coastal wetlands, which have been recognized as a significant source of methylmercury (MeHg), the most toxic form of Hg among marine food webs (Chen et al., 2012; Schäfer et al., 2010). MeHg is a potent neurotoxin and can have detrimental impacts on long-distance migrants due to impaired takeoff and endurance flight performance (Carlson et al., 2014; Ma et al., 2018b), lowering their ability to avoid predators and eventually lowering their survival in the wild (Ma et al., 2018a). Many shorebird species have some of the longest migration distances covered by any wildlife. Hence, Hg contamination could be a severe threat to them, but little is known about the bioaccumulation of Hg and particularly MeHg in shorebirds. More investigation on this topic is urgently needed to protect shorebird health and ensure their conservation.

At breeding grounds, it has been well described that Hg contamination in shorebirds exceeded toxicological thresholds. Picone et al. (2019) found that Hg emerged as a major threat to the conservation of the Kentish Plovers in the Venice Lagoon area since the average concentration in feathers was clearly above the adverse-effect threshold of 5 mg/kg, a level associated with impairment in the reproductive success. Notably, reproductive success was negatively correlated with paternal Hg which was adversely related to egg volume (Hargreaves et al., 2011). There was variation in blood Hg among the Arctic region: blood Hg at Barrow of the Alaskan breeding ground were higher than those breeding in the Eastern Canadian Arctic and the Yukon Arctic (Perkins et al., 2016; Hargreaves et al., 2011). In addition, micro-habitat preference for shorebird individuals seems to play a key role in differential Hg accumulation among waterbird species (Perkins et al., 2016). Furthermore, Eagles-Smith et al. (2009) used radio-tracking to determine habitat preferences in American Avocets *Recurvirostra americana* and Black-necked Stilts. They found that stilts, which tend to use more vegetated areas such as marshes, had higher Hg concentrations in blood.

Although Asia suffers from the highest rates of Hg emission (UN Environment, 2019), research in this region remains scarce. Recently, Su et al. (2020) reported feather Hg concentrations in Charadrius plovers (Kentish and White-faced Plover) *Charadrius dealbatus* right after "White-faced Plover from coastal populations of China were below the toxicological threshold of reproduction impairment. Meanwhile, the population breeding in inland China was found to have elevated feather Hg, indicating that Hg risk may cause concern at an unknown molting site (Su et al., 2020). Thus, it is important to incorporate the knowledge of molt pattern to aid shorebird contamination studies.

In general, Se is thought to be a detoxification agent for Hg toxicity and a ratio of Se: Hg may indicate an interaction between these two toxic elements which may be mutually detoxifying, and the ratio above 21 suggests the potential for the amelioration of Hg toxicity (Koeman et al., 1973). A few studies reported the molar ratio between Se and Hg in shorebirds. Specifically, Long-billed Dowitcher *Limnodromus scolopaceus* had concentrations of Se and Hg (in pooled samples) which were highly correlated in a mean molar ratio of 19:1, similar to that reported in other birds (Hui et al., 2001). Yet, Burger et al. (2014) reported a varied range of Se: Hg molar ratios among tissues, with brain and fat having the highest ratio of 141:1, while breast feathers have the lowest ratio of 31.41:1. There is only one study but no strong correlation between Se and Hg in breeding shorebirds (Hargreaves et al., 2010). Yet, whether and how Se detoxifies the Hg toxicity in shorebirds still needs more

investigation (Burger and Gochfeld, 2021).

3.4.3. Lead

Lead (Pb) as a toxic substance, threatens humans and wildlife (Pokras and Kneeland, 2009). Several avian groups i.e., birds of prey, scavengers, and waterbirds are at high risk to exposure to Pb poisoning from ingestion of ammunition, fishing weights, carcasses or injured animals contaminated by Pb ammunition residues (details see review Eisler, 1988). The Common Loon *Gavia immer* in New Hampshire, the USA is a good example – approximately 48% of collected adult mortalities was due to Pb fishing tackle from 1989 to 2012 (Grade et al., 2018). Furthermore, sublethal effects caused by Pb contamination include immunotoxic toxicity (Vallverdú-Coll et al., 2019), as well as damaged nervous system and kidney and liver function (Mudge, 1983).

Shorebirds are not usually affected by lead poisoning directly. Cases were reported of Pb poisoning in the British estuaries, which was contaminated by a nearby industrial smelter and vehicle exhausts (Bull et al., 1983; Ferns and Anderson, 1997). In the liver of a Dunlin, the Pb concentration was up to 31.0 mg/kg (wet weight), which was considered Pb poisoning in raptors. In fact, shorebirds can be exposed to Pb via atmospheric pollution in remote breeding regions, as well as waste disposal at wetland associate habitats (Haig et al., 2014). Lead accumulation in various tissues (e.g., bone, blood, feather, and even beak) has been widely reported in approximately 25 shorebird species across the Arctic (Hargreaves et al., 2010), Delaware Bay (a key stopover for multiple flyways; Burger et al., 1993, 2014, 2015, 2019a,b; Tsipoura et al., 2017); South America (Burger et al., 2018; Scherer et al., 2015), and the Europe (Blomqvist et al., 1987; Ferns and Anderson, 1997; Lucia et al., 2014, 2012a, b; Picone et al., 2019; Roodbergen et al., 2008; Vidal et al., 2021).

Along the EAAF, Pb is a potential threat to migratory shorebirds based on a series of research conducted in Korea (Kim et al., 2007a,b; Kim and Koo, 2010). Particularly, Pb was found elevated in Red-necked Stints *Calidris ruficollis*, suggesting this contamination should reflect the accumulation in their breeding grounds since this species only spends a short period at Korean stopover; furthermore, Pb concentration did not match the sediment concentrations in Korea (Kim et al., 2010). In addition, bone Pb in Terek Sandpipers *Xenus cinereus*, Great Knots, and Red-necked Stints from the Okgu Mudflat, Korea were all found greater than 10 mg/kg (dry weight), and were even higher than that in birds of prey from the USA and Spain, suggesting chronic exposure along this flyway.

3.4.4. Cadmium

Cadmium (Cd) is a ubiquitous pollutant found in many ecosystems from both natural and anthropogenic sources such as mining activities and industrial processes. For birds, Cd has been associated with kidney toxicity, decreased growth (Eisler, 1985), and altered behavior (Heinz et al., 1983). Similar to Pb, a large spectrum of tissues including kidney, liver, bone, muscle, egg, and blood have been investigated for Cd concentrations among shorebird species (e.g., Hargreaves et al., 2011; Kim et al., 2007a, 2007b). Generally, like in other avian species, kidney contained the highest Cd concentrations, followed by liver (e.g., Dunlin, Ferns and Anderson, 1994), while blood, feather, egg and muscle contained very low or non-detected levels (e.g., Lucia et al., 2014; Hargreaves et al., 2011; Kim et al., 2007a, b; Scherer et al., 2015).

Additionally, gender and age contribute to Cd accumulation in shorebirds, although Cd biomagnification through food webs is complex (Ferns and Anderson, 1994; McFarland et al., 2002; St Clair et al., 2015). Notably, feeding preference seems to affect Cd bioaccumulation: shorebirds feeding on intertidal invertebrates accumulated even higher Cd than gulls foraging on marine fish (Vermeer and Castilla, 1991); while individuals foraging at estuaries had higher Cd compared with terrestrial foragers (St Clair et al., 2015).

In the avian liver or kidney, it is suggested that 6 mg/kg is a poison threshold for Cd, while below 1 mg/kg is considered a background level

(Scheuhammer, 1987). There were considerable studies examining Cd concentrations for more than 20 shorebird species, yet Cd in liver or kidney generally was at background level. Only a few studies raise concerns about Cd contamination. During autumn migrations, the elevated kidney Cd concentrations (highest in Red-necked Stint, geometric mean, 84 mg/kg, dry weight) in shorebirds collected from two major stopover sites Yeongjong Island and Okgu Mudflat in Korea, indicated long term exposure along the EAAF (Kim et al., 2007a, 2010). Besides, Dunlins from Bristol Channel, UK also contained elevated concentrations of kidney Cd (geometric mean, range between 1.9 and 18.2 mg/kg, dry weight) but no histological changes were observed in internal tissues (Ferns and Anderson, 1994). Specifically, Hudsonian Whimbrels and oystercatcher species *Haematopus sp.* collected from their wintering site near a copper mine in Chile showed the alarming Cd concentrations up to 89.7 mg/kg (wet weight) in the liver (Vermeer and Castilla, 1991), the highest concentration recorded in shorebirds. Furthermore, since wader hunting exists commonly in this region, biomonitoring shorebird prey such as Pacific Sand Crab *Emerita analoga*, which widely occurs on North American and South American coasts, may be feasible to assess both ecosystem and human health (Blomqvist et al., 1987).

3.4.5. Persistent organochlorines

Evidence has revealed that exposure to POPs has proven to have chronic negative consequences in other avian groups (details see Guigueno and Fernie, 2017; Hao et al., 2021). Some legacy POPs, such as PCBs, impact avian fitness via decreased immunity against viruses (Sagerup et al., 2009), developmental deformity (Handel and Van Hemert, 2015), slower feather growth (Bustnes et al., 2002) and hinder orientation during flights (Flahr et al., 2015). The researchers further confirmed that lipids may play an important role in the distribution of POPs in tissues, suggesting migratory individuals with a high composition of fatness likely face a high level of POPs. Together, these altered biological variables/ behavioral changes may either increase energy cost or decrease refueling and flight performance in those species that undergo long-distance migration, eventually reduce migration success. In addition, at breeding grounds, POPs impact population stability via delayed egg laying (Eng et al., 2013), and reduced hatching success (Tartu et al., 2014). Recent studies focus on environmentally relevant concentrations of emerging compounds suggest reproductive failure (e.g., Fernie et al., 2009; Marteinson et al., 2011) and developing embryos (Yu et al., 2022).

A suite of legacy chlorinated compounds and their metabolites (e.g., DDTs, PCBs, PBDEs, HCCPDs and HCHs) have been examined in shorebirds in multiple regions such as Alaska, the Yellow Sea, the Gulf of Mexico, North America, and Western Europe. Among them, DDTs are of particular interest since they have been found widely in environments, even though they have been banned in agriculture use. For instance, within the remote Arctic breeding grounds, Braune and Noble (2009) found that DDTs were detected in about 98% of sampled Canadian shorebirds. Along the EAAF, shorebirds from both wintering grounds of Vietnam and India, and the breeding grounds of Lake Baikal, Russia, had elevated concentrations of DDTs. Particularly, the short-distance migrant, *Charadrius dubius* right after "Little Ringed Plover Little Ringed Plover, collected from India, showed an alarming concentration average of 57000 ng/g of DDTs (fat weight, homogenized whole body), indicating heavily polluted local contamination in India (Kunisue et al., 2003). Due to serious consequences in organochlorine pesticides (HCHs and DDTs), their usage and manufacture was restricted and even banned in many countries for decades. Yet, Melville et al. (2016) raised concerns about DDTs due to their alternative usage in antifouling paint for fish vessels in the Yellow Sea region. In the South China Sea, an important wintering or stopover region for many migrants, a decreasing trend of DDTs levels in mussels were observed, but this region is still one of the most polluted areas worldwide (Sun et al., 2020). Further north in the Yellow and Bohai Sea, Jin et al. (2017) reported 14 organochlorine pesticide residues (e.g., p, p'-DDE) in muscle and adipose tissue (e.g.,

endosulfan sulfate and p, p'-DDT) samples of Great Knots and Red Knots opportunistically collected at stopover sites in Yalujiang and Chongming. Furthermore, at Cangzhou wetland, Western Bohai Bay, DDTs were the dominant contaminant in eggs of Kentish Plover (Zheng et al., 2018). Wang et al. (2019) suggested that this could be attributed to the higher pesticide residues on the wintering grounds of Kentish Plovers, however, the contamination source is still difficult to determine due to uncertainties of their migration ecology.

There is very limited information regarding the impacts of POPs on shorebirds. As mentioned above, only one experimental study revealed that under non-optimum temperature, captive Killdeers were vulnerable to the exposure of anthropogenic PCB-126, the known endocrine disruptor at early life stage (Lunny et al., 2020). Besides, similar to previous findings of eggshell thinning in raptors (e.g., American Kestrel *Falco sparverius*, Lincer, 1975), Blus et al. (1985) found egg shells collected in Oregon and northern California in the 1950s were about 6% thinner than the pre-DDT period samples ($p < 0.05$), yet other factors that might have caused variation of eggshell thickness were not investigated. Overall, the POPs (e.g., PCBs) that already showed negative effects in other avian species, should be considered as potentially harmful to shorebirds if no data exists. And, research focusing on ecological implications from shorebird ecology perspectives should be encouraged.

4. Conclusions and future perspectives

This review represents the first attempt to summarize baseline data for shorebird exposure to hazardous pollutants. Over one-third of the research focused on descriptive knowledge concerning the baseline of shorebird contaminant exposure, but the sources of contamination remain uncertain. For instance, some studies using non-invasive tissues like feathers cannot easily trace their source of exposure (Burger et al., 1993; Kim and Koo, 2008). Exposure to particular pollutants has proven to induce negative effects on shorebirds, such as embryotoxicity and post-hatch development impairment caused by PCBs (Lunny et al., 2020); reduced reproduction performance caused by Hg and Pb (Hargreaves et al., 2011); and impaired refueling and/or migration performance related to oil contamination (Bianchini et al., 2021; Bianchini and Morrissey, 2018a,b; Maggini et al., 2017b). Most studies focused on Hg, Cd, Pb, Se, oil contamination, PCBs, and DDTs in species belonging to the Charadriidae and Scolopacidae families. Yet, the impacts of various pollutants on shorebird health have not been fully investigated. Besides, a large proportion of the research was conducted in North America. Further, studies in American key staging regions, i.e., the Gulf of Mexico, have focused on oil pollution and identified significant negative impacts on shorebirds. In the migratory hub of Bohai Bay along the EAAF, concerns about POPs are being raised. Exposure to contamination on the non-breeding grounds, particularly in East Africa and South Asia; as well as the breeding region in Siberian Arctic, is understudied and more research should be conducted in these regions. Based on our current understanding, we identified three areas that must be prioritized in future research.

1) **Determine the exposure and assess the risk of pollutants on shorebirds in their annual life cycle via advanced techniques.** Shorebirds are undoubtedly among the greatest endurance athletes for their incredible migration during the dynamic life cycle. Shorebirds seem especially vulnerable to environmental contaminants, as they forage primarily on invertebrates in coastal wetlands habitats, which are subjects to substantial inputs of various pollutants from adjacent ecosystems. Assessing the health status of shorebirds, particularly during the migratory stage, is difficult as they are rarely caught and handled. Additionally, the sources of pollution that migrants are exposed to as well as changes caused by pollution at physiological, biological, and behavioral scales need to be investigated. Recent studies in seabirds and songbirds have combined both field and laboratory work, shedding light on a new direction of future

research (Eng et al., 2019; Seewagen et al., 2019). For instance, Eng et al. (2019) and Seewagen et al. (2019) both captured wild passerines and dosed the birds up to the environmental relevant levels of specific toxins, then applied trackers to assess impacts on the behavior of treated and untreated individuals. For migratory shorebirds that undergo long-distance migration, this could be a useful means to assess the health effects of exposure to environmental relevant dose of toxicants in the wild.

Furthermore, there is insufficient data to quantify pollution risk to shorebird populations. Researchers have suggested that shorebirds, which feed on invertebrates such as mollusks and crustaceans, occupy a relatively low trophic level compared to raptors and piscivore species (Colwell, 2010), and would therefore be expected to be exposed to a lower burden of pollutants. Regarding the proper thresholds for shorebirds, Perkins et al. (2016) and Su et al. (2020) suggested lower thresholds for shorebirds, instead of using the thresholds built for piscivore birds. In short, there is an urgent need assess the risk thresholds in shorebirds to facilitate the conservation and monitoring of declining shorebird populations.

2) Screen emergent contaminants that may impact shorebirds.

There is a paucity of information about loads of emergent contaminants such as pharmaceuticals, plastic debris, and antibiotic residuals, which have been widely detected in marine and inland aquatic ecosystems. Recently, non-steroidal anti-inflammatory drugs and negative impacts have been observed in polychaeta (Świacka et al., 2021), a key dietary component for shorebirds (Choi et al., 2017). Moreover, plastic debris is generally an issue in coastal ecosystems and has already been proven to cause health issues in coastal organisms. Microplastics are currently one of the most common contaminants present in the environment. It has been confirmed that microplastics are present in shorebird food chains in intertidal habitats of Southern Europe and West Africa (Lourenço et al., 2017). Furthermore, recent studies (Rossi et al., 2019; Zhu et al., 2019) confirmed that synthetic plastic fragments were found in the stomachs of American Oystercatcher *Haematopus alliates* sampled on the Southern coast of Brazil, as well as Whimbrel located in the Southern China Sea. The researchers suggested that this debris was likely absorbed through unintentional ingestion directly from substrate consumption or indirectly through prey consumption. In addition, according to recent research, migratory shorebirds could be the reservoirs and potential spreaders of antibiotic-resistant bacteria and antibiotic resistance genes (Navedo et al., 2021), causing pollution in a pristine region. Furthermore, antibiotics might cause the alteration of gut microbiota in birds (Morgun et al., 2015). More research on antibiotic residuals is urgently needed at important staging regions like the Yellow Sea, which receives a large amount of antibiotic residuals, likely from domestic sewage, livestock wastewater, and pond aquaculture discharge (Du et al., 2017). Therefore, studies screening emergent contaminants are especially needed at locations where shorebirds are potentially exposed to them.

3) Standardized long-term non-invasive biomonitoring protocols.

The increasing production and usage of various pollutants (e.g., Cr and Pb) threaten coastal wetlands, which contain the most important shorebird habitats, such as mudflat and estuarine (British Geological Survey, 2019; C. Li et al., 2022). To track the effectiveness in regulating the production and/or emission of environmental pollutants, raptors (birds of prey and owls) and seagulls are usually used as bioindicators for regional programs in biomonitoring toxins for decades (Champoux et al., 2017; Gómez-Ramírez et al., 2014; Marth et al., 2000). For wild birds, the long-term data suggest that DDTs, PCBs, and Cd have been declining at a regional scale in the last few decades (Braune and Muir, 2017; Norstrom and Hebert, 2006), while Hg in environmental media is still not clear, probably due to complex mechanisms of bioavailability (Y. Li et al., 2022). Shorebird populations are seldom monitored for a substantial time to observe the spatial and temporal trends of exposure to environmental pollutants,

not to mention their possible ecological consequences.

Continued measure is needed to assess environmental risks and outcomes of chemical regulation and is essential for conserving migratory shorebirds effectively. For instance, the ecosystem recovery from Se-contaminated drainage is generally slow: even after 20 years of freshwater management, Se concentration in livers of Black-necked Stilts continued to decline but remained at levels associated with potential reproductive impairment (Pavegio and Kilbride, 2007; Pavegio et al., 1992). Thus, long-term biomonitoring is crucial for understanding whether the ecosystem is recovering. Unfortunately, there are no accepted or well-defined standards, guidelines, or protocols for these avian groups with large annual movements. To avoid biological variations, a standard protocol for tissues and species is required to quantify the exposure levels in a large geographic region. We recommended using non-invasive tissue e.g., feathers for biomonitoring. Feather samples are valuable as they contain a variety of information, including molting grounds based on stable isotopes, which is particularly useful for estimation of xenobiotic contamination in threatened/endangered species with trans-boundary annual movements (Ma et al., 2021). To prevent impairing their long-distance flight ability, small specific feathers should be considered. Primary coverts, instead of primary feathers, might have a less negative impact on flight performance for long-distance migrants. Although feathers are often used as a non-invasive biomonitoring tool for contamination exposure, several studies have not considered the molt ecology for migratory shorebirds, leading to estimation of body burden without a clear contamination source (e.g., Burger et al., 1993; Pandiyan et al., 2020). Thus, when considering an effective and reliable biomonitoring strategy, it is essential to choose species or populations with a well-characterized molt pattern; if necessary, further research should be conducted to fill this gap if the molt pattern remains unclear.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We thank Yuling Liang for processing the data. We thank the anonymous reviewers for their constructive comments. L.G. was supported by the China Thousand Young Talents Program (K18291101), Shenzhen Government (Y01296116), and the High-level Special Funding of the Southern University of Science and Technology (G02296302, G02296402). C.-Y. C. was supported by the Shenzhen Science and Technology Program (JCYJ20190809115207397).

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at [doi:10.1016/j.ecoenv.2022.113868](https://doi.org/10.1016/j.ecoenv.2022.113868).

References

- Ackerman, J.T., Eagles-Smith, C.A., Takekawa, J.Y., Demers, S.A., Adelsbach, T.L., Bluso, J.D., Keith Miles, A., Warnock, N., Suchanek, T.H., Schwarzbach, S.E., 2007. Mercury concentrations and space use of pre-breeding American Avocets and Black-necked Stilts in San Francisco Bay. *Sci. Total Environ.* 384, 452–466. <https://doi.org/10.1016/j.scitotenv.2007.04.027>.
- Andres, B.A., 1999. Effects of persistent shoreline oil on breeding success and chick growth in Black Oystercatchers. *Auk* 116, 640–650. <https://doi.org/10.2307/4089325>.
- Argüelles-Ticó, A., Küpper, C., Kelsh, R.N., Kosztolányi, A., Székely, T., van Dijk, R.E., 2016. Geographic variation in breeding system and environment predicts melanin-based plumage ornamentation of male and female Kentish Plovers. *Behav. Ecol. Sociobiol.* 70, 49–60. <https://doi.org/10.1007/s00265-015-2024-8>.

- Bianchini, K., Morrissey, C.A., 2018a. Assessment of shorebird migratory fueling physiology and departure timing in relation to polycyclic aromatic hydrocarbon contamination in the Gulf of Mexico. *Environ. Sci. Technol.* 52, 13562–13573.
- Bianchini, K., Morrissey, C.A., 2018b. Polycyclic aromatic hydrocarbon exposure impairs pre-migratory fueling in captive-dosed Sanderling (*Calidris alba*). *Ecotoxicol. Environ. Saf.* 161, 383–391.
- Bianchini, K., Crump, D., Farhat, A., Morrissey, C.A., 2021. Polycyclic Aromatic Hydrocarbons Alter the Hepatic Expression of Genes Involved in Sanderling (*Calidris alba*) Pre-migratory Fueling. *Environ. Toxicol. Chem.* 40, 1983–1991. <https://doi.org/10.1002/etc.5056>.
- Blomqvist, S., Frank, A., Petersson, L.R., 1987. Metals in liver and kidney tissues of autumn-migrating Dunlin *Calidris alpina* and Curlew Sandpiper *Calidris ferruginea* staging at the Baltic Sea. *Mar. Ecol. Prog. Ser.* 35, 1–13. <https://doi.org/10.3354/meps035001>.
- Blus, L.J., Henny, C.J., Krynski, A.J., 1985. Organochlorine-induced mortality and residues in Long-billed Curlews from Oregon. *Condor* 87, 563–565. <https://doi.org/10.2307/1367968>.
- Braune, B.M., Muir, D.C.G., 2017. Declining Trends of Polychlorinated Naphthalenes in Seabird Eggs from the Canadian Arctic, 1975–2014. *Environ. Sci. Technol.* 51, 3802–3808. <https://doi.org/10.1021/acs.est.7b00431>.
- Braune, B.M., Noble, D.G., 2009. Environmental contaminants in Canadian shorebirds. *Environ. Monit. Assess.* 148, 185–204. <https://doi.org/10.1007/s10661-007-0150-0>.
- British Geological Survey, 2019. World Mineral Statistics Dataset. (<https://www.bgs.ac.uk/mineralsuk/statistics/home.html>).
- Bruederle, A., Hodler, R., 2019. Effect of oil spills on infant mortality in Nigeria. *Proc. Natl. Acad. Sci. U. S. A.* 116, 5467–5471. <https://doi.org/10.1073/pnas.1818303116>.
- Bull, K.R., Every, W.J., Freestone, P., Hall, J.R., Osborn, D., Cooke, A.S., Stowe, T., 1983. Alkyl lead pollution and bird mortalities on the Mersey Estuary, UK, 1979–1981. *Environ. Pollut. Ser. A, Ecol. Biol.* 31, 239–259. [https://doi.org/10.1016/0143-1471\(83\)90062-4](https://doi.org/10.1016/0143-1471(83)90062-4).
- Burger, J., 1997. Effects of oiling on feeding behavior of Sanderlings and Semipalmated Plovers in New Jersey. *Condor* 99, 290–298. <https://doi.org/10.2307/1369935>.
- Burger, J., Gochfeld, M., 2010. Effects of lead on birds (Laridae): A review of laboratory and field studies. *J. Toxicol. Environ. Heal. Part B* 3, 59–78. <https://doi.org/10.1080/109374000281096>.
- Burger, J., Gochfeld, M., 2021. Biomonitoring selenium, mercury, and selenium: mercury molar ratios in selected species in Northeastern US estuaries: risk to biota and humans. *Environ. Sci. Pollut. Res.* 28, 18392–18406. <https://doi.org/10.1007/s11356-020-12175-z>.
- Burger, J., Gochfeld, M., Niles, L., Dey, A., Jeitner, C., Pittfield, T., Tspoura, N., 2014. Metals in tissues of migrant Semipalmated Sandpipers (*Calidris pusilla*) from Delaware Bay, New Jersey. *Environ. Res.* 133, 362–370. <https://doi.org/10.1016/j.envres.2014.05.020>.
- Burger, J., Mizrahi, D., Jeitner, C., Tspoura, N., Mobley, J., Gochfeld, M., 2019a. Metal and metalloids levels in blood of Semipalmated Sandpipers (*Calidris pusilla*) from Brazil, Suriname, and Delaware Bay: Sentinels of exposure to themselves, their prey, and predators that eat them. *Environ. Res.* 173, 77–86. <https://doi.org/10.1016/j.envres.2019.02.048>.
- Burger, J., Mizrahi, D., Tspoura, N., Jeitner, C., Gochfeld, M., 2018. Mercury, lead, cadmium, cobalt, arsenic and selenium in the blood of Semipalmated Sandpipers (*Calidris pusilla*) from Suriname, South America: Age-related differences in wintering site and comparisons with a stopover site in New Jersey, USA. *Toxics* 6, 1–11. <https://doi.org/10.3390/toxics6020027>.
- Burger, J., Seyboldt, S., Morganstein, N., Clark, K., 1993. Heavy metals and selenium in feathers of three shorebird species from Delaware Bay. *Environ. Monit. Assess.* 28, 189–198. <https://doi.org/10.1007/BF00547037>.
- Burger, J., Tspoura, N., Gochfeld, M., 2017. Metal levels in blood of three species of shorebirds during stopover on Delaware Bay reflect levels in their food, Horseshoe Crab Eggs. *Toxics* 5, 20. <https://doi.org/10.3390/toxics5030020>.
- Burger, J., Tspoura, N., Niles, L., Dey, A., Jeitner, C., Gochfeld, M., 2019b. Heavy metals in Biota in Delaware Bay, NJ: Developing a food web approach to contaminants. *Toxics* 7, 1–16. <https://doi.org/10.3390/toxics7020034>.
- Burger, J., Tspoura, N., Niles, L.J., Gochfeld, M., Dey, A., Mizrahi, D., 2015. Mercury, lead, cadmium, arsenic, chromium and selenium in feathers of shorebirds during migrating through Delaware Bay, New Jersey: Comparing the 1990s and 2011/2012. *Toxics* 3, 63–74. <https://doi.org/10.3390/toxics3010063>.
- Bursian, S.J., Dean, K.M., Harr, K.E., Kennedy, L., Link, J.E., Maggini, I., Pritsos, C., Pritsos, K.L., Schmidt, R.E., Guglielmo, C.G., 2017. Effect of oral exposure to artificially weathered Deepwater Horizon crude oil on blood chemistries, hepatic antioxidant enzyme activities, organ weights and histopathology in Western Sandpipers (*Calidris mauri*). *Ecotoxicol. Environ. Saf.* 146, 91–97. <https://doi.org/10.1016/j.ecoenv.2017.03.045>.
- Bustnes, J.O., Folstad, I., Erikstad, K.E., Fjeld, M., Miland, Ø.O., Skaare, J.U., 2002. Blood concentration of organochlorine pollutants and wing feather asymmetry in glaucous gulls. *Funct. Ecol.* 16, 617–622.
- Champoux, L., Boily, M., Fitzgerald, G., 2017. Thyroid hormones, retinol and clinical parameters in relation to mercury and organohalogen contaminants in great blue heron (*Ardea herodias*) Nestlings from the St. Lawrence River, Quebec, Canada. *Arch. Environ. Contam. Toxicol.* 72, 200–214. <https://doi.org/10.1007/s00244-017-0364-2>.
- Carlson, J.R., Cristol, D.A., Swaddle, J.P., 2014. Dietary mercury exposure causes decreased escape takeoff flight performance and increased molt rate in European starlings (*Sturnus vulgaris*). *Ecotoxicology* 23, 1464–1473. <https://doi.org/10.1007/s10646-014-1288-5>.
- Chapagain, S.K., Shrestha, S., Du Laing, G., Verloo, M., Kazama, F., 2009. Spatial distribution of arsenic in the intertidal sediments of River Scheldt, Belgium. *Environ. Int.* 35, 461–465. <https://doi.org/10.1016/j.envint.2008.07.019>.
- Chen, L., Xu, Z., Ding, X., Zhang, W., Huang, Y., Fan, R., Sun, J., Liu, M., Qian, D., Feng, Y., 2012. Spatial trend and pollution assessment of total mercury and methylmercury pollution in the Pearl River Delta soil, South China. *Chemosphere* 88, 612–619. <https://doi.org/10.1016/j.chemosphere.2012.03.041>.
- Choi, C.-Y., Battley, P.F., Potter, M.A., Ma, Z., Melville, D.S., Sukkaweemane, P., 2017. How migratory shorebirds selectively exploit prey at a staging site dominated by a single prey species. *Auk* 134, 76–91. <https://doi.org/10.1642/auk-16-58.1>.
- Colwell, M.A., 2010. *Shorebird Ecology, Conservation, and Management*. Univ of California Press.
- Dietz, R., Sonne, C., Basu, N., Braune, B., O'Hara, T., Letcher, R.J., Scheuhammer, T., Andersen, M., Andreasen, C., Andriashek, D., 2013. What are the toxicological effects of mercury in Arctic biota? *Sci. Total Environ.* 443, 775–790.
- Du, J., Zhao, H., Liu, S., Xie, H., Wang, Y., Chen, J., 2017. Antibiotics in the coastal water of the South Yellow Sea in China: Occurrence, distribution and ecological risks. *Sci. Total Environ.* 595, 521–527. <https://doi.org/10.1016/j.scitotenv.2017.03.281>.
- Eagles-Smith, C.A., Ackerman, J.T., De La Cruz, S.E.W., Takekawa, J.Y., 2009. Mercury bioaccumulation and risk to three waterbird foraging guilds is influenced by foraging ecology and breeding stage. *Environ. Pollut.* 157, 1993–2002. <https://doi.org/10.1016/j.envpol.2009.03.030>.
- Eisler, R., 1985. Cadmium hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish and Wildlife Service Biological Report 85 (1.2). *Contam. Hazard Rev. Report No. 2*.
- Eisler, R., 1988. Lead hazards to fish, wildlife, and invertebrates: a synoptic review. U.S. Fish and Wildlife Service Biological Report 85 (1.14). *Contam. Hazard Rev. Report No. 14*.
- Eng, M.L., Stutchbury, B.J.M., Morrissey, C.A., 2019. A neonicotinoid insecticide reduces fueling and delays migration in songbirds. *Science* 365, 1177–1180. <https://doi.org/10.1126/science.aaw9419>.
- Eng, M.L., Williams, T.D., Elliott, J.E., 2013. Developmental exposure to a brominated flame retardant: An assessment of effects on physiology, growth, and reproduction in a songbird, the zebra finch. *Environ. Pollut.* 178, 343–349. <https://doi.org/10.1016/j.envpol.2013.03.037>.
- Fernández, G., García-Hernández, J., Cruz-Acevedo, E., Castillo-Guerrero, J.A., 2019. Patterns of mercury accumulation in tissues of Western Sandpipers (*Calidris mauri*) wintering on the Coast of Sinaloa, Mexico. *Waterbirds* 41, 438–442. <https://doi.org/10.1675/063.041.0408>.
- Fernie, K.J., Shutt, J.L., Letcher, R.J., Ritchie, I.J., Bird, D.M., 2009. Environmentally relevant concentrations of DE-71 and HBCD alter eggshell thickness and reproductive success of American Kestrels. *Environ. Sci. Technol.* 43, 2124–2130. <https://doi.org/10.1021/es8027346>.
- Ferns, P.N., Anderson, J.I., 1997. Lead in the diet and body tissues of Dunlins, *Calidris alpina*, from the Bristol Channel, UK. *Environ. Pollut.* 96, 35–42. [https://doi.org/10.1016/S0269-7491\(97\)00008-0](https://doi.org/10.1016/S0269-7491(97)00008-0).
- Ferns, P.N., Anderson, J.I., 1994. Cadmium in the diet and body tissues of dunlins *Calidris alpina*, from the Bristol channel, UK. *Environ. Pollut.* 86, 225–231.
- Flahr, L.M., Michel, N.L., Zahara, A.R.D., Jones, P.D., Morrissey, C.A., 2015. Developmental exposure to Aroclor 1254 alters migratory behavior in juvenile European starlings (*Sturnus vulgaris*). *Environ. Sci. Technol.* 49, 6274–6283.
- Flickinger, E.L., King, K.A., 1972. Some Effects of Aldrin-Treated Rice on Gulf Coast Wildlife. *J. Wildl. Manag.* 36, 706–727. <https://doi.org/10.2307/3799422>.
- Gibson, D., Catlin, D.H., Hunt, K.L., Fraser, J.D., Karpanty, S.M., Friedrich, M.J., Bimbi, M.K., Cohen, J.B., Maddock, S.B., 2017. Evaluating the impact of man-made disasters on imperiled species: Piping Plovers and the Deepwater Horizon oil spill. *Biol. Conserv.* 212, 48–62. <https://doi.org/10.1016/j.biocon.2017.05.027>.
- Goede, A.A., 1993. Selenium in eggs and parental blood of a Dutch marine wader. *Arch. Environ. Contam. Toxicol.* 25, 79–84. <https://doi.org/10.1007/BF00230715>.
- Goede, A.A., Wolterbeek, H.T., 1994. The possible role of selenium in antioxidantation in marine waders; a preliminary study. *Sci. Total Environ.* 144, 241–246.
- Gómez-Ramírez, P., Shore, R.F., van den Brink, N.W., van Hattum, B., Bustnes, J.O., Duke, G., Fritsch, C., García-Fernández, A.J., Helander, B.O., Jaspers, V., Krone, O., Martínez-López, E., Mateo, R., Movalli, P., Sonne, C., 2014. An overview of existing raptor contaminant monitoring activities in Europe. *Environ. Int.* 67, 12–21. <https://doi.org/10.1016/j.envint.2014.02.004>.
- Grade, T.J., Pokras, M.A., Laflamme, E.M., Vogel, H.S., 2018b. Population-level effects of lead fishing tackle on common loons. *J. Wildl. Manag.* 82, 155–164. <https://doi.org/10.1002/jwmg.21348>.
- Guigueno, M.F., Fernie, K.J., 2017. Birds and flame retardants: A review of the toxic effects on birds of historical and novel flame retardants. *Environ. Res.* 154, 398–424. <https://doi.org/10.1016/j.envres.2016.12.033>.
- Haig, S.M., D'Elia, J., Eagles-Smith, C., Fair, J.M., Gervais, J., Herring, G., Rivers, J.W., Schulz, J.H., 2014. The persistent problem of lead poisoning in birds from ammunition and fishing tackle. *Condor* 116, 408–428. <https://doi.org/10.1650/CONDOR-14-36.1>.
- Handel, C.M., Van Hemert, C., 2015. Environmental contaminants and chromosomal damage associated with beak deformities in a resident North American Passerine. *Environ. Toxicol. Chem.* 34, 314–327. <https://doi.org/10.1002/etc.2799>.
- Hao, Y., Zheng, S., Wang, P., Sun, H., Matsiko, J., Li, W., Li, Y., Zhang, Q., Jiang, G., 2021. Ecotoxicology of persistent organic pollutants in birds. *Environ. Sci. Process. Impacts* 23, 400–416. <https://doi.org/10.1039/d0em00451k>.
- Hargreaves, A.L., Whiteside, D.P., Gilchrist, G., 2011. Concentrations of 17 elements, including mercury, in the tissues, food and abiotic environment of arctic shorebirds. *Sci. Total Environ.* 409, 3757–3770. <https://doi.org/10.1016/j.scitotenv.2011.06.043>.

- Hargreaves, A.L., Whiteside, D.P., Gilchrist, G., 2010. Concentrations of 17 elements, including mercury, and their relationship to fitness measures in arctic shorebirds and their eggs. *Sci. Total Environ.* 408, 3153–3161. <https://doi.org/10.1016/j.scitotenv.2010.03.027>.
- Hartman, C.A., Ackerman, J.T., Herzog, M.P., 2019. Mercury exposure and altered parental nesting behavior in a wild songbird. *Environ. Sci. Technol.* 53, 5396–5405. <https://doi.org/10.1021/acs.est.8b07227>.
- Heinz, G.H., Haseltine, S.D., Sileo, L., 1983. Altered avoidance behavior of young black ducks fed cadmium. *Environ. Toxicol. Chem.* 2, 419–421.
- Henkel, J.R., Sigel, B.J., Taylor, C.M., 2014. Oiling rates and condition indices of shorebirds on the northern gulf of Mexico following the deepwater horizon oil spill. *J. F. Ornithol.* <https://doi.org/10.1111/jof.12080>.
- Hiller, B.J., Barclay, J.S., 2011. Concentrations of heavy metals in American woodcock harvested in Connecticut. *Arch. Environ. Contam. Toxicol.* 60, 156–164. <https://doi.org/10.1007/s00244-010-9525-2>.
- Hui, C.A., 1998. Metal and trace element burdens in two shorebird species at two sympatric wintering sites in southern California. *Environ. Monit. Assess.* 50, 233–247. <https://doi.org/10.1023/A:1005850112994>.
- Hui, C.A., Takekawa, J.Y., Warnock, S.E., 2001. Contaminant profiles of two species of shorebirds foraging together at two neighboring sites in south San Francisco Bay. *California* 107–121.
- Irons, D.B., Kendall, S.J., Erickson, W.P., McDonald, L.L., Lance, B.K., 2000. Nine years after the Exxon Valdez oil spill: Effects on marine bird populations in Prince William Sound, Alaska. *Condor* 102, 723–737. <https://doi.org/10.2307/1370300>.
- IUCN 2022. The IUCN Red List of Threatened Species. Version 2021–3. (<https://www.iucnredlist.org>).
- Jin, X., Dai, L.-S., Chen, Q.-F., Zhou, Y.-H., Qiu, Yan-Ling, Ma, Z.-J., 2017. Contamination of organochlorine pesticides in Great Knots (*Calidris tenuirostris*) and Red Knots (*C. canutus*). *Chin. J. Zool.* 52, 1–10. <https://doi.org/10.13859/j.cjz.201701001>.
- Kenow, K.P., Grasman, K.A., Hines, R.K., Meyer, M.W., Gendron-Fitzpatrick, A., Spalding, M.G., Gray, B.R., 2007. Effects of methylmercury exposure on the immune function of juvenile Common Loons (*Gavia immer*). *Environ. Toxicol. Chem.* 26, 1460–1469.
- Kim, J., Koo, T.H., 2010. Acute and/or chronic contaminations of heavy metals in shorebirds from Korea. *J. Environ. Monit.* 12, 1613–1618. <https://doi.org/10.1039/c0em00042f>.
- Kim, J., Park, S.K., Koo, T.H., 2007a. Lead and cadmium concentrations in shorebirds from the Yeongjong Island, Korea. *Environ. Monit. Assess.* 134, 355–361. <https://doi.org/10.1007/s10661-007-9626-1>.
- Kim, J., Park, S.K., Koo, T.H., 2007b. Trace elements and pollutants concentrations in shorebirds from Yeongjong Island, Korea in the East Asian-Australian migration flyways. *Ecotoxicology* 16, 403–410. <https://doi.org/10.1007/s10646-007-0141-5>.
- Kim, J., Koo, T.H., 2008. Heavy metal concentrations in feathers of Korean shorebirds. *Arch. Environ. Contam. Toxicol.* 55, 122–128. <https://doi.org/10.1007/s00244-007-9089-y>.
- King, M.D., Elliott, J.E., Williams, T.D., 2021. Effects of petroleum exposure on birds: A review. *Sci. Total Environ.* 755, 142834 <https://doi.org/10.1016/j.scitotenv.2020.142834>.
- Koeman, J.H., Peeters, W.H.M., Koudstaal-Hol, C.H.M., Tjioe, P.S., de Goeij, J.J.M., 1973. Mercury-selenium correlations in marine mammals. *Nature* 245, 385–386. <https://doi.org/10.1038/245385a0>.
- Kunise, T., Watanabe, M., Subramanian, A., Sethuraman, A., Titenko, A.M., Qui, V., Prudente, M., Tanabe, S., 2003. Accumulation features of persistent organochlorines in resident and migratory birds from Asia. *Environ. Pollut.* 125, 157–172. [https://doi.org/10.1016/S0269-7491\(03\)00074-5](https://doi.org/10.1016/S0269-7491(03)00074-5).
- Lambeck, R.H.D., Nieuwenhuize, J., van Liere, J.M., 1991. Polychlorinated biphenyls in oystercatchers (*Haematopus ostralegus*) from the oosterschelde (Dutch Delta area) and the Western Wadden Sea, that died from starvation during severe winter weather. *Environ. Pollut.* 71, 1–16. [https://doi.org/10.1016/0269-7491\(91\)90040-4](https://doi.org/10.1016/0269-7491(91)90040-4).
- Levengood, J.M., Smith, R.V., Gay, D.A., Davis, M.A., Stafford, J.D., 2014. Mercury in migrating shorebirds in the Illinois river valley. *Waterbirds* 37, 225–229. <https://doi.org/10.1675/063.037.0212>.
- Li, C., Wang, H., Liao, X., Xiao, R., Liu, K., Bai, J., Li, B., He, Q., 2022. Heavy metal pollution in coastal wetlands: A systematic review of studies globally over the past three decades. *J. Hazard. Mater.* <https://doi.org/10.1016/j.jhazmat.2021.127312>.
- Li, Y., Chen, L., Liang, S., Zhou, H., Liu, Y.-R., Zhong, H., Yang, Z., 2022. Looping Mercury Cycle in Global Environmental-Economic System Modeling. *Environ. Sci. Technol.* 56, 2861–2879. <https://doi.org/10.1021/acs.est.1c03936>.
- Lincer, J.L., 1975. DDE-induced eggshell-thinning in the American Kestrel: A comparison of the field situation and laboratory results. *J. Appl. Ecol.* 12, 781. <https://doi.org/10.2307/2402090>.
- Lindberg, P., Odsjö, T., Reutergerd, L., 1985. Residue levels of polychlorobiphenyls, EDDT, and mercury in bird species commonly preyed upon by the peregrine falcon (*Falco peregrinus Tunst.*) in Sweden. *Arch. Environ. Contam. Toxicol.* 14, 203–212. <https://doi.org/10.1007/BF01055613>.
- Lourengo, P.M., Serra-Gonçalves, C., Ferreira, J.L., Catty, T., Granadeiro, J.P., 2017. Plastic and other microfibers in sediments, macroinvertebrates and shorebirds from three intertidal wetlands of southern Europe and west Africa. *Environ. Pollut.* 231, 123–133. <https://doi.org/10.1016/j.envpol.2017.07.103>.
- Lucia, M., Bocher, P., Chambosse, M., Delaporte, P., Bustamante, P., 2014. Trace element accumulation in relation to trophic niches of shorebirds using intertidal mudflats. *J. Sea Res.* 92, 134–143. <https://doi.org/10.1016/j.seares.2013.08.008>.
- Lucia, M., Bocher, P., Cosson, R.P., Churlaud, C., Bustamante, P., 2012a. Evidence of species-specific detoxification processes for trace elements in shorebirds. *Ecotoxicology* 21, 2349–2362. <https://doi.org/10.1007/s10646-012-0991-3>.
- Lucia, M., Bocher, P., Cosson, R.P., Churlaud, C., Robin, F., Bustamante, P., 2012b. Insight on trace element detoxification in the Black-tailed Godwit (*Limosa limosa*) through genetic, enzymatic and metallothionein analyses. *Sci. Total Environ.* 423, 73–83. <https://doi.org/10.1016/j.scitotenv.2012.02.005>.
- Lunny, E., Eng, M.L., Gurney, K.E.B., Morrissey, C.A., 2020. Incubation temperature and PCB-126 exposure interactively impair shorebird embryo and post-hatch development. *Environ. Res.* 188, 109779 <https://doi.org/10.1016/j.envres.2020.109779>.
- Ma, Y., Branfireun, B.A., Hobson, K.A., Guglielmo, C.G., 2018a. Evidence of negative seasonal carry-over effects of breeding ground mercury exposure on survival of migratory songbirds. *J. Avian Biol.* 49 <https://doi.org/10.1111/jav.01656>.
- Ma, Y., Hobson, K.A., Kardynal, K.J., Guglielmo, C.G., Branfireun, B.A., 2021. Inferring spatial patterns of mercury exposure in migratory boreal songbirds: Combining feather mercury and stable isotope ($\delta^2\text{H}$) measurements. *Sci. Total Environ.* 762, 143109 <https://doi.org/10.1016/j.scitotenv.2020.143109>.
- Ma, Y., Perez, C.R., Branfireun, B.A., Guglielmo, C.G., 2018b. Dietary exposure to methylmercury affects flight endurance in a migratory songbird. *Environ. Pollut.* 234, 894–901. <https://doi.org/10.1016/j.envpol.2017.12.011>.
- Ma, Z., Melville, D.S., Liu, J., Chen, Y., Yang, H., Ren, W., Zhang, Z., Piersma, T., Li, B., 2014. Rethinking China's new great wall. *Science* 346, 912–914. <https://doi.org/10.1126/science.1257258>.
- Maggini, I., Kennedy, L.V., Bursian, S.J., Dean, K.M., Gerson, A.R., Harr, K.E., Link, J.E., Pritsos, C.A., Pritsos, K.L., Guglielmo, C.G., 2017a. Toxicological and thermoregulatory effects of feather contamination with artificially weathered MC 252 oil in Western Sandpipers (*Calidris mauri*). *Ecotoxicol. Environ. Saf.* 146, 118–128. <https://doi.org/10.1016/j.ecoenv.2017.04.025>.
- Maggini, I., Kennedy, L.V., Macmillan, A., Elliott, K.H., Dean, K., Guglielmo, C.G., 2017b. Light oiling of feathers increases flight energy expenditure in a migratory shorebird. *J. Exp. Biol.* 220, 2372–2379.
- Marteinson, S.C., Kimmins, S., Bird, D.M., Shutt, J.L., Letcher, R.J., Ritchie, I.J., Fernie, K.J., 2011. Embryonic Exposure to the Polybrominated Diphenyl Ether Mixture, DE-71, Affects Testes and Circulating Testosterone Concentrations in Adult American Kestrels (*Falco sparverius*). *Toxicol. Sci.* 121, 168–176. <https://doi.org/10.1093/toxsci/ktf027>.
- Marth, P., Martens, D., Schramm, K.-W., Schmitzer, J., Oxynos, K., Kettrup, A., 2000. Environmental specimen banking. Herring gull eggs and brems as bioindicators for monitoring longterm and spatial trends of chlorinated hydrocarbons. *Pure Appl. Chem.* 72, 1027–1034. <https://doi.org/10.1351/pac200072061027>.
- McFarland, C.N., Bendell-Young, L.I., Guglielmo, C., Williams, T.D., 2002. Kidney, liver and bone cadmium content in the western sandpiper in relation to migration. *J. Environ. Monit.* 4, 791–795. <https://doi.org/10.1039/B206045K>.
- Melville, D.S., Chen, Y., Ma, Z., 2016. Shorebirds along the Yellow Sea coast of China face an uncertain future—a review of threats. *Emu-Austral Ornithol.* 116, 100–110. <https://doi.org/10.1071/MU15045>.
- Miller, D.S., Peakall, D.B., Kinter, W.B., 1978. Ingestion of crude oil: sublethal effects in Herring Gull chicks. *Science* 199, 315–317. <https://doi.org/10.1126/science.145655>.
- Mora, M.A., Taylor, R.J., Brattin, B.L., 2007. Potential ecotoxicological significance of elevated concentrations of strontium in eggshells of passerine birds. *Condor* 109, 199–205.
- Morgun, A., Dzutsev, A., Dong, X., Greer, R.L., Sexton, D.J., Ravel, J., Schuster, M., Hsiao, W., Matzinger, P., Shulzhenko, N., 2015. Uncovering effects of antibiotics on the host and microbiota using transkingdom gene networks. *Gut* 64, 1732–1743. <https://doi.org/10.1136/gutjnl-2014-308820>.
- Mudge, G.P., 1983. The incidence and significance of ingested lead pellet poisoning in British wildfowl. *Biol. Conserve* 27, 333–372.
- Murphy, S.M., Day, R.H., Wiens, J.A., Parker, K.R., 1997. Effects of the Exxon Valdez Oil Spill on Birds: Comparisons of Pre- and Post-Spill Surveys in Prince William Sound, Alaska. *Condor* 99, 299–313. <https://doi.org/10.2307/1369936>.
- Navedo, J.G., Araya, V., Verdugo, C., 2021. Upraising a silent pollution: Antibiotic resistance at coastal environments and transference to long-distance migratory shorebirds. *Sci. Total Environ.* 777, 146004 <https://doi.org/10.1016/j.scitotenv.2021.146004>.
- Norstrom, R.J., Hebert, C.E., 2006. Comprehensive re-analysis of archived herring gull eggs reconstructs historical temporal trends in chlorinated hydrocarbon contamination in Lake Ontario and Green Bay, Lake Michigan, 1971–1982. *J. Environ. Monit.* 8, 835–847. <https://doi.org/10.1039/b602378a>.
- Pandiyani, J., Mahboob, S., Jagadheesan, R., Elumalai, K., Krishnappa, K., Al-Misned, F., Ali Kaimkhani, Z., Govindarajan, M., 2020. A novel approach to assess the heavy metal content in the feathers of shorebirds: A perspective of environmental research. *J. King Saud. Univ. - Sci.* 32, 3065–3071. <https://doi.org/10.1016/j.jksus.2020.08.014>.
- Pavegio, F.L., Kilbride, K.M., 2007. Selenium in Aquatic Birds from Central California. *J. Wildl. Manag.* 71, 2550–2555. <https://doi.org/10.2193/2006-250>.
- Pavegio, F.L., Bunck, C.M., Heinz, G.H., 1992. Selenium and boron in aquatic birds from central California. *J. Wildl. Manag.* 56, 31–42.
- Pedersen, H.C., Saether, M., 1999. Effects of cadmium on parental behaviour in free-living willow ptarmigan hens. *Ecotoxicology* 8, 1–7. <https://doi.org/10.1023/A:1008836908985>.
- Perez, C.R., Moye, J.K., Cacula, D., Dean, K.M., Pritsos, C.A., 2017. Low level exposure to crude oil impacts avian flight performance: The Deepwater Horizon oil spill effect on migratory birds. *Ecotoxicol. Environ. Saf.* 146, 98–103. <https://doi.org/10.1016/j.ecoenv.2017.05.028>.
- Perkins, M., Ferguson, L., Lanctot, R.B., Stenhouse, I.J., Kendall, S., Brown, S., Gates, H. R., Hall, J.O., Regan, K., Evers, D.C., 2016. Mercury exposure and risk in breeding

- and staging Alaskan shorebirds. *Condor* 118, 571–582. <https://doi.org/10.1650/CONDOR-16-36.1>.
- Picone, M., Corami, F., Gaetan, C., Basso, M., Battiston, A., Panzarin, L., Volpi Ghirardini, A., 2019. Accumulation of trace elements in feathers of the Kentish Plover *Charadrius alexandrinus*. *Ecotoxicol. Environ. Saf.* 179, 62–70. <https://doi.org/10.1016/j.ecoenv.2019.04.051>.
- Pokras, M., Kneeland, M.R., 2009. Understanding Lead Uptake and Effects across Species Lines: A Conservation Medicine Based Approach. In: Watson, R.T., Fuller, M., Pokras, M., W.G.H. (Eds.), *Ingestion of Lead from Spent Ammunition: Implications for Wildlife and Humans*. The Peregrine Fund, Boise. <https://doi.org/10.4080/ilsa.2009.0101>.
- Pratte, I., Noble, D.G., Mallory, M.L., Braune, B.M., Provencher, J.F., 2020. The influence of migration patterns on exposure to contaminants in Nearctic shorebirds: a historical study. *Environ. Monit. Assess.* 192 <https://doi.org/10.1007/s10661-020-8218-1>.
- Roodbergen, M., Klokk, C., van der Hout, A., 2008. Transfer of heavy metals in the food chain earthworm Black-tailed Godwit (*Limosa limosa*): Comparison of a polluted and a reference site in The Netherlands. *Sci. Total Environ.* 406, 407–412. <https://doi.org/10.1016/j.scitotenv.2008.06.051>.
- Rossi, L.C., Scherer, A.L., Petry, M.V., 2019. First record of debris ingestion by the shorebird American Oystercatcher (*Haematopus palliatus*) on the Southern coast of Brazil. *Mar. Pollut. Bull.* 138, 235–240. <https://doi.org/10.1016/j.marpolbul.2018.11.051>.
- Saalfeld, D.T., Matz, A.C., McCaffery, B.J., Johnson, O.W., Bruner, P., Lanctot, R.B., 2016. Inorganic and organic contaminants in Alaskan shorebird eggs. *Environ. Monit. Assess.* 188. <https://doi.org/10.1007/s10661-016-5270-y>.
- Sagerup, K., Larsen, H.J.S., Skaare, J.U., Johansen, G.M., Gabrielsen, G.W., 2009. The Toxic Effects of Multiple Persistent Organic Pollutant Exposures on the Post-Hatch Immunity Maturation of Glaucous Gulls. *J. Toxicol. Environ. Heal. A-Curr. ISSUES* 72, 870–883. <https://doi.org/10.1080/15287390902959516>.
- Schäfer, J., Castelle, S., Blanc, G., Dabrin, A., Masson, M., Lanceleur, L., Bossy, C., 2010. Mercury methylation in the sediments of a macrotidal estuary (Gironde Estuary, south-west France). *Estuar. Coast. Shelf Sci.* 90, 80–92. <https://doi.org/10.1016/j.ecss.2010.07.007>.
- Scherer, J.F.M., Scherer, A.L., Barbieri, E., Petry, M.V., Valiati, V.H., 2015. Trace elements concentrations in Buff-breasted Sandpiper sampled in Lagoa do Peixe National Park, Southern Brazil. *Braz. J. Biol.* 75, 932–935. <https://doi.org/10.1590/1519-6984.02914>.
- Scheuhammer, A.M., 1987. The chronic toxicity of aluminium, cadmium, mercury, and lead in birds: A review. *Environ. Pollut.* 46, 263–295. [https://doi.org/10.1016/0269-7491\(87\)90173-4](https://doi.org/10.1016/0269-7491(87)90173-4).
- Schwemmer, P., Covaci, A., Das, K., Lepoint, G., Adler, S., Garthe, S., 2015. Assessment of contaminant levels and trophic relations at a World Heritage Site by measurements in a characteristic shorebird species. *Environ. Res.* 136, 163–172. <https://doi.org/10.1016/j.envres.2014.10.021>.
- Seewagen, C.L., Ma, Y., Morbey, Y.E., Guglielmo, C.G., 2019. Stopover departure behavior and flight orientation of spring-migrant Yellow-rumped Warblers (*Setophaga coronata*) experimentally exposed to methylmercury. *J. Ornithol.* 160, 617–624. <https://doi.org/10.1007/s10336-019-01641-2>.
- Smith, H.G., Bean, D.C., Clarke, R.H., Loyn, R., Larkins, J., Hassell, C., Greenhill, A.R., 2022. Presence and antimicrobial resistance profiles of *Escherichia coli*, *Enterococcus* spp. and *Salmonella* sp. in 12 species of Australian shorebirds and terns. *Zoonoses Public Health* 1–10. <https://doi.org/10.1111/zph.12950>.
- Spalding, M.G., Frederick, P.C., McGill, H.C., Bouton, S.N., Richey, L.J., Schumacher, I. M., Blackmore, C.G.M., Harrison, J., 2000. Histologic, neurologic, and immunologic effects of methylmercury in captive great egrets. *J. Wildl. Dis.* 36, 423–435.
- St Clair, C.T., Baird, P., Ydenberg, R., Elner, R., Bendell, L.L., 2015. Trace elements in Pacific Dunlin (*Calidris alpina pacifica*): patterns of accumulation and concentrations in kidneys and feathers. *Ecotoxicology* 24, 29–44. <https://doi.org/10.1007/s10646-014-1352-1>.
- Stock, M., Herber, R., Geron, H., 2007. Cadmium levels in oystercatcher *Haematopus ostralegus* from the German Wadden Sea. *Mar. Ecol. Prog. Ser.* 53, 227–234. <https://doi.org/10.3354/meps053227>.
- Su, T., Lin, X., Huang, Q., Jiang, D., Zhang, C., Zhang, X., Dingle, C., Goodale, E., Que, P., Wang, R., Liu, Y., 2020. Mercury exposure in sedentary and migratory Charadrius plovers distributed widely across China. *Environ. Sci. Pollut. Res.* 27, 4236–4245. <https://doi.org/10.1007/s11356-019-06873-6>.
- Sun, R., Yu, J., Liao, Y., Chen, J., Wu, Z., Mai, B., 2020. Geographical distribution and risk assessment of dichlorodiphenyltrichloroethane and its metabolites in *Perna viridis* mussels from the northern coast of the South China Sea. *Mar. Pollut. Bull.* 151, 110819 <https://doi.org/10.1016/j.marpolbul.2019.110819>.
- Świacka, K., Michnowska, A., Maculewicz, J., Caban, M., Smolarz, K., 2021. Toxic effects of NSAIDs in non-target species: A review from the perspective of the aquatic environment. *Environ. Pollut.* 273, 115891 <https://doi.org/10.1016/j.envpol.2020.115891>.
- Tang, Z., Huang, Q., Nie, Z., Yang, Y., 2015. Pollution threatens migratory shorebirds. *Science* 350, 1176–1177. <https://doi.org/10.1126/science.350.6265.1176-c>.
- Tartu, S., Gabrielsen, G.W., Blevin, P., Ellis, H., Bustnes, J.O., Herzke, D., Chastel, O., 2014. Endocrine and Fitness Correlates of Long-Chain Perfluorinated Carboxylates Exposure in Arctic Breeding Black-Legged Kittiwakes. *Environ. Sci. Technol.* 48, 13504–13510. <https://doi.org/10.1021/es503297n>.
- Tavares, P.C., McGill, R., Newton, J., Pereira, E., Duarte, A., Furness, R.W., 2009. Relationships between carbon sources, trophic level and mercury exposure in generalist shorebirds revealed by stable isotope ratios in chicks. *Waterbirds* 32, 311–321. <https://doi.org/10.1675/063.032.0211>.
- Tavares, P.C., Monteiro, L.R., Lopes, R.J., Correia Santos, M.M., Furness, R.W., 2004. Intraspecific variation of mercury contamination in chicks of Black-winged Stilt (*Himantopus himantopus*) in Coastal Wetlands from Southwestern Europe. *Bull. Environ. Contam. Toxicol.* <https://doi.org/10.1007/s00128-004-0265-8>.
- Tsipoura, N., Burger, J., Niles, L., Dey, A., Gochfeld, M., Peck, M., Mizrahi, D., 2017. Metal Levels in Shorebird Feathers and Blood During Migration Through Delaware Bay. *Arch. Environ. Contam. Toxicol.* 72, 562–574. <https://doi.org/10.1007/s00244-017-0400-2>.
- Tsuji, L.J.S., Martin, I.D., Martin, E.S., LeBlanc, A., Dumas, P., 2008. Spring-harvested game birds in the Western James Bay region of Northern Ontario, Canada: the amount of organochlorines in matched samples of breast muscle, skin, and abdominal fat. *Environ. Monit. Assess.* 146, 91–104. <https://doi.org/10.1007/s10661-007-0062-z>.
- UN Environment, 2019. *Global Mercury Assessment*. Geneva, Switzerland.
- UNEP, 2021. *Stockholm Convention on persistent organic pollutants*. (<http://chm.pops.int>), access: March 2021.
- Vallverdú-Coll, N., Mateo, R., Mougeot, F., Ortiz-Santaliestra, M.E., 2019. Immunotoxic effects of lead on birds. *Sci. Total Environ.* 689, 505–515. <https://doi.org/10.1016/j.scitotenv.2019.06.251>.
- Van Gils, J.A., Lisovski, S., Lok, T., Meissner, W., Ozarowska, A., De Fouw, J., Rakhimberdiev, E., Soloviev, M.Y., Piersma, T., Klaassen, M., 2016. Body shrinkage due to Arctic warming reduces Red Knot fitness in tropical wintering range. *Science* 352, 819–821. <https://doi.org/10.1126/science.aad6351>.
- Vermeer, K., Castilla, J.C., 1991. High cadmium residues observed during a pilot study in shorebirds and their prey downstream from the El Salvador copper mine, Chile. *Bull. Environ. Contam. Toxicol.* 46, 242–248. <https://doi.org/10.1007/BF01691944>.
- Vidal, M., García-Miranda, O., Domínguez, J., 2021. Sex-related and spatial variation in trace elements in feathers of the Kentish Plover (*Charadrius alexandrinus*) from different coastal areas of the Iberian Peninsula. *Sci. Total Environ.* 766, 144628 <https://doi.org/10.1016/j.scitotenv.2020.144628>.
- Wang, D., Zheng, S., Wang, P., Matsiko, J., Sun, H., Hao, Y., Li, Y., Zhang, Z., Que, P., Meng, D., Zhang, Q., Jiang, G., 2019. Effects of migration and reproduction on the variation in persistent organic pollutant levels in Kentish Plovers from Cangzhou Wetland, China. *Sci. Total Environ.* 670, 122–128. <https://doi.org/10.1016/j.scitotenv.2019.03.039>.
- Wania, F., Mackay, D., 1993. Global fractionation and cold condensation of low volatility organochlorine compounds in polar regions. *Ambio* 22, 10–18.
- Warnock, N., Schwarzbach, S.E., 1995. Incidental kill of dunlin and killdeer by strychnine. *J. Wildl. Dis.* 31, 566–569. <https://doi.org/10.7589/0090-3558-31.4.566>.
- Yong, D.L., Liu, Y., Low, B.W., Espanola, C.P., Choi, C.-Y., Kawakami, K., 2015. Migratory songbirds in the East Asian-Australasian Flyway: A review from a conservation perspective. *Bird. Conserv. Int.* 25, 1–37. <https://doi.org/10.1017/S0959270914000276>.
- Yu, Y.J., Li, Z.R., Zhu, Y., Li, L.Z., Zhang, L.H., Xiang, M.D., Zeng, E.Y., 2022. Significance of biotransformation and excretion on the enantioselective bioaccumulation of hexabromocyclododecane (HBCDD) in laying hens and developing chicken embryos. *J. Hazard. Mater.* 422 <https://doi.org/10.1016/j.jhazmat.2021.126749>.
- Zamani-Ahmadmahmoodi, R., Esmaili-Sari, A., Savabieasfahani, M., Ghasempouri, S.M., Bahramifar, N., 2010. Mercury pollution in three species of waders from shadegan wetlands at the head of the Persian gulf. *Bull. Environ. Contam. Toxicol.* 84, 326–330. <https://doi.org/10.1007/s00128-010-9933-z>.
- Zhang, P., Song, J., Yuan, H., 2009. Persistent organic pollutant residues in the sediments and mollusks from the Bohai Sea coastal areas, North China: An overview. *Environ. Int.* 35, 632–646. <https://doi.org/10.1016/j.envint.2008.09.014>.
- Zheng, S., Wang, P., Sun, H., Matsiko, J., Hao, Y., Meng, D., Li, Y., Zhang, G., Zhang, Q., Jiang, G., 2018. Tissue distribution and maternal transfer of persistent organic pollutants in Kentish Plovers (*Charadrius alexandrinus*) from Cangzhou Wetland, Bohai Bay, China. *Sci. Total Environ.* 612, 1105–1113. <https://doi.org/10.1016/j.scitotenv.2017.08.323>.
- Zhu, C., Li, D., Sun, Y., Zheng, X., Peng, X., Zheng, K., Hu, B., Luo, X., Mai, B., 2019. Plastic debris in marine birds from an island located in the South China Sea. *Mar. Pollut. Bull.* 149, 110566 <https://doi.org/10.1016/j.marpolbul.2019.110566>.

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
American Avocet	<i>Recurvirostra americana</i>	LC	Stable	2004-2006	San Francisco Bay	USA	PAF	PB/B	blood, liver, kidney, muscle, feather	Hg	Eagles-Smith et al 2009
				2005-2006	San Francisco Bay	USA	PAF	B	blood	Hg	Ackerman et al, 2007
American Golden Plover	<i>Pluvialis dominica</i>	LC	Decreasing	2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba, Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				1990-1992	Cape Churchill, Quill Lakes, Bay of Fundy, Tofino, Boundary Bay	Canada	AAF	B	carcass	HCB, Σ CBz, β -HCH, Σ HCH, oxychl, trans-nonachl, HE, Σ CHL, p,p' DDE, Σ DDT, Dieldrin, Σ Merex, Σ PCB42, Hg, Se, Cd, As	Braune et al 2009
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
American Oystercatcher	<i>Haematopus palliatus</i>	LC	Stable	2007 - 2015	Southern coast of Brazil	Brazil	mutiple	W	gastrointestinal tract	plastic fragment	Rossi et al 2019
American Woodcock	<i>Scolopax minor</i>	LC	Decreasing	2002-2003	Connecticut	USA		Local/Sto	liver, kidney, muscle, bone	As, Cd, Cr, Hg, Pb, Se	Hiller et al 2011
Baird's Sandpiper	<i>Calidris bairdii</i>	LC	Stable	2008–2009	Barrow, Alaska	USA	mutiple	B	blood, feather	Hg	Perkins et al 2016
Black-tailed Godwit	<i>Limosa limosa</i>	NT	Decreasing	2007-2011	Pertuis, Charentais, Atlantic coast	France	EAF	W/Sto	liver, kidney, muscle, feather	Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Zn	Lucia et al 2012a

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Bar-tailed Godwit	<i>Limosa lapponica</i>	NT	Decreasing	2002-2004	polder Blokland and polder Zeevang	The Netherlands	EAF	B	egg, feather	Cd, Cr,Cu, Hg, Ni, Pb, Zn	Roodbergen et al 2009
				2013-2015	The Tejo estuary; The Banc d'Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
				1996–1997	Lake Baikal	Russia	EAAF	B	whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				2007-2010	Pertuis Charentais, Atlantic coast	France	EAF		liver, kidney, muscle, feather	Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Zn	Lucia et al 2012b
				2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				1997	Con Lu Island	Vietnam			whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2013-2015	The Tejo estuary; The Banc d'Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Black Oystercatcher	<i>Haematopus bachmani</i>	LC	Unknown	2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				1992-1993	Prince William Sound, Alask	USA	mutiple	B	behavior	T/V Exxon Valdez oil spill	Andres 1999
				1988,1989-1991	Prince William Sound, Alask	USA	mutiple	B	behavior	T/V Exxon Valdez oil spill	Murphy et al 1997
Black Turnstone	<i>Arenaria melanocephala</i>	LC	Stable	2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
Black-bellied Plover	<i>Pluvialis squatarola</i>	LC	Decreasing	2008	Nunavut	Canada	AAF	B	blood, egg, feather, RBC	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2010
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				1994	San Diego area	USA	mutiple	W	liver	Al, As, Cd, Cu, Fe, Hg, Mg, Mn, Mo, Ni, Se, Zn	Hui 1998

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordan, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2011
				1990-1992	Cape Churchill, Quill Lakes, Bay of Fundy, Tofino, Boundary Bay	Canada	AAF	B	carcass	HCB, Σ CBz, β -HCH, Σ HCH,oxychl,trans-nonachl,HE, Σ CHL,p,p' DDE, Σ DDT,Dieldrin, Σ Mirex, Σ PCB42, Hg, Se, Cd, As	Braune et al 2009
				1981	Western Washington	USA	PAF	W	carcass	p,p'-DDE,p,p'-TDE, p,p'-DDT, Σ DDT, PCBs, HCB, Chlordane isomers, Dieldrin, Heptachlor epoxide	Schick et al 1987
Black-necked Stilt	<i>Himantopus mexicanus</i>	LC	Increasing	2011	Nunavut	Canada	AAF	B	egg	Hg	McCloskey et al 2013
				2004-2006	San Francisco Bay	USA	PAF	PB/B	blood, liver, kidney,muscle, feather	Hg	Eagles-Smith et al 2009
				1985-1988	the Grasslands Watershed of Central California	USA	PAF	B, W	liver	Se, B	Paveglio et al 1992
				1986-2005	the Grasslands Watershed of Central California	USA	PAF	W	liver	Se	Paveglio et al 2007

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Black-winged Stilt	<i>Himantopus himantopus</i>	LC	Increasing	2005-2006	San Francisco Bay, CA	USA	PAF	PB	blood	Hg	Ackerman et al, 2007
				2006-2007	Persian Gulf	Iran	M/BSF	Local	feather,kidney, liver and muscle	Hg	Zamani-Ahmadmahmoodi et al 2010
				1997-1999	mutiple sites	Portugal	EAF	B	feather,egg, liver	Hg	Tavares et al 2004
				2000-2003	the Western Wadden Sea	Portugal	EAF	B	feather	Hg	Tavares et al 2009
				na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni, Zn	Pandiyan et al 2020
Broad-billed Sandpiper	<i>Calidris falcinellus</i>	LC	Decreasing	1995	Cuddalore, Parangipettai, Pudukottai and Mandapan in southern India	India	EAAF	Local	whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Buff-breasted Sandpiper	<i>Calidris subruficollis</i>	NT	Decreasing	2011	Lagoa do Peixe National Park	Brazil	mutiple	W	feather	Cd, Co, Cu, Mn, Pb, Zn	Scherer et al 2015
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ-PCB, Σ-DDT	Saalfeld et al 2016
Common Sandpiper	<i>Actitis hypoleucos</i>	LC	Decreasing	2011	Zayanderud Dam	Iran	M/BSF	W/local	liver, kidney, feathers	Hg	Malekian et al 2015

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni, Zn	Pandiyam et al 2020
				1995, 1998	Cuddalore, Parangipettai, Pudukottai and Mandapan in southern India	India	EAAF		whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
Common Greenshank	<i>Tringa nebularia</i>	LC	Stable	1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Common Snipe	<i>Gallinago gallinago</i>	LC	Decreasing	1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Curlew Sandpiper	<i>Calidris ferruginea</i>	NT	Decreasing	1981-1983	the Baltic Sea	Sweden	EAF	Stagging	liver, kidney	Ca, Cd, Co, Cu, Fe, Mg, Mn, Pb, V and Zn	Blomqvist et al 1987
				na	Point Calimere Wildlife Sanctuary (PWLS) and Pichavaram Mangrove Forest (PMF)	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyam et al 2020
				1995	Cuddalore, Parangipettai, Pudukottai and Mandapan in southern India	India	EAAF		whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
				2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Double-banded Plover	<i>Charadrius bicinctus</i>	NT	Decreasing	2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Dunlin	<i>Calidris alpina</i>	LC	Decreasing	2008–2009	North American Arctic	USA	mutiple	B	blood, feather	Hg	Perkins et al 2016
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	feather, liver	Zn, Mn, Cu, Pb, Cd	Kim & Koo, 2008
				2003-2011	the Pertuis Charentais	France	EAF	W & stop	liver, kidney, muscle, feather	Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Zn	Lucia et al 2014
				2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2011
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle, bone	Pb, Cd	Kim et al 2007a
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle, bone	Cd,Cu, Mn, Pb, Zn	Kim et al 2007b
				1990-1992	Cape Churchill, Quill Lakes, Bay of Fundy, Tofino, Boundary Bay	Canada	EAF	B/Sto/W	carcass	HCB, Σ CBz, β -HCH, Σ HCH,oxychl,trans-nonachl,HE, Σ CHL,p,p' DDE, Σ DDT,Dieldrin, Σ Mirex, Σ PCB42, Hg, Se, Cd, As	Braune et al 2009

Species	<i>Scientific name</i>	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1979-1981	the Mersey Estuary	UK	EAF, M/BSF	W	brain, bone, kidney, Pb liver, muscle		Bull et al 1983
				1979-1980	the Bristol channel	UK	EAF, M/BSF	W	kidney	Cd	Ferns & Anderson, 1993
				1979-1982, 1991	the Bristol Channel	UK	mutiple	B	beak, gut, kidney, leg, liver, muscle, sternum, bone	Pb	Ferns & Anderson, 1997
				1981	Western Washington	USA	PAF	W	carcass	p,p'-DDE, p,p'-TDE, p,p'-DDT, Σ DDT, PCBs, HCB, Chlordane isomers, Dieldrin, Heptachlor epoxide	Schick et al 1987
				2009- 2011	Chongming Island, shanghai	China	EAAF	Sto	muscle	BDE28, BDE47, BDE99, BDE100, BDE153, BDE154, BDE183, BDE209, Sum-PBDEs	Huang et al 2013
				2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/Sto	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014
				1990	California	USA	mutiple	Sto	stomach contents	Strychnine	Warnock & Schwarzbach, 1995
	na				Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2013-2015	The Tejo estuary; The Banc d'Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
	<i>Calidris alpina pacifica</i>			2007	the Fraser River Delta, British Columbia and Skagit Bay, Washington	Canada, USA		Sto	feather,kidney	Cd, Cu, Pb, Zn, Hg, Se	St Clair et al 2015
Eurasian Oystercatcher	<i>Haematopus ostralegus</i>	NT	Decreasing	1987-1988	the mainland coast of the Dutch Wadden Sea	The Netherlands	EAF	B	Se-dependent enzyme glutathione peroxidase	Se	Goede & Wolterbeek, 1994
					the coast of the Dutch Wadden Sea	The Netherlands			Se distribution in the flight feathers	Se	Goede, 1991
				1981-1982	EI Salvador Copper Mine	Chile	PAF	W	liver, stomach	Cd, Cu	Vermeer and Castilla 1991
				1987	The Wadden Sea/Wangerooge	Germany	EAF	W	kidney, liver, feather,	Cd	Stock et al 2007
				1986	Dutch Wadden Sea	The Netherlands	EAF	B	egg, yolk, white, plasma, red blood cells	Se	Goede 1993a

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1986-1990	Dutch Wadden Sea	The Netherlands	EAF	B	plasma, RBC, liver, Se lung, kidney, spleen, gallbladder, testis, salt gland, sm. intestine, heart, pancreas, gizzard, preen gland, brain, breast muscle		Goede 1993b
				1986-1987	the central and eastern Oosterschelde and the western Wadden Sea	The Netherlands		W	liver lipid	PCBs	Lambeck et al. 1991
Eurasian Curlew	<i>Numenius arquata</i>	NT	Decreasing	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
Eurasian Golden Plover	<i>Pluvialis apricaria</i>	LC	Increasing	1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Great Knot	<i>Calidris tenuirostris</i>	EN	Decreasing	1994-1995	Yeongjong Island	Korea	EAAF	Sto	feather, liver	Zn, Mn, Cu, Pb, Cd	Kim & Koo, 2008
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Pb, Cd	Kim et al 2007a
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Cd,Cu, Mn, Pb, Zn	Kim et al 2007b

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
					Chongming Island, Shanghai	China	EAAF	Sto	muscle	BDE28, BDE47, BDE99, BDE100, BDE153, BDE154, BDE183, BDE209, Sum-PBDEs	Huang et al 2013
				2008-2012	Yellow Sea and Bohai Sea	China	EAAF	Sto	adipose tissue, muscle	α -HCH, β -HCH, γ -HCH, δ -HCH, Sum-HCHs, Heptachlor, Aldrin, Heptachlor epoxide, <i>g</i> -Chlordane, Endosulfan I, <i>a</i> -Chlordane, <i>p</i> , <i>p'</i> -DDE, Dieldrin, Endrin, Endosulfan II, <i>p</i> , <i>p'</i> -DDD, Endrin aldehyde, Endosulfan sulfate, Methoxychlor	Jin et al 2017
				1999–2000	Okgu Mudflat,	Korea	EAAF	Sto	liver	Fe,Zn,Cu,Pb,Cd	Kim et al 2009
				1999-2000	Okgu Mudflat	Korea	EAAF	Sto	bone, liver, kidney	Zn,Cu,Pb,Cd	Kim & Koo, 2010
				1999-2000	Okgu Mudflat	Korea	EAAF	Sto	liver, feather	Fe, Zn, Cu, Pb, Cd	Kim & Koo, 2012
				1999–2000	Okgu Mudflat	Korea	EAAF	Sto	liver, feather	Fe,Zn,Cu,Pb,Cd	Kim & Oh, 2018
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
Greater Yellowleg	<i>Tringa melanoleuca</i>	LC	Stable	1990-1992	Cape Churchill, Quill Lakes, Bay of Fundy, Tofino, Boundary Bay	Canada	EAF	B/Sto/W	Carcass	HCB, Σ CBz, β -HCH, Σ HCH,oxychl,trans-nonachl,HE, Σ CHL, <i>p</i> , <i>p'</i> -DDE, Σ DDT,Dieldrin, Σ Mirex, Σ PCB42, Hg, Se, Cd, As	Braune et al 2009
Green Sandpiper	<i>Tringa ochropus</i>	LC	Increasing	2011	Zayanderud Dam	Iran	M/BSF	W/local	liver, kidneys, and tail and breast and tail feathers	Hg	Malekian et al 2015

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Grey Plover	<i>Pluvialis squatarola</i>	LC	Decreasing	1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagos archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
Hudsonian Godwit	<i>Limosa haemastica</i>	LC	Decreasing	1990-1992	Cape Churchill, Quill Lakes, Bay of Fundy, Tofino, Boundary Bay	Canada	EAF	B/Sto/W	Carcass	HCB,ΣCBz,β-HCH, ΣHCH,oxychl,trans-nonachl,HE,ΣCHL,p,p' DDE,ΣDDT,Dieldrin,ΣMir ex,ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				2017-2018	Pacific Patagonian coasts	Chile	PAF	Sto	fecal	ARB and ARGs	Navedo et al 2021
				2005	Northern Ontario	Canada			breast muscle, skin, abdominal fat	DDTs, PCBs, CHL	Tsuji et al 2008
Hudsonian Whimbrel	<i>Numenius hudsonicus</i>	NT	Decreasing	1981-1982	EI Salvador Copper Mine	Chile	PAF	W	liver, stomach	Cd, Cu	Vermeer and Castilla 1991
Kentish Plover	<i>Charadrius alexandrinus</i>	LC	Decreasing	2016	coastal/inland	China	EAAF	B	blood, feather	Hg	Su et al 2018
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	feather, liver	Zn, Mn, Cu, Pb, Cd	Kim & Koo, 2008
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Pb, Cd	Kim et al 2007a
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Cd,Cu, Mn, Pb, Zn	Kim et al 2007b
				2018	Delaware Bay	USA	AAF	Sto	feather	As,Cd,Co, Cr, Cu, Ni, Pb,Hg,Se, Zn, V	Picone et al 2019
				2015-2017	Western Bohai Bay	China	EAAF	Sto	egg, muscle and liver	DDTs, p,p'-DDE, HCHs, PCBs, PBDEs, HCB, PCDD/Fs	Wang et al 2019
				2015-2016	Western Bohai Bay	China	EAAF	Sto	egg, muscle and liver	PCDD/Fs, PCBs, PBDEs, OCPs,HCB, Mirex, PeCB, HCHs	Zheng et al 2018

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2009-2011	Chongming Island, Shanghai	China	EAAF	Sto	muscle	BDE28, BDE47, BDE99, BDE100, BDE153, BDE154, BDE183, BDE209, Sum-PBDEs	Huang et al 2013
				2000-2003	the Western Wadden Sea	Portugal	EAF	B	feather	Hg	Tavares et al 2009
				2009-2013	the Iberian Peninsula	Spain	EAF	B	feather	Al, Cr, Mn, Cu, Zn, As, Se, Cd, Hg, Pb	Vidal et al 2021
				na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				1995-1998	Cuddalore, Parangipettai, Pudukottai and Mandapan in southern India; coastal areas of Caratagan in Philippines	India, Philippines	CAF	Local	Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
Killdeer	<i>Charadrius vociferus</i>	LC	Decreasing	1990	California	USA	mutiple	Sto	stomach contents	Strychnine	Warnock & Schwarzbach, 1995
				2007	Illinois	USA	NA	Sto	liver	Hg	Levengood et al 2014
				2020	Saskatchewan	Canada	NA	NA	egg	PCB-126	Lunny et al 2020
Least Sandpiper	<i>Calidris minutilla</i>	LC	Decreasing	2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/stopover	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Lesser Sandplover	<i>Charadrius mongolus</i>	LC	Unknown	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
Lesser Yellowleg	<i>Tringa flavipes</i>	LC	Decreasing	1991-1993	Bay of Fundy, Cape Churchill, Araya Peninsula, Delaware Bay	Canada, USA	PAF, EAAF, MAF	B/Sto/W	carcass, liver, kidney	Cd, Se, Hg, Organochlorines	Pratte et al 2020
				1990-1992	mutiple sites	Canada	AAF	Sto	carcass	HCB,ΣCBz,β-HCH, ΣHCH,oxychl,trans-nonachl,HE,ΣCHL,p,p' DDE,ΣDDT,Dieldrin,ΣMir ex,ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
Little ringed Plover	<i>Charadrius dubius</i>	LC	Stable	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1995	Cuddalore, Parangipettai, Pudukottai and Mandapan in southern India; coastal areas of Caratagan in Philippines	India, Philippines	CAF	Local	Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
Little Stint	<i>Calidris minuta</i>	LC	Increasing	na	Point Calimere Wildlife Sanctuary (PWLS) and Pichavaram Mangrove Forest (PMF)	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Long-billed Curlew	<i>Numenius americanus</i>	LC	Decreasing	1978-1979, 1981-1983	Umatilla and Morrow counties, Oregon	USA	multiple	B	eggshell, brain	DDE, DDT, Dieldrin, Heptachlor epoxide, Oxychlorodane, CCH, TNCH, CNCH, TCH, Endrin, PCBs	Blus et al 1985
Long-billed Dowitcher	<i>Limnodromus scolopaceus</i>	LC	Unknown	2008-2009	North American Arctic	USA	multiple	B/Staging	blood, feather	Hg	Perkins et al 2016
				2012-2013	North American Arctic	USA	multiple	B	blood, feather	Hg	Stenhouse et al 2014
				1991-1992	San Francisco Bay	USA	PAF	W	liver, muscle and fat	Ag, As, Cd, Cr, Cu, Hg, Mn, Ni, Se, V, Zn, OCPs, PCBs, DDTs and HCBs	Hui et al 2001
				1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB, ΣCBz, β-HCH, ΣHCH, oxychl, trans-nonachl, HE, ΣCHL, p,p' DDE, ΣDDT, Dieldrin, ΣMir ex, ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				2002-2004	Alaska	USA	multiple	B	egg	Al, As, B, Ba, Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorodane, p,p'-DDE, p,p'-DDT, Σ-PCB, Σ-DDT	Saalfeld et al 2016
Marbled Godwit	<i>Limosa fedoa</i>	LC	Decreasing	1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB, ΣCBz, β-HCH, ΣHCH, oxychl, trans-nonachl, HE, ΣCHL, p,p' DDE, ΣDDT, Dieldrin, ΣMir ex, ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				2005	Northern Ontario	Canada				breast muscle, skin, abdominal fat	DDTs, PCBs, CHL

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Marsh Sandpiper	<i>Tringa stagnatilis</i>	LC	Decreasing	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyani et al 2020
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
Mongolian Plover	<i>Charadrius mongolus</i>	LC	Unknown	1996–1997	Lake Baikal	Russia		B	Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	feather, liver	Zn, Mn, Cu, Pb, Cd	Kim & Koo, 2008
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Pb, Cd	Kim et al 2007a
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Cd,Cu, Mn, Pb, Zn	Kim et al 2007b
					Chongming Island, shanghai	China	EAAF	Sto	muscle	BDE28, BDE47, BDE99, BDE100, BDE153, BDE154, BDE183, BDE209, Sum-PBDEs	Huang et al 2013
Northern Lapwing	<i>Vanellus vanellus</i>	NT	Decreasing	1994	Caratagan	Philippines	EAAF	Sto& W	whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
Northern Lapwing	<i>Vanellus vanellus</i>	NT	Decreasing	1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
				1996-1997	Lake Baikal	Russia		B	Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Pacific Golden Plover	<i>Pluvialis fulva</i>	LC	Decreasing	2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
Painted Snipe	<i>Rostratula benghalensis</i>	LC	Decreasing	1994	Caratagan	Philippines	local	Local	whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
Pectoral Sandpiper	<i>Calidris melanotos</i>	LC	Stable	2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
Pied Oystercatcher	<i>Haematopus longirostris</i>	LC	Unknown	2007 2016–2019	Illinois tidal habitats on the Australian coast	USA Australia	NA EAAF	Sto W/local	liver cloacal swab	Hg AMR	Levengood et al 2014 Smith et al 2021
Pied Avocet	<i>Recurvirostra avosetta</i>	LC	Unknown	2000-2003	the Western Wadden Sea	Portugal	EAF	B	feather	Hg	Tavares et al 2009
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea- Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
Piping Plover	<i>Charadrius melodus</i>	NT	Increasing	2010	the Gulf of Mexico and Atlantic southeast	USA	AAF	Sto	behavior	Oil spill	Gibson et al 2017

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Red-capped Plover	<i>Charadrius ruficapillus</i>	LC	Unknown	2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Red Knot	<i>Calidris alba</i>	LC	Unknown	2015-2017	Gulf of Mexico	Canada	mutiple	Stagging	Plasma Metabolite Concentrations, body mass and staging duration	PAH	Bianchini and Morrissey, 2018 a
				1991-1992	Delaware Bay	USA	AAF	W	feather	Pb,Cd,Hg,Se,Mn,Cr	Burger et al 1993
				2007-2011	Pertuis Charentais	France	EAF	Sto	liver, kidney, muscle	Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Zn	Lucia et al 2012
				NA	Delaware Bay	USA	mutiple	Sto	blood, feather, egg, muscle	Cd, Pb, Hg, Se	Burger et al 2019b
				2011-2012	Delaware Bay	USA	AAF	Sto	blood, feather	Pb,Cd,Hg,As,Cr	Tsipoura et al 2017
				1991-1992, 1995, 2011–2012	Delaware Bay	USA	AAF	Sto	feather	As,Cd,Cr,Pb,Hg,Se	Burger et al 2015
				2008-2012	Yellow Sea and Bohai Sea	China	EAAF	Sto	adipose tissue, muscle	α -HCH, β -HCH, γ -HCH, δ -HCH, Sum-HCHs, Heptachlor, Aldrin, Heptachlor epoxide, g-Chlordane, Endosulfan I, a-Chlordane, p, p'-DDE, Dieldrin, Endrin, Endosulfan II, p, p'-DDD, Endrin aldehyde, Endosulfan sulfate, Methoxychlor	Jin et al 2017
				2011-2012	Delaware Bay	USA	mutiple	Sto	blood, feather	As, Cd, Cr, Pb, Hg	Burger et al 2017
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2011-2012	Delaware Bay and Barnegat Bay	USA	AAF	Sto	blood	Se, Hg and Se: Hg	Burger & Gochfeld, 2021
Red Phalarope	<i>Phalaropus fulicarius</i>	LC	Unknown	2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2011
Red-necked Phalarope	<i>Phalaropus lobatus</i>	LC	Decreasing	2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
Red-necked Stint	<i>Calidris ruficollis</i>	NT	Decreasing	1999-2000	Okgu Mudflat	Korea	EAAF	Sto	bone, liver, kidney	Zn,Cu,Pb,Cd	Kim & Koo, 2010
				1999–2000	Okgu Mudflat,	Korea	EAAF	Sto	liver	Fe,Zn,Cu,Pb,Cd	Kim et al 2009
				1999-2000	Okgu Mudflat	Korea	EAAF	Sto	liver, feather	Fe, Zn, Cu, Pb, Cd	Kim & Koo, 2012

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Red Shank	<i>Tringa totanus</i>	LC	Unknown	2003-2011	the Pertuis Charentais	France	EAF	W & stop	liver, kidney, muscle, feather	Ag, As, Cd, Co, Cr, Cu, Fe, Hg, Mn, Ni, Pb, Se, Zn	Lucia et al 2014
				na	Point Calimere Wildlife Sanctuary (PWLS) and Pichavaram Mangrove Forest (PMF)	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
Red-wattled Plover	<i>Hoplopterus indicus</i>	LC	Unknown	2006-2007	Persian Gulf	Iran	M/BSF	Local	feather, kidney, liver and muscle	Hg	Zamani-Ahmadm Mahmoodi et al 2010
Ringed Plover	<i>Charadrius hiaticula</i>	LC	Decreasing	2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
Rock sandpiper	<i>Calidris ptilocnemis</i>	LC	Decreasing	2008–2009	Yukon Delta NWR	USA	mutiple	B	blood	Hg	Perkins et al 2016

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
Ruddy Turnstone	<i>Arenaria interpres</i>	LC	decreasing	2008	Nunavut	Canada	AAF	B	blood, egg, feather, RBC	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2010
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2011
				1994	Caratagan	Philippines	EAAF	W/Sto	whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
				2011-2012	Delaware Bay and Barnegat Bay	USA	AAF	Sto	blood	Se, Hg and Se: Hg	Burger & Gochfeld, 2021
				2011 2016–2019	Nunavut tidal habitats on the Australian coast	Canada Australia	AAF EAAF	B W/local	egg cloacal swab	Hg AMR	McCloskey et al 2013 Smith et al 2021
				NA	Delaware Bay	USA	mutiple	Sto	blood, feather, egg, muscle	Cd, Pb, Hg, Se	Burger et al 2019b

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Ruff	<i>Philomachus pugnax</i>	LC	decreasing	1996-1997	Lake Baikal	Russia		B	Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisie et al 2003
				1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Sanderling	<i>Calidris alba</i>	LC	Unknown	2016	Saskatchewan	Canada	NA	Stagging	refuelling physiology	PAH	Bianchini and Morrissey, 2018 b
				2015-2017	Gulf of Mexico	Canada	mutiple	Stagging	plasma metabolite concentrations, body mass and stagging duration	PAH	Bianchini and Morrissey, 2018 a
				2021	Saskatchewan	Canada	NA	pre migration	liver	PAH	Bianchini et al 2021
				1991-1992	Delaware Bay	USA	AAF	W	feather	Pb,Cd,Hg,Se,Mn,Cr	Burger et al 1993
				1981-1982	EI Salvador Copper Mine	Chile	PAF	W	liver, stomach	Cd, Cu	Vermeer and Castilla 1991
				NA	Delaware Bay	USA	mutiple	Sto	blood, feather, egg, muscle	Cd, Pb, Hg, Se	Burger et al 2019b
				1991-1992,1995,2011-2012	Delaware Bay	USA	AAF	Sto	feather	As,Cd,Cr,Pb,Hg,Se	Burger et al 2015
				2011-2012	Delaware Bay	USA	AAF	Sto	blood, feather	Pb,Cd,Hg,As,Cr	Tsipoura et al 2017
				1981	Western Washington	USA	PAF	W	carcass	p,p'-DDE,p,p'-TDE, p,p'-DDT, Σ DDT, PCBs, HCB, Chlordane isomers, Dieldrin, Heptachlor epoxide	Schick et al 1987
1996	Delaware Bay	USA	AAF	Sto	foraging behavior	Oil spill	Burger et al 1997				
2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/stopover	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014				
2011-2012	Delaware Bay	USA	mutiple	Sto	blood, feather	As, Cd, Cr, Pb, Hg	Burger et al 2017				
2013-2015	The Tejo estuary; The Banc d' Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017				

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2011-2012	Delaware Bay and Barnegat Bay	USA	AAF	Sto	blood	Se, Hg and Se: Hg	Burger & Gochfeld, 2021
				2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Semipalmated Plover	<i>Charadrius semipalmatus</i>	LC	Stable	2008	Nunavut	Canada	AAF	B	blood, egg, feather, RBC	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Ti, V, Zn	Hargreaves et al 2010
				1991-1993	Bay of Fundy	Canada	mutiple	B/Sto/W	liver, kidney	Cd, Se, Hg, Organochlorines	Pratte et al 2020
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba, Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordan, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Ti, V, Zn	Hargreaves et al 2011
				1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB, Σ CBz, β -HCH, Σ HCH, oxychl, trans-nonachl, HE, Σ CHL, p,p'-DDE, Σ DDT, Dieldrin, Σ Mir ex, Σ PCB42, Hg, Se, Cd, As	Braune et al 2009
				1989-1990	Bay of Fundy	Canada	AAF	B	adipose tissue	Squalene, PCBs, DDTs, DDEs	Napolitano et al 1992
				1996	Delaware Bay	USA	AAF	Sto	foraging behavior	Oil spill	Burger et al 1997

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/stopover	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014
Semipalmated Sandpiper	<i>Calidris pusilla</i>	NT	Decreasing	2011	Nunavut	Canada	AAF	B	egg	Hg	McCloskey et al 2013
				2008–2009	Barrow, Alaska	USA	mutiple	B	blood	Hg	Perkins et al 2016
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				2013	Delaware Bay and Suriname	Suriname, USA	AAF	W/S	blood	Hg, Pb, Cd, Co, As and Se	Burger et al 2018
				1991-1992	Delaware Bay	USA	AAF	W	feather	Pb,Cd,Hg,Se,Mn,Cr	Burger et al 1993
				1991-1993	Bay of Fundy	Canada	mutiple	B/Sto/W	liver, kidney	Cd, Se, Hg, Organochlorines	Pratte et al 2020
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				2011-2016	Delaware Bay	USA, Suriname and Brazil		W	blood	As,Cd,Cr,Co,Pb,Hg,Se	Burger et al 2019a
				NA	Delaware Bay	USA	mutiple	Sto	blood, feather, egg, muscle	Cd, Pb, Hg, Se	Burger et al 2019b
				2011-2012	Delaware Bay	USA	AAF	Sto	blood, feather	Pb,Cd,Hg,As,Cr	Tsipoura et al 2017
1991-1992,1995, 2011–2012	Delaware Bay	USA	AAF	Sto	feather	As,Cd,Cr,Pb,Hg,Se	Burger et al 2015				

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB,ΣCBz,β-HCH, ΣHCH,oxychl,trans-nonachl,HE,ΣCHL,p,p' DDE,ΣDDT,Dieldrin,ΣMir ex,ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				2011	Lagoa do Peixe National Park	Brazil	mutiple	W	feather	Cd, Co, Cu, Mn, Pb, Zn	Scherer et al 2011
				1989-1990	Bay of Fundy	Canada	AAF	B	adipose tissue	Squalene, PCBs, DDTs, DDEs	Napolitano et al 1992
				2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/stopover	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014
				NA	Delaware Bay	USA	mutiple	Sto	brain, feather, fat, liver, muscle	As,Cd,Cr,Pb,Mn,Hg,Se	Burger et al 2014
				2011-2012	Delaware Bay	USA	mutiple	Sto	blood, feather	As, Cd, Cr, Pb, Hg	Burger et al 2017
				2011-2012	Delaware Bay and Barnegat Bay	USA	AAF	Sto	blood	Se, Hg and Se: Hg	Burger & Gochfeld, 2021
Sharp-tailed Sandpiper	<i>Calidris acuminata</i>	LC	Stable		Chongming Island, shanghai	China	EAAF	Sto	muscle	BDE28, BDE47, BDE99, BDE100, BDE153, BDE154, BDE183, BDE209, Sum-PBDEs	Huang et al 2013
				2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Short-billed Dowitcher	<i>Limnodromus griseus</i>	LC	Decreasing	1991-1993	Bay of Fundy	Canada	mutiple	B/Sto/W	liver, kidney	Cd, Se, Hg, Organochlorines	Pratte et al 2020
				1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB,ΣCBz,β-HCH, ΣHCH,oxychl,trans-nonachl,HE,ΣCHL,p,p' DDE,ΣDDT,Dieldrin,ΣMir ex,ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				1989-1990	Bay of Fundy	Canada	AAF	B	adipose tissue	Squalene, PCBs, DDTs, DDEs	Napolitano et al 1992

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
Sooty Oystercatcher	<i>Haematopus fuliginosus</i>	LC	Stable	2016–2019	tidal habitats on the Australian coast	Australia	EAAF	W/local	cloacal swab	AMR	Smith et al 2021
Spotted Redshank	<i>Tringa erythropus</i>	LC	Stable	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
Spotted Sandpiper	<i>Actitis macularia</i>	LC	Decreasing	2002	The Elk River	Canada	mutiple	B	egg	Se	Harding et al 2005
Spur-winged Lapwing	<i>Vanellus spinosus</i>	LC	Increasing	2015	the Mediterranean Sea coast of Turkey	Turkey	M/BSF	B	feather	CHLs, HCHs, DDTs, HCCPDs and PCBs	Arıkan et al 2018
Stilt sandpiper	<i>Calidris himantopus</i>	LC	Increasing	2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlorane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
Terminck's Stint		LC	Unknown	na	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
Terek Sandpiper	<i>Xenus cinereus</i>	LC	Decreasing	1994-1995	Yeongjong Island	Korea	EAAF	Sto	feather, liver	Zn, Mn, Cu, Pb, Cd	Kim & Koo, 2008
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Pb, Cd	Kim et al 2007a

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1994-1995	Yeongjong Island	Korea	EAAF	Sto	liver, kidney, muscle and bone	Cd,Cu, Mn, Pb, Zn	Kim et al 2007b
				1999–2000	Okgu Mudflat,	Korea	EAAF	Sto	liver	Fe,Zn,Cu,Pb,Cd	Kim et al 2009
				1999-2000	Okgu Mudflat	Korea	EAAF	Sto	bone, liver, kidney	Zn,Cu,Pb,Cd	Kim & Koo, 2010
				1999-2000	Okgu Mudflat	Korea	EAAF	Sto	liver, feather	Fe, Zn, Cu, Pb, Cd	Kim & Koo, 2012
				2008	Wadden Sea	Germany	EAF	B	Serum, feather	PCBs,OxC,HCB, α -, β -, γ -HCHs,DDT,PBDEs	Schwemmer et al 2015
				1995		India			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Kunisue et al 2003
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
Western Sandpiper	<i>Calidris mauri</i>	LC	Decreasing	2013	Ontario	Canada	na	Migration	flight energy expenditure	Oil spill	Maggini et al 2017
				2012-2013	North American Arctic	USA	mutiple	B	blood, feather	Hg	Stenhouse et al 2014
				1991-1992	San Francisco Bay	USA	PAF	W	liver	Ag, As, Cd, Cr, Cu, Hg, Mn, Ni, Se, V, Zn	Hui et al 2001
				2010-2011	coast of Sinaloa	Mexico	mutiple	W, pre migration	muscle, liver	Hg	Fernández et al 2019
				2002-2004	Alaska	USA	mutiple	B	egg	Al, As, B, Ba,Be, Cr, Cu, Cd, Fe, Hg, Mg, Mn, Mo, Ni, Pb, Se, Sr, Zn, V, Endosulfan I, Hexachlorobenzene, Heptachlor epoxide, Oxychlordane, p,p'-DDE, p,p'-DDT, Σ -PCB, Σ -DDT	Saalfeld et al 2016
				1981	Western Washington	USA	PAF	W	carcass	p,p'-DDE,p,p'-TDE, p,p'-DDT, Σ DDT, PCBs, HCB, Chlordane isomers, Dieldrin, Heptachlor epoxide	Schick et al 1987

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/stopover	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014
				1990	Nueces Bay, Texas	USA	multiple	W	Pooled stomach contents and carcasses	organochlorine pesticides and metabolites, PCBs	Rattner et al 1995
				2013	Ontario	Canada	NA	NA	thermoregulatory ability	Mississippi Canyon 252 (MC252) oil	Bursian et al 2017
				2013	Ontario	Canada	NA	NA	thermoregulatory ability	Mississippi Canyon 252 (MC252) oil	Maggini et al 2017
				1995-1996	the Fraser Estuary, British Columbia, Canada; Playa el Agallito, Chitre, Panama	Canada	PAF	Sto & W	liver, kidney, bone	Cd	McFarland et al 2002
Whimbrel	<i>Numenius phaeopus</i>	LC	Decreasing	1990-1992	Quill Lakes	Canada	EAF	Sto	Carcass	HCB, ΣCBz, β-HCH, ΣHCH, oxychl, trans-nonachl, HE, ΣCHL, p,p' DDE, ΣDDT, Dieldrin, ΣMir ex, ΣPCB42, Hg, Se, Cd, As	Braune et al 2009
				1997	Con Lu Island	Vietnam			Whole body	CHLs, HCHs, DDTs, HCB and PCBs	Minh et al 2002
				2013-2015	The Tejo estuary; The Banc d'Arguin; The Bijagós archipelago	Mauritania, Guinea-Bissau, Portugal		Sto	faeces	Plastic and microfibers	Lourenço et al 2017
				2017	Yongxing Island, Xisha archipelago	China	EAAF	W	Esophagus, stomach and intestine	plastic debris	Zhu et al 2019

Species	Scientific name	Conservation status	Population trend	Sampling period	Location	Country	Flyway	Life stage	Tissue sampled/endpoint	Types of pollutants	Reference
				1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985
White-faced Plover	<i>Charadrius dealbatus</i>	DD	Unknown	2016	coastline/inland	China	EAAF	B	blood, feather	Hg	Su et al 2018
White-rumped Sandpiper	<i>Calidris fuscicollis</i>	LC	Decreasing	2008	Nunavut	Canada	AAF	B	blood, egg, feather	As, Be, Cd, Co, Cr, Cu, Fe, Hg, Mn, Mo, Ni, Pb, Sb, Se, Tl, V, Zn	Hargreaves et al 2011
White-tailed Plover	<i>Vanellus leucurus</i>	LC	Unknown	2011	Nunavut	Canada	AAF	B	egg	Hg	McCloskey et al 2013
Willet	<i>Tringa semipalmata</i>	LC	Stable	2006-2007	Persian Gulf	Iran	M/BSF	Local	feather, kidney, liver and muscle	Hg	Zamani-Ahmadmahmoodi et al 2010
Wilson's Plover	<i>Charadrius wilsonia</i>	LC	Decreasing	1994	San Diego area	USA	mutiple	W	liver	Al, As, Cd, Cu, Fe, Hg, Mg, Mn, Mo, Ni, Se, Zn	Hui 1998
				2010-2012	The coastline of the Northern Gulf of Mexico	Mexico		W/Sto	blood and plasma, Proportions of total oiled shorebirds observed	Deepwater Horizon oil spill	Henkel et al 2014
Wood Sandpiper	<i>Tringa glareola</i>	LC	Stable	NA	Point Calimere Wildlife Sanctuary and Pichavaram Mangrove Forest	India	CAF	W	feather	Cu, Cr, Co, Pb, Hg, Ni and Zn	Pandiyan et al 2020
				1976-1977	the county of Gallivare, Norrbotten	Sweden		B	muscle	DDTs, PCBs, Hg	Lindberg et al 1985

Data source

DD

LC

NT

EN

PB

B

Sto

W

[BirdLife International \(2022\) IUCN Red List for birds. Downloaded from http://www.birdlife.org on 20/05/2022.](http://www.birdlife.org)

Data Deficient

Least Concern

Near Threatened

Endangered

Pre breeding

Breeding

Stopover

Wintering