

**Evaluating the Clinical Care of Traumatic Brain Injury
Patients and Identifying Opportunities for Quality
Improvement in Neurosurgery at Mulago National Referral
Hospital in Kampala Uganda**

by

Benjamin Jyhhan Kuo

Duke Global Health Institute
Duke University

Date: _____
Approved:

Michael M. Haglund, Supervisor

Henry E. Rice

Joao Ricardo N. Vissoci

Thesis submitted in partial fulfillment of
the requirements for the degree of
Master of Science in the Duke Global Health Institute
in the Graduate School of Duke University

2017

ABSTRACT

**Evaluating the Clinical Care of Traumatic Brain Injury
Patients and Identifying Opportunities for Quality
Improvement in Neurosurgery at Mulago National Referral
Hospital in Kampala Uganda**

by

Benjamin Jyhhan Kuo

Duke Global Health Institute
Duke University

Date: _____

Approved:

Michael M. Haglund, Supervisor

Henry E. Rice

Joao Ricardo N. Vissoci

An abstract of a thesis submitted in partial
fulfillment of the requirements for the degree
of Master of Science in the Duke Global Health Institute
in the Graduate School of Duke University

2017

Copyright by
Benjamin Jyhhan Kuo
2017

Abstract

Background: Traumatic Brain Injury (TBI) is disproportionately concentrated in low- and middle-income countries (LMICs), with the odds of dying from TBI in Uganda more than 4 times higher than in high income countries (HICs). The objectives of this study are to describe the quality of care and determine risk factors predictive of poor outcomes for TBI patients presenting to Mulago National Referral Hospital (MNRH), Kampala Uganda.

Methods: We used a prospective neurosurgical registry based on Research Electronic Data Capture (REDCap) to systematically collect variables spanning 8 categories. Univariate and multivariate analysis were conducted to determine significant predictors of mortality.

Results: 563 TBI patients were enrolled from 1 June – 30 November 2016. 102 patients (18%) received surgery, 29 patients (5.1%) intended for surgery failed to receive it, and 251 patients (45%) received non-operative management. Overall mortality was 9.6%, which ranged from 4.7% for mild and moderate TBI to 55% for severe TBI patients with GCS 3-5. Within each TBI severity category (mild, moderate, severe GCS 6-8, severe GCS 3-5), mortality differed by management pathway. The variables predictive of mortality were: moderate to severe TBI (GCS 9-12, GCS 6-8, and GCS 3-5), more than one intracranial bleed, failure to receive surgery, high dependency unit admission, ventilator support outside of surgery, and hospital arrival delayed by more than 4 hours.

Conclusions: The overall mortality rate of 9.6% in Uganda for TBI is high, and likely underestimates the true TBI mortality. Furthermore, the wide-ranging mortality (3-82%), high

ICU fatality, and negative impact of care delays suggest shortcomings with the current triaging practices. Lack of surgical intervention when needed was highly predictive of mortality in TBI patients. Further research into the determinants of surgical interventions, quality of step-up care, and prolonged care delays are needed to better understand the complex interplay of variables that affect patient outcome. These insights guide the development of future interventions and resource allocation to improve patient outcomes.

Contents

Abstract	iv
List of Tables	viii
List of Figures	ix
Acknowledgements.....	x
1. Introduction.....	1
2. Methods.....	3
2.1 Study Design.....	3
2.2 Setting.....	4
2.3 Participants	4
2.4 Measures.....	5
2.5 Ethical Consideration	5
2.6 Analysis	6
3. Results.....	7
3.1 Outcome	7
3.2 Patient Demographics.....	8
3.3 Diagnostics	9
3.4 Clinical Assessment	10
3.5 Patient Baseline and Risk Factors	11
3.6 Care Continuum	11
3.7 Management	12

3.8 Predictors of Mortality	13
4. Discussion	16
4.1 Summary	16
4.2 Benchmarking against High Income Countries.....	16
4.3 Predictors of Mortality	18
4.4 Limitations.....	20
4.5 Future Studies.....	20
5. Conclusion	22
Appendix A – TBI Management Streams, Discharge, and Death Patterns at MNRH	23
References.....	24

List of Tables

Table 1. Variables Captured in Registry.....	5
Table 2. Demographic variable association with mortality	8
Table 3. Diagnostic variable association with mortality.....	9
Table 4. Clinical variable association with mortality.	10
Table 5. Patient baseline and risk factor variable association with mortality.....	11
Table 6. Care seeking and reaching variable association with mortality.....	11
Table 7. Management variable association with mortality	12

List of Figures

Figure 1. Mulago National Referral Hospital (MNRH) traumatic brain injury (TBI) prospective rapid electronic data capture (REDCap) registry workflow.	4
Figure 2. Mortality of traumatic brain injury (TBI) patients presenting to Mulago National Referral Hospital (MNRH) by TBI severity and management pathway	7
Figure 3. Multivariate model of predictors of traumatic brain injury (TBI) mortality by adjusted odds ratios (aOR)	14

Acknowledgements

I would first like to thank the following for their guidance and support.

My thesis advisor Professor Haglund, MD, PhD, the head of Duke Global Neurosurgery and Neuroscience (DGNN), not only inspired me to dream big, but also provided a phenomenal platform and support structure to carry out my research. He encouraged me to explore beyond my comfort zone and was there to steer me on track at crucial times.

Professor Rice, MD, a member of my thesis committee, provided his most passionate mentorship all the way in the beginning when I was still a prospective candidate at the Duke Global Health Institute (DGHI) Master's Program. His consistent input from research to classwork were the impetus for me formulating both my research project and career direction.

Joao Ricardo Vissoci, PhD, generously assisted with not only his statistical expertise, but untiringly encouraged me to keep the long-term vision in mind when I was figuring out my role in the growing DGNN group and planning my summer fieldwork.

The members of DGNN and our external collaborators who provided invaluable input throughout the various stages of this project: Silvia Vacca, BS; Emily Smith, PhD; Catherine Staton, MD; Tony Fuller, MSc; Tu Tran, MSc; Jihad Abdelgadir, MD.

Our Ugandan partners for their support in championing this project. In particular, I would like to acknowledge Dr. Michael Muhumuza, Chair of the Department of Neurosurgery at Mulago National Referral Hospital (MNRH), Dr. Lydia Nanjula, Neurosurgery Resident at MNRH, and our research assistants Josephine Nanjima, Lynet Kamusiime, and Joseph Kakaga.

Finally, my parents provided abundant love and support throughout my years of study in medical and graduate school.

Benjamin Kuo

1. Introduction

Annual global death from injury is estimated to be 5 million, of which 90% occur in low- and middle-income countries (LMICs).^{1,2} Rising injury mortality can be attributed in part to the 50% increase in road traffic injuries (RTI) over the past two decades, now accounting for a quarter of deaths from injury.¹ Furthermore, this estimation is likely to increase by the year 2030 when RTI is projected to become the fifth leading cause of death.³ Referred to as the “silent epidemic”,⁴ traumatic brain injury (TBI) is the most critical sequelae of RTIs with the greatest contribution to mortality and disability adjusted life years (DALYs).⁵

The global incidence of TBI from RTIs is disproportionately concentrated in LMICs with rates of 150-170 per 100,000 TBI in Sub Saharan Africa (SSA) compared to 106 per 100,000 globally.⁶ Furthermore, this burden is compounded by greater risk factors and limited healthcare capacity in LMICs.⁵ In a study comparing mortality from injury between Kampala Uganda and the United States, 25% of all deaths in Kampala were due to injuries compared to 7% in the United States, and the odds of dying from injury were 4.2 times higher in Uganda.⁷

The disparities in the injury incidence and outcome between LMICs and resource-rich settings have led to increased health outcomes research for TBIs and their associated risk factors in LMICs.^{8,9} While there have been increasing TBI studies in LMICs over the last decade, there is still a need for more robust prospective registries.¹⁰ In Uganda, a trauma registry implemented in 2004 at the Mulago National Referral Hospital (MNRH) showed that RTI is the major contributor (60%) of overall mortality in the casualty department.¹⁰ Within the critical care department, TBI was the most common admission diagnosis and contributed to an overall mortality rate of 44%.¹¹ In another analysis evaluating severe TBI patients, the mortality rate was 25.8%.¹² While the prior registry provides information on injury incidence and burden, it's limited in scope and doesn't

follow patients longitudinally throughout their hospital stay nor does it focus specifically on TBIs. And although these retrospective analyses are helpful for benchmarking TBI outcomes, they make it hard to identify specific quality improvement initiatives. The relationship among epidemiology, patient risk factors, clinical care, and TBI outcomes are still relatively unknown at MNRH. The objectives of this study are to describe the quality of care and determine risk factors predictive of poor outcomes for TBI patients presenting to a single tertiary hospital in Uganda.

2. Methods

2.1 Study Design

A prospective registry of TBI patients was created using Research Electronic Data Capture (REDCap). All TBI patients presenting between 1 June 2016 and 30 November 2016 to the emergency department at MNRH and referred to the neurosurgery team were included in this study. This study is a secondary data analysis of this de-identified dataset for this 6-month period. The registry data are collected over 70 hours per week, 7 days a week by two trained Ugandan research nurses. The research assistants (RAs) observed patient care throughout their hospital stay, reviewed patient charts, collected data initially on paper forms, and entered all data into the REDCap registry following patient discharge. Quality analysis occurred at two stages: 1) during data collection by the local neurosurgery resident supervising the RAs to clarify clinical variables and check for accuracy, and 2) during data analysis by the local research program manager supervising the accuracy of data entry into REDCap. Data quality was assessed among the research team every 4 weeks and discrepancies were investigated by crosschecking paper collection forms as well as the original patient charts (Figure 1).




Figure 1. Mulago National Referral Hospital (MNRH) traumatic brain injury (TBI) prospective rapid electronic data capture (REDCap) registry workflow.

2.2 Setting

The study was conducted at MNRH, which is located on Mulago Hill in the northern part of the city of Kampala. MNRH is the largest hospital in the country and serves as a teaching hospital for Makerere University College of Health Sciences.

2.3 Participants

At MNRH, TBI patients are admitted through the emergency ward. From the emergency ward, patients are either admitted to the neurosurgical ward, remain in emergency ward pending CT diagnostic, or are discharged. All TBI patients presenting to MNRH under the care of the neurosurgical team were included in the study.

2.4 Measures

A total of 60 variables were tracked in the registry and divided into 8 categories: demographics, patient baseline and risk factors of in-hospital mortality, clinical assessment, diagnostics, management, care continuum, complications, and discharge status (Table 1). Relevant risk factor variables were determined from prior studies involving the National Trauma Databank.¹³ Clinical assessment and diagnostics were recorded as described in the patient chart.

Table 1. Variables Captured in Registry.

Demographics	Age, Gender, Education, Occupation, District, Injury type, Primary care visit
Patient Baseline and Risk factors	Comorbidity, Angina, Fever, Seizure, Open wound, Transfusion, Oxygen support, Advanced airway
Clinical Assessment	Admission GCS, Change in GCS between admission and second assessment, Abnormal Pupillary Exam (Anisocoria, Light reflex, Pupil dilation), Cranial nerve deficits, Limb weakness, Polytrauma, Continence
Diagnostics	CBC, Admission vitals, Neuro exams done, CT diagnosis, Midline shift, Subjective hematoma size, Basal cistern status
Management	Management pathway, ICU require, ICU received, HDU received, BP and pulse, Ventilator support, Ventilator disruption, Physiotherapy
Care Continuum	Time from injury to seeking care, Time taken to reach hospital, Duration from arrival to neuro team review, duration from neuro review to CT results, duration from CT results to surgery, and duration from surgery to discharge
Complications	CSF leak, Oxygen requirement > 24 hour, Transfusion, Reoperation, Pneumonia, SSI, Infection, Tracheostomy, Seizure, Pressure sores, DVT, Other
Discharge Status	Death, Date of death, Discharge destination
Demographics	Age, Gender, Education, Occupation, District, Injury type, Primary care visit

GCS, Glasgow coma scale; CBC, complete blood count; CT, computed tomography; ICU, intensive care unit; HDU, high dependency unit; BP, blood pressure; CSF, cerebral spinal fluid; SSI, surgical site infection; DVT, deep vein thrombosis

2.5 Ethical Consideration

Ethical approval was provided by the Duke University and Stanford University Institutional Review Boards (IRB) (protocols 69190 and 34346, respectively) as well as the Mulago National Referral Hospital Research and Ethics Committee in Uganda (protocol 1020).

2.6 Analysis

Statistical analyses were performed using Stata version 14.0 (Stata Corp, College Station, TX). Univariate analysis of association between variables with mortality was performed using cross-tabulation, chi square, and Fisher's exact test when appropriate. Risk factors significant at the $p < 0.05$ level on univariate analysis were entered into the multivariate model for multivariate analysis. The final multivariate model was constructed through stepwise backward elimination until all the remaining variables were significant at the $p < 0.05$ level. Total cases, mortality counts, univariate p-values, unadjusted odds ratios (OR), adjusted OR, and 95% confidence intervals (CI) were reported.

3. Results

3.1 Outcome

The overall mortality rate was 9.6% (54 cases) (Figure 2). Eight died after surgery, 18 died while awaiting surgical intervention (per the patient chart), 19 died during non-operative management, and 9 died prior to CT done and admission. Mortality rate for mild TBI was 3.4% and increased to 13.8%, 15%, and 55% for moderate and severe TBI (GCS 6-8, GCS 3-5), respectively. Highest rates of death were seen in patients who failed to receive surgery and those without CT results.




Figure 2. Mortality of traumatic brain injury (TBI) patients presenting to Mulago National Referral Hospital (MNRH) by TBI severity and management pathway

3.2 Patient Demographics

During the 6-month period from 1 June 2016 to 30 November 2016, 563 patients met the inclusion criteria to be enrolled in the TBI registry at MNRH. The average age was 29 (IQR 20-36), and mortality increased by age group from 5% in the 0-14 age group to 17% in the above 45 age group (Table 2). More than 70% of patients were in the 15-44 age group and 86% were male. The most common cause of TBI was road traffic injury (62%), of which most were pedestrians and motorcyclists. Half of the patients had less than a secondary education, and 43% were referred from a primary care center.

Table 2. Demographic variable association with mortality

Demographics	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
Total patient	563 (100.0)			
Age (years)				
0-14	78 (13.8)	4 (5.1)	Ref	
15-29	243 (43.4)	18 (7.4)	0.510	
30-44	160 (28.3)	17 (10.6)	0.170	
≥ 45	81 (14.3)	14 (17.3)	0.009	3.87 (1.21-12.32)
Male gender	488 (86.4)	50 (10.3)	0.167	
Type of injury				
Assault	136 (24.4)	12 (8.8)	0.686	
Fall	64 (11.3)	12 (18.8)	0.007	2.76 (1.32-5.79)
Road traffic injury	350 (62.0)	27 (7.7)	Ref	
Motorcycle	96 (27.4)	11 (11.5)	0.090	
Car	16 (4.6)	2 (12.5)	0.296	
Pedestrian	192 (55.0)	11 (5.7)	Ref	
Education				
None	40 (20.0)	2 (5.0)	0.315	
Primary	60 (30.0)	3 (5.0)	0.622	
Secondary	88 (44.0)	3 (3.4)	Ref	
University and above	13 (6.4)	1 (7.7)	0.466	
Occupation				
Unemployed	74 (29.5)	4 (5.4)	Ref	
Self employed	180 (71.7)	12 (6.7)	0.707	
Formal employment	58 (23.1)	5 (8.6)	0.471	
Primary care referral	242 (42.8)	20 (8.3)	0.630	

3.3 Diagnostics

Documented admission vitals (HR, RR, BP, Temperature) and lab testing in the patient chart were seen in 95 out of 563 patients (17%), most of which were patients admitted for surgical intervention (Table 3). CT results were available for 440 out of 563 patients (78%), of which 9.3% were normal results. Intracranial hemorrhages diagnosed in descending order were: epidural (18%), acute subdural (15%), parenchymal (5.5%), intraventricular (1.8%), subarachnoid (3.6%). An additional 5.7% had more than one intracranial hemorrhage. Non-intracranial hemorrhage pathologies represented nearly 40% of CT findings: fracture (9.3%), edema (4.5%), contusion (12%), and more than one of the above without intracranial bleed (13%). Nearly all (98.6%) had documented admission GCS, cranial nerve and neuro examinations.

Table 3. Diagnostic variable association with mortality

Diagnostic	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
Admission vitals recorded	95 (16.8)	9 (9.5)	0.685	
Laboratory CBC done	95 (16.8)	8 (8.4)	0.449	
CT done	440 (77.9)	43 (9.8)		
Normal	41 (9.3)	1 (2.4)	Ref	
Acute subdural hematoma	66 (15.0)	10 (15.2)	0.044	8.00 (1.30-58.06)
Epidural hematoma	79 (18.0)	8 (10.1)	0.199	
Parenchymal hematoma	24 (5.5)	1 (4.2)	0.665	
Intraventricular hematoma	8 (1.8)	0 (0)	NA	
Subarachnoid hematoma	16 (3.6)	3 (18.8)	0.069	
More than one intracranial hemorrhage	25 (5.7)	7 (28.0)	0.007	18.80 (1.78-135.95)
Fracture only	41 (9.3)	0 (0)	NA	
Edema only	20 (4.5)	0 (0)	NA	
Contusion only	53 (12.0)	3 (5.8)	0.702	
More than one of the following without intracranial bleed: fracture, edema, contusion	57 (13.0)	10 (17.5)	0.033	8.51 (1.04-69.39)
Midline shift	48 (10.9)	8 (16.7)	0.158	
Basal cistern compressed or absent	44 (10.0)	5 (11.4)	0.423	
Hematoma size big or massive	28 (6.36)	6 (21.4)	0.254	

3.4 Clinical Assessment

Mild TBI (GCS 13-15) were seen in 324 patients (57.5%), moderate TBI (GCS 9-12) in 152 patients (27.0%), and severe TBI (GCS 3-8) in 87 patients (15.5%), of which 65 (11.5%) had a GCS 6-8 and 22 (4%) had a GCS 3-5 (Table 4). More than two-thirds of patients had either no documented change in GCS or an improvement of 1 or more between initial and second assessment. Abnormal pupillary findings were seen in 40 (7%).

Table 4. Clinical variable association with mortality.

Clinical Variables	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
TBI Severity				
GCS 13-15	324 (57.5)	11 (3.4)	Ref	
GCS 9-12	152 (27.0)	21 (13.8)	<0.001	4.56 (2.14-9.73)
GCS 6-8	65 (11.5)	10 (15.4)	<0.001	5.17 (2.10-12.76)
GCS 3-5	22 (3.9)	12 (54.6)	<0.001	34.15 (12.16-80.87)
GCS Change				
No change	195 (34.6)	12 (6.2)	Ref	
Improves by 1 or more	196 (34.8)	23 (11.7)	0.350	
Drops by 1	26 (4.6)	3 (11.5)	0.680	
Drops by 2 or more	40 (7.1)	10 (25.0)	0.017	4.69 (1.33-16.6)
Abnormal Pupillary Exam	40 (7.1)	12 (40.0)	<0.001	8.51 (3.66-18.07)
Cranial nerve deficit	35 (6.2)	13 (37.1)	<0.001	7.53 (3.52-16.11)
Limb weakness	61 (10.8)	18 (29.5)	<0.001	5.70 (2.97-10.93)
Polytrauma	176 (31.3)	19 (10.8)	0.438	
Incontinence	69 (12.3)	22 (31.9)	<0.001	7.38 (3.93-13.85)

TBI, traumatic brain injury; GCS, Glasgow coma scale, Abnormal pupillary exam includes presence of any of the following: anisocoria, pupil dilation, and light reflex deficits

3.5 Patient Baseline and Risk Factors

Risk factors in decreasing order of prevalence were (Table 5): comorbidity (16.5%), open wound (15%), advanced airway support (12%), seizure (11%), oxygen requirement (7%), transfusion (4%), and fever (1%).

Table 5. Patient baseline and risk factor variable association with mortality.

	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
Comorbidity	93 (16.5)	11 (11.8)	0.464	
Angina	9 (1.6)	2 (22.2)	0.208	
Oxygen support	39 (6.9)	17 (43.6)	<0.001	10.13 (4.95-20.72)
Fever	6 (1.1)	1 (16.7)	0.592	
Seizure	63 (11.2)	17 (27.0)	0.051	
Transfusion	21 (3.7)	1 (4.8)	0.453	
Open wound	85 (15.0)	10 (11.8)	0.274	
Advanced airway	66 (11.7)	20 (30.3)	<0.001	5.92 (3.15-11.11)

3.6 Care Continuum

Nearly a quarter (23%) waited a day before seeking care and 168 (30%) took more than 4 hours to reach MNRH (Table 6). Nearly half (47%) experienced an injury to arrival timeframe longer than 4 hours. The median arrival to review by the neurosurgery team was 2 hours, review to CT result was 17 hours, CT result to surgery was 71 hours, and surgery to discharge was 78 hours.

Table 6. Care seeking and reaching variable association with mortality.

	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
Seeking care: waited at least a day	128 (22.7)	18 (14.1)	0.033	1.91 (1.1-3.4)
Reaching care: > 4 hours	168 (29.8)	21 (12.5)	0.046	1.82 (1.1-3.2)
Injury to Arrival				
≤ 4 hours	298 (52.9)	24 (8.1)	Ref	
> 4 and ≤ 24 hours	128 (22.7)	15 (11.7)	0.232	
> 1 and ≤ 2 days	37 (6.6)	2 (5.4)	0.573	
> 2 days	74 (13.1)	9 (12.2)	0.269	

Arrival to Neuro review				
≤ 1 hour	239 (42.5)	22 (9.2)	Ref	
> 1 and ≤ 4 hours	132 (23.4)	18 (13.6)	0.190	
> 4 hours	189 (33.6)	13 (6.9)	0.385	
Neuro review to CT results				
≤ 4 hour	81 (26.7)	13 (16.1)	Ref	
> 4 and ≤ 24 hours	221 (72.9)	14 (6.3)	0.011	0.35 (0.16-0.79)
> 24 hours	89 (29.4)	9 (10.1)	0.253	
CT results to surgery				
≤ 4 hour	13 (12.7)	4 (30.8)	0.025	7.20 (1.27-40.68)
> 4 and ≤ 24 hours	15 (14.7)	0	NA	
> 24 hours	60 (58.8)	4 (6.7)	Ref	

3.7 Management

Surgery was performed for 102 patients (18%) presenting to MNRH with TBI (Table 7) and intended but not provided for 29 patients (5.1%) due to limited operating theatre capacity and delays along the neurosurgical care continuum. Non-operative management was provided to 251 patients (45%). Nearly a third (181 patients, 32.1%) were not admitted from the emergency department, of which 111 (19.7%) did not receive CT testing.

Of all 563 TBI patients, 74 patients (13.1%) required intensive care unit (ICU) support, but only 34 patients (46%) received it. ICU availability was predominantly offered to severe TBI patients (57.1%) compared to moderate TBI (32%) and mild TBI (28.6%) cases. Nearly all patients entering ICUs (94.1%) received continuous BP and pulse monitoring; 79.4% received mechanical ventilation, of which nearly a third (29.6%) had documented ventilator disruption. 35 patients (6.2%) entered the HDU and only 21 of 59 patients received adequate physiotherapy – completed as requested by the neurosurgical team in the patient chart.

Table 7. Management variable association with mortality

	Total Cases (%)	Died (%)	Univariate p-Value	Unadjusted OR (95% CI)
Management Pathway				
Surgery received	102 (18.1)	8 (7.8)	Ref	
Non-operative	251 (44.6)	19 (7.6)	0.887	

Not admitted CT done	70 (12.4)	0	NA	
Not admitted CT not done	111 (19.7)	9 (8.1)	0.943	
Surgery not received	29 (5.1)	18 (62.1)	< 0.001	19.23 (6.79-54.45)
HDU received	35 (6.2)	10 (28.6)	< 0.001	6.66 (2.86-15.51)
ICU needed	74 (13.1)	35 (47.3)	< 0.001	24.60 (12.65-47.84)
ICU received	34 (6.0)	18 (52.9)	< 0.001	15.41 (7.25-33.73)
BP and pulse monitoring in ICU	32 (94.1)	16 (50.0)	< 0.001	12.97 (6.02-27.95)
Ventilator support				
Not needed	489	19 (3.9)	Ref	
Received	27 (36.5)	16 (59.3)	< 0.001	36.06 (14.75-88.18)
Needed but not received	47 (63.5)	19 (41.3)	< 0.001	17.52 (8.28-36.74)
Ventilator disruption	8 (29.6)	3 (37.5)	0.015	6.12 (1.42-26.39)
Physiotherapy				
Not needed	504 (89.5)	24 (5.5)	Ref	
Received	21 (3.7)	4 (19.1)	0.016	4.21 (1.25-12.88)
Not adequate or none	38 (6.7)	4 (10.5)	0.191	
Any morbidity	98 (17.4)	15 (26.3)	< 0.001	4.27 (2.17-8.37)

CT, computed tomography; HDU, high dependency unit; ICU, intensive care unit; BP, blood pressure

Percentages by columns

ICU received, Ventilator support percentages calculated using ICU needed as denominator.

BP and Pulse monitoring percentage calculated using ICU received as denominator

Ventilator disruption percentage calculated using Ventilator received as denominator

3.8 Predictors of Mortality

The two demographic variables associated with mortality on univariate analysis, age ≥ 45 and TBI due to fall, were not retained in the multivariate model (Table 2).

While acute subdural hematoma, more than one intracranial bleed, and more than one minor pathology without intracranial bleed were associated with mortality on univariate analysis, only more than one intracranial bleed was retained in the multivariate model with adjusted OR of 7.0 (2.1-23.5) (Table 3 and Figure 3).




Figure 3. Multivariate model of predictors of traumatic brain injury (TBI) mortality by adjusted odds ratios (aOR)

Eight of the thirteen clinical assessment variables were associated with mortality on univariate analysis. In decreasing order of unadjusted OR, they are: GCS 3-5, abnormal pupillary exam, cranial nerve deficit, incontinence, limb weakness, GCS 6-8, drops in GCS of 2 or more, and GCS 9-12. Following multivariate analysis, only TBI severity (GCS 3-5 aOR 11.6, 3.0-44.3, GCS 6-8 aOR 5.6, 2.0-15.8, GCS 9-12 aOR 3.3, 1.3-8.0) was retained (Table 4 and Figure 3).

Although two patient risk factors were associated with mortality on univariate analysis, oxygen support and advanced airway on initial assessment, they were not significant in the multivariate analysis (Table 5). Among the care continuum variables, only reaching care > 4 hours was retained after multivariate analysis (aOR 3.6, 1.3-8.8) (Table 6 and Figure 3).

All nine of the management variables were associated with mortality on univariate analysis. In decreasing order of unadjusted OR, they are: ventilator support, ICU need, failure to receive surgery, ICU received, continuous BP and pulse monitoring, HDU care, ventilator disruption, any morbidity, and physiotherapy received. In multivariate modelling, three management variables remained significant in predicting mortality: surgery intended but not received (aOR 21.3, 4.5-78.5), ventilator support needed outside of surgery (received aOR 4.6, 1.1-21.9 and not received aOR 8.7, 2.0-38.8), and HDU care (aOR 3.5, 1.8-8.7) (Table 7 and Figure 3).

4. Discussion

4.1 Summary

To our knowledge, this is the first prospective TBI registry established in Uganda spanning eight variable categories to understand the biggest drivers for poor outcomes in TBI patients and to highlight areas for quality improvement. Within the 6-month study period, 563 patients presented to MNRH with acute TBI predominantly from RTC (62%). 102 patients (18%) received surgery, and 8 died postoperatively (8%). 29 (5%) patients failed to receive surgery due to infrastructural limitations, of which 18 died (62%). 251 patients (45%) received non-operative management, with a mortality rate of 7.6%. 28 variables were significantly associated with mortality on univariate analysis, and 9 variables remained significant on multivariate analysis: moderate to severe TBI (GCS 3-5, 6-8, and 9-12), more than one intracranial bleed, surgery intended but not received, HDU care, ventilator support needed and received or not received, and reaching care in greater than 4 hours.

4.2 Benchmarking against High Income Countries

In-hospital mortality for TBI patients was 9.6%, which is comparable to most contemporary studies in high income countries (HIC).^{14,15} ICU and ventilator status were associated with dramatically increased mortality: 60% and 67%, respectively. Moreover, mortality was 40% for patients in need of ventilator support outside of surgery who did not receive it. In comparison, studies in the United States and Italy have reported 26.5% and 17.2% mortality for TBI patients admitted to the ICU, respectively.^{16,17} A separate United States study in patients with severe isolated TBI identified mechanical ventilation as a modifiable risk factor for in-hospital mortality.¹³ Comparable Kenyan, Nigerian, and Tanzanian studies have found 54%, 52.2%, and 47% mortality, respectively¹⁸⁻²⁰, with the Nigerian study additionally reporting poorer

outcomes for patients receiving mechanical ventilation. When this study was conducted, MNRH had three to four functional ventilators, all located in the six-bed ICU, thereby providing an opportunity for targeted quality initiatives. Additional research is needed to understand the baseline factors of patients receiving such services and the most effective utilization to improve patient outcomes.

We found that lack of access to timely surgical intervention was also associated with increased mortality. Surgical patients had the lowest mortality of all management pathways. However, when significant provider delays and infrastructural bottlenecks prevent patients from receiving timely surgery, mortality soars from 7.8% to 62%. Based on clinical assessment and diagnostics, these patients were admitted to the neurosurgical ward and planned for surgery. However, due to diagnostic delays, cost, limited theatre space, or other factors, they died preoperatively.

The impact of timely surgery is especially apparent in mild TBI patients whose mortality rises from 3.1% when operated to 37.5% when surgery was intended but not delivered. Of the 11 mild TBI patient deaths, four had skull fractures, three had epidural hematomas, one had a subdural hematoma, one had a normal CT, and two awaited CT results. Three experienced a drop in GCS of two or more from admission to next recording and one only had an admission GCS recorded. Half were managed non-operatively, two were postoperative, and three awaited surgery. The causes of death were sepsis (n=2), meningitis (n=1), aspiration (n=1), and unknown (n=7). Given the limited clinical documentation for this group, poor monitoring and management of neurological status deterioration may contribute to mortality. Standardized clinical documentation coupled with further research is necessary to identify gaps in the management of mild TBI patients and the role of timely surgery.

Prior HIC studies have shown a drop in mortality by 50% in patients who underwent craniotomy or hematoma drainage within four hours of arrival to the emergency department²¹, highlighting the importance of timely clinical assessment and neurosurgical intervention in TBI patients. At MNRH, the importance of timely CT imaging is further emphasized in severe TBI patients of which nearly one in three are not admitted to neurosurgery due to the lack of CT scan. Of those not admitted to neurosurgery, nearly half died in-hospital, suggesting an even higher actual mortality rate, as post-hospital outcomes are not recorded. Interestingly, obtaining CT scan within 4 hours of arrival had a 7.2 OR of death compared to delayed CT findings, which suggests that sicker patients may be triaged first, but without improved outcomes. Therefore, further research is needed to understand the magnitude, causes, and impact of delays in MNRH.

Similarly, pre-hospital delays can negatively impact patient outcomes. Delays in seeking care longer than 24 hours and in reaching care longer than four hours were strongly associated with high mortality. To address delays in seeking care, future interventions may target patient education with respect to the time sensitive nature of TBI. Delays in reaching care point to infrastructural factors such as limited access to emergency vehicles, navigating unpaved roads and public transportation, and highly congested traffic. The strong association between transport delays and poor outcomes emphasizes the need for future research and interventions aimed at identifying and improving infrastructural bottlenecks.

4.3 Predictors of Mortality

Factors predictive of TBI mortality in Uganda can be divided into delays in reaching care, injury severity (GCS<13, more than one intracranial bleed on CT), and hospital interventions (surgery and step-up care). As previously discussed, timely access to neurosurgical

intervention is essential for TBI patients, therefore delays in reaching care negatively impact patient outcomes. Patient baseline and injury risk factors are consistent with previous studies which report worse outcomes for patients with more severe TBI.^{14,22} Similarly, over half of patients with more than one intracranial bleed had moderate to severe TBI.

With respect to hospital interventions, the classification of a patient for whom surgery was intended but not received proved to be highly predictive of mortality. These patients experienced a *more than tenfold mortality than all other patients*. One reason they were not operated is their choice to self-discharge against medical advice. Therefore, the high mortality rate reported may be an underestimation of the true mortality rate, since patient outcomes post-discharge are unknown. For patients dying in the hospital prior to their surgery, reasons for surgical delays are multifaceted, including limitations in theatre space, delayed procurement of supplies, and diagnostic logistics. The dramatic increase in mortality for patients not receiving necessary surgical care stresses the importance of increasing access to neurosurgery in future interventions to address morbidity and mortality.

With respect to diagnostic capacity, patients at MNRH are faced with challenges in access and cost similar to patients in other LMIC settings.²³ To obtain a head CT scan - necessary for surgical decision making - caregivers typically arrange their own transportation to nearby private Nakasero Hospital, the only facility within close proximity to MNRH with a functional CT scanner, and pay for the scan and films out of pocket. The financial burden of a CT scan, which can range from \$70 to \$132²⁴, can be almost equal to the average monthly income in Uganda.²⁵ Further investigation into the causes of surgical delays is necessary to guide future interventions.

Similarly, receiving step-up care through HDU admission and mechanical ventilation was predictive of mortality. Although ICU admission was not retained as a predictive factor in the

multivariate analysis, the co-linearity between ICU care and mechanical ventilation, coupled with the strong association between ICU admission and high mortality, emphasize the opportunity for targeted interventions in step-up care to impact patient outcomes.

4.4 Limitations

In an effort to provide a comprehensive perspective on TBI patient management at MNRH, 60 variables throughout the care continuum are tracked per patient. As a result, this study was limited by the ability to collect complete data for all patients throughout their hospital stay. Moreover, the quality of data collection was limited by availability of clinical documentation. For example, only 17% of patients had admission vitals recorded in the patient charts, with the proportion of the other 83% that were not assessed or assessed but not recorded unknown. Because the RAs are not clinically trained, they are unable to collect missing data points such as admission vitals not documented in the charts. To mitigate the burden on clinical staff, data collection forms were printed and placed in the patient charts. In order to improve sustainability and reduce redundancy, future data collection efforts should aim to improve form completion by integrating prospective data collection into the local workflow.

4.5 Future Studies


This study provides a framework for future TBI studies and registry development for other patient populations in LMIC. Review of our prospective neurosurgical registry estimates the burden of TBI to account for 80% of patients managed by the neurosurgery department at MNRH. As such, future studies may expand to include all neurosurgical patients at MNRH. The patient outcomes presented in this study also call for further research into the underlying infrastructural and systems-based factors affecting mortality. The drastically high mortality in patients for whom surgery was intended but not received warrants further exploration into the

causes of surgical delays. By identifying bottlenecks in the care continuum, future interventions may address infrastructural or provider delays affecting patient outcomes. Moreover, the prospective nature of the registry allows for up-to-date statistics on the management of TBI patients at MNRH to the clinical team. In a setting where patient records are in paper format, regular and timely reports based on an electronic registry may provide high-level insights to the clinical team that would otherwise be unavailable.

5. Conclusion

While there is increasing literature describing the disproportional burden of TBI in LMICs, its epidemiological pattern, and increased surgical capacity programs, there remains a need to systematically examine how demographics, diagnostics, patient risk profiles, clinical data, care continuum delays, and management pathways interrelate and affect patient outcomes. Our prospective study examined 563 TBI patients across 8 variable categories presenting to MNRH over a 6-month period to determine significant predictors of poor outcomes and highlight areas for improvement and further research. Although the mortality rate is comparable to other LMICs, it ranged from 3-82% depending on the intervention type, its availability, and disease severity, with tenfold increase in mortality for patients intended for surgery who failed to receive it. As TBI burden remains consistently high with a complex interplay of factors contributing to the heterogeneous mortality rate, further research and collaborative efforts in complementing better resource allocation practices to existing neurosurgical residency education are imperative.

Appendix A – TBI Management Streams, Discharge, and Death Patterns at MNRH



References

1. Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet*. 2012;380(9859):2095-2128. doi:10.1016/S0140-6736(12)61728-0.
2. Hofman K, Primack A, Keusch G, Hrynkow S. Addressing the growing burden of trauma and injury in low- and middle-income countries. *Am J Public Health*. 2005;95(1):13-17. doi:10.2105/AJPH.2004.039354.
3. World Health Organization. Global status report on road safety. *Inj Prev*. 2015:318. doi:http://www.who.int/violence_injury_prevention/road_safety_status/2015/en/.
4. Langlois JA, Rutland-Brown W, Wald MM. The Epidemiology and Impact of Traumatic Brain Injury A Brief Overview. *J Head Trauma Rehabil*. 2006;21(5):375-378. doi:00001199-200609000-00001 [pii].
5. Hyder, A.A.; Colleen, A.w.; Puvanachandra, P., Gururaj, G.; Kobusingye OC. The impact of traumatic brain injuries : A global perspective. *NeuroRehabilitation*. 2007;22:341-353.
6. Chandran A, Hyder AA, Peek-Asa C. The global burden of unintentional injuries and an agenda for progress. *Epidemiol Rev*. 2010;32(1):110-120. doi:10.1093/epirev/mxq009.
7. Jayaraman S, Ozgediz D, Miyamoto J, et al. Disparities in injury mortality between uganda and the United States: Comparative analysis of a neglected disease. *World J Surg*. 2011;35(3):505-511. doi:10.1007/s00268-010-0871-z.
8. Bhalla K, Naghavi M, Shahraz S, Bartels D, Murray CJL. Building national estimates of the burden of road traffic injuries in developing countries from all available data sources: Iran. *Inj Prev*. 2009;15(3):150-156. doi:10.1136/ip.2008.020826.
9. Bhalla K, Harrison J, Abraham J, et al. Data sources for improving estimates of the global burden of injuries: Call for contributors. *PLoS Med*. 2009;6(1):0022-0024. doi:10.1371/journal.pmed.1000001.
10. Demyttenaere S V, Nansamba C, Nganwa A, Mutto M, Lett R, Razek T. Injury in Kampala, Uganda: 6 years later. *Can J Surg*. 2009;52:E146-50. http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2769114/pdf/052e146.pdf.
11. Kwizera A, Dünser M, Nakibuuka J. National intensive care unit bed capacity and ICU patient characteristics in a low income country. *BMC Res Notes*. 2012;5:475. doi:10.1186/1756-0500-5-475.
12. Tran TM, Fuller AT, Kiryabwire J, et al. Distribution and characteristics of severe traumatic brain injury at Mulago National Referral Hospital in Uganda. *World Neurosurg*. 2015;83(3):269-277. doi:10.1016/j.wneu.2014.12.028.
13. Krishnamoorthy V, Vavilala MS, Mills B, Rowhani-rahbar A. Demographic and clinical risk factors associated with hospital mortality after isolated severe traumatic brain injury : a cohort study. *J Intensive Care*. 2015:1-8. doi:10.1186/s40560-015-0113-4.
14. Gagné M, Moore L, Sirois M-J, Simard M, Beaudoin C, Kuimi BLB. Performance of International Classification of Diseases–based injury severity measures used to predict in-hospital mortality and intensive care admission among traumatic brain-injured patients. *J Trauma Acute Care Surg*. 2017;82(2):374-382. doi:10.1097/TA.0000000000001319.
15. Selassie AW, Fakhry SM, Ford DW. Population-Based Study of the Risk of In-Hospital Death After Traumatic Brain Injury: The Role of Sepsis. *J Trauma Inj Infect Crit Care*. 2011;71(5):1226-1234. doi:10.1097/TA.0b013e318226ecfc.
16. Gupta D, Sharma D, Kannan N, et al. Guideline Adherence and Outcomes in Severe Adult Traumatic Brain Injury for the CHIRAG (Collaborative Head Injury and Guidelines) Study. *World Neurosurg*. 2016;89:169-179. doi:10.1016/j.wneu.2015.12.097.
17. Di Saverio S, Gambale G, Coccolini F, et al. Changes in the outcomes of severe trauma patients from 15-year experience in a Western European trauma ICU of Emilia Romagna region (1996–2010). A population cross-sectional survey study. *Langenbeck's Arch Surg*. 2014;399(1):109-126.

- doi:10.1007/s00423-013-1143-9.
18. Opondo EA, Mwangombe NJM. Outcome of severe traumatic brain injury at a critical care unit: a review of 87 patients. *Ann African Surg.* 2009;1(1). doi:10.4314/aas.v1i1.45788.
 19. Tobi KU, Azeez AL, Agbedia SO. Outcome of traumatic brain injury in the intensive care unit: a five-year review. *South African J Anaesth Analg.* 2016;22(5):135-139. doi:10.1080/22201181.2016.1206293.
 20. Staton CA, Msilanga D, Kiwango G, et al. A prospective registry evaluating the epidemiology and clinical care of traumatic brain injury patients presenting to a regional referral hospital in Moshi, Tanzania: challenges and the way forward. *Int J Inj Contr Saf Promot.* 2015;(August):1-9. doi:10.1080/17457300.2015.1061562.
 21. Kim YJ. The Impact of Time from ED Arrival to Surgery on Mortality and Hospital Length of Stay in Patients With Traumatic Brain Injury. *J Emerg Nurs.* 2011;37(4):328-333. doi:10.1016/j.jen.2010.04.017.
 22. Rates of TBI-related Deaths by Age Group — United States, 2001–2010 | Concussion | Traumatic Brain Injury | CDC Injury Center. *Centers Dis Control Prev.* 2016. https://www.cdc.gov/traumaticbraininjury/data/rates_deaths_byage.html.
 23. Mwita CC, Muthoka J, Maina S, Mulingwa P, Gwer S. Early management of traumatic brain injury in a Tertiary hospital in Central Kenya: A clinical audit. *J Neurosci Rural Pract.* 2016;7(1):97-101. doi:10.4103/0976-3147.165390.
 24. Abdelgadir J, Tran T, Muhindo A, et al. Estimating the Cost of Neurosurgical Procedures in a Low-income Setting: An Observational Economic Analysis. *World Neurosurg.* 2017. doi:10.1016/j.wneu.2017.02.048.
 25. Ubos. Uganda National Household Survey 2009/2010. *Uganda Bur Stat.* 2010:222. doi:UGA-UBOS-UBOS-2012-v01.