

AIDS-Associated *Penicillium marneffe* Infection of the Central Nervous System

Thuy Le,^{1,3} Nguyen Huu Chi,² Ngo T. Kim Cuc,² Tran Phu Manh Sieu,² Cecilia M. Shikuma,³ Jeremy Farrar,¹ and Jeremy N. Day¹

¹Wellcome Trust Major Overseas Programmes, Oxford University Clinical Research Unit, ²Hospital for Tropical Diseases, Ho Chi Minh City, Vietnam; and ³Hawaii Center for AIDS, University of Hawaii at Manoa, Honolulu, Hawaii

***Penicillium marneffe* is an important human immunodeficiency virus–associated opportunistic infection endemic in Southeast Asia. Central nervous system infection has not been described. We report the first case series of 21 human immunodeficiency virus–infected patients who presented with a syndrome consistent with acute central nervous system infection and who had *Penicillium marneffe* isolated from cerebrospinal fluid.**

Penicillium marneffe is emerging as an important opportunistic pathogen among human immunodeficiency virus (HIV)–infected and immunocompromised residents of (and travelers to) Southeast Asia, Northeastern India, and Southern China [1]. No definitive mode of acquisition has been found, but inhalation has been implicated [2, 3]. The infection has been described in both immunocompromised (>80%) and immunocompetent individuals [4]. Immunocompromised individuals develop disseminated disease involving the reticuloendothelial, skin, respiratory, and gastrointestinal systems. *P. marneffe* is commonly isolated from skin lesions (90%), blood (76%), bone marrow (100%), and lymph nodes (34%) [5]. There are 2 reports of isolation of *P. marneffe* from cerebrospinal fluid (CSF) of patients with penicilliosis [5, 6]. However, clinical details were lacking in these cases, and despite the fact that other members of the *Penicillium* genus are known to cause central nervous system (CNS) disease, penicilliosis presenting

as a predominantly CNS infection has not been described [6]. We report the clinical characteristics and outcomes of 21 HIV-infected adult patients who presented with symptoms consistent with a CNS infection and who had *P. marneffe* isolated from CSF.

Patients. The Hospital for Tropical Diseases in Ho Chi Minh City is the largest referral center for infectious diseases in Vietnam, with >5000 HIV-infected patients seen yearly. From January 2004 through December 2009, 677 incident cases of penicilliosis were retrospectively identified from our hospital records. Diagnosis was confirmed by culture of *P. marneffe* from skin scrapings, blood, lymph nodes, bone marrow, or other body tissue. Fifty-eight (8.6%) of these patients had lumbar puncture performed for suspected CNS infection. A definitive diagnosis of CNS infection was established by CSF culture and/or microscopy (Gram, India, and Zielh-Neelsen stains) for 29 patients (50%); 20 were due to *P. marneffe*, 7 to *Cryptococcus neoformans*, 1 to *Mycobacterium tuberculosis*, and 1 to both *P. marneffe* and *C. neoformans*. The clinical features and outcomes of the 21 patients with *P. marneffe* cultured from CSF samples were reviewed and are reported in Table 1. The study was approved by the Ethical Committee of the Hospital for Tropical Diseases.

The 21 patients were admitted to Hospital for Tropical Diseases from April 2004 through July 2009. All were HIV infected; the median CD4 count was 11 cells/ μ L (interquartile range [IQR], 10–12 cells/ μ L). The median age was 28 years (IQR, 25–31 years); 76% were men. The median duration of illness was 9 days (IQR, 3–30 days). Patients experienced fever (90%), anorexia (57%), fatigue (52%), cough (33%), and diarrhea (29%) prior to the onset of altered mentation that prompted the hospital visits (48%). The median duration of altered mentation was 2 days (IQR, 1.25–2 days). The median temperature was 38.3°C (IQR, 37°C–39°C). On examination, characteristic umbilicated skin lesions were present in 10 patients (photograph of skin lesions in Figure 1), oral thrush in 9, hepatosplenomegaly in 12, and lymphadenopathy in 3 patients. Blood tests showed anemia in 20 patients (median hemoglobin level, 7.2 g/dL; IQR, 5.6–9 g/dL), thrombocytopenia in all patients (median platelet count, 85,000 cells/ μ L; IQR, 31,000–125,500 cells/ μ L), and elevated transaminase levels in 18 patients (median alanine transaminase level, 220 units/L; IQR, 109–306 units/L; median aspartate transaminase level, 130 units/L; IQR, 86–181 units/L). Symptoms of altered mentation, including confusion,

Received 25 June 2010; accepted 20 August 2010; electronically published 5 November 2010.

Reprints or correspondence: Dr Thuy Le, Wellcome Trust Major Overseas Programmes, Oxford University Clinical Research Unit, Vietnam. 190 Ben Ham Tu, District 5, Ho Chi Minh City, Vietnam (thuy@oucru.org).

Clinical Infectious Diseases 2010;51(12):1458–1462

© 2010 by the Infectious Diseases Society of America. All rights reserved.
1058-4838/2010/5112-0017\$15.00

DOI: 10.1086/657400

agitation, or drowsiness, were the presenting features in 10 patients and developed in all remaining patients during hospitalization. Headache was the presenting symptom in 2 patients. Signs of meningeal irritation were absent. Generalized convulsions and facial nerve palsy were observed in 2 and 1 patients, respectively. The CSF analysis is shown in Table 1. The CSF opening pressure was not routinely measured; in 1 patient it measured 180 mm CSF. CSF was clear in all patients and acellular in 14 patients. Among the 7 patients with CSF pleocytosis, the median nucleated cell count was 80 cells/mL (IQR, 19–170 cells/mL), with neutrophil predominance in 3 patients. CSF protein level was elevated (>0.45 g/dL) in 71% of patients, with a median protein level of 0.7 g/dL (IQR, 0.5–0.88 g/dL). The median CSF/serum glucose ratio was 0.65 (IQR, 0.5–0.78); 5 patients had CSF/serum glucose ratios <0.5 . Gram stain of CSF was negative in all patients. India ink stain of CSF was positive in 1 of 19 patients tested, and *C. neoformans* was cultured from blood and CSF samples of this patient (patient 11). Ziehl-Neelsen stain of CSF was negative in all 17 patients examined. A diagnosis of tuberculosis was made in 2 patients by positive Ziehl-Neelsen stain of sputum and lymph node biopsy. The mean time to identify *P. marneffei* from CSF culture was 4.4 days (range, 2–8 days) and from blood culture was 4.3 days (range, 3–9 days). Blood culture was performed in 18 patients; 13 grew *P. marneffei*, 1 grew *Escherichia coli*, 1 grew *Enterococcus* species, 1 grew unidentified gram-negative rods, and 1 grew both *P. marneffei* and *C. neoformans*.

Outcomes. Three patients survived and had experienced improvement of symptoms at hospital discharge, 1 was transferred to another hospital for tuberculosis treatment after his condition deteriorated, 12 died within 24–72 h after hospital admission, and 5 were taken from the hospital to die at home, which is common practice in Vietnam. Patients taken home to die were moribund and received no further effective medical care, and all were expected to have died, giving an overall mortality of 81%. Among the 18 patients with a poor outcome, the diagnosis was not established prior to death, and no antifungal drugs were given in 12 patients. Two patients received 2 doses of itraconazole (400 mg/day) on the basis of characteristic skin lesions, 1 received 1 dose of amphotericin B on the basis of preliminary blood culture results indicating growth of yeasts, and the remaining 3 patients received 7–17 days of itraconazole or amphotericin B treatment but had concurrent comorbid conditions. The 3 patients who survived and had improvement of symptoms at hospital discharge started receiving amphotericin B within 24 h after developing CNS symptoms, on the basis of detection of unidentified yeasts from blood culture (patients 10 and 14) or positive India ink stain of CSF (patient 11), and all received a total of 14 days of amphotericin B therapy before switching to itraconazole or fluconazole.

Discussion. This report describes a new clinical syndrome associated with *P. marneffei* infection in HIV-infected patients. The syndrome is characterized by an acute onset of altered mental status with confusion, agitation, or depressed consciousness in the setting of a subacute febrile illness with non-specific constitutional symptoms. Symptoms of increased cranial pressure and signs of meningeal inflammation were notably uncommon or absent. Characteristic umbilicated skin lesions were present in only one-half of the patients. CSF analysis varied from acellular to mild pleocytosis, with normal to mildly elevated protein levels and normal to mildly low glucose levels. CSF microscopy for *P. marneffei* was negative. CSF culture for *P. marneffei* took a mean of 4 days for identification. *P. marneffei* was not always isolated from blood cultures of these patients. The disease course was rapidly progressive, and inpatient mortality was very high. Early initiation of amphotericin B was administered in 5 patients, 3 of whom survived, whereas all 15 patients who did not receive amphotericin B or itraconazole died.

To our knowledge, *P. marneffei* has never been described as a CNS pathogen. *P. marneffei* was isolated from the meninges of 1 patient in a review of 155 published penicilliosis cases and from 3 of 20 CSF specimens in a case series of 80 patients with penicilliosis from Thailand [4, 5]. However, the clinical features of those patients were not provided. The patients in this case series had CNS symptoms consistent with a CNS infection, and *P. marneffei* was the single pathogen isolated from CSF samples from 20 of 21 patients. It is not possible to exclude the involvement of *M. tuberculosis* by CSF and sputum microscopy and of other opportunistic viral pathogens not tested in this case series; therefore, it remains uncertain whether the CNS syndrome in these patients can be wholly attributed to *P. marneffei*. Concurrent growth of *C. neoformans* and *P. marneffei* has been observed from blood and CSF samples of HIV-infected patients at our hospital [7], as occurred for patient 11 in this case series. *P. marneffei* grows much slower (at least 24–48 h later) than *C. neoformans* from clinical specimens, and neither has been observed to have competitive cultural advantages over the other. Unlike *M. tuberculosis*, cryptococcus is unlikely to have been missed, although cryptococcal antigen tests, if they were clinically available, would have been more informative. Selected members of the other 225 *Penicillium* species are known to cause CNS disease. *Penicillium commune* was isolated from multiple brain and lung autopsy specimens from a patient with acute leukemia who was receiving antibiotics and steroids [8]. *Penicillium chrysogenum* was isolated from CSF and brain biopsy samples of 2 nonimmunocompromised individuals with CNS symptoms [9, 10]. An unidentified *Penicillium* species was isolated from multiple brain lesions of a patient with chronic liver disease at autopsy [6]. These reports demonstrate the neu-

Table 1. Clinical Features of 21 HIV-Infected Patients with *Penicillium marneffei* Isolated from Cerebrospinal Fluid (CSF)

Patient	Age, years	Sex	Duration of illness, days	Clinical characteristic			CSF analysis			Treatment and/or hospital course	Outcome
				Presenting symptoms	Presenting sign(s)	Cell counts, cells/mL	Protein level, ^a g/dL	CSF/serum glucose ratio			
1	25	F	3	Fever, fatigue, anorexia, diarrhea	Alert, wasting, thrush, skin lesions	NC, 1; RBC, 1	0.9	3/5	Itra 400 mg/day for 17 days; MS change	Died 96 h after MS change; concurrent GNR, sepsis	
2	23	M	30	Fever, cough, 24 h of MS change	GCS 8; thrush, tachypnea, hepatosplenomegaly	NC, 8; RBC, 1	0.5	1.2/3.1	No prescription	Died 24 h after admission	
3	29	F	30	Fever, cough, anorexia, 24 h of MS change	Decreasing GCS, thrush, skin lesions, hepatosplenomegaly	NC, 30 (58% N, 42% L); RBC, 2800	0.68	1.6/3.2	No prescription	Died 72 h after admission	
4	28	M	7	Fever, cough, fatigue, diarrhea	Thrush	NC, 1; RBC, 1	0.7	3/3.9	No prescription; MS change and convulsion	Died 24 h after MS change	
5	28	M	14	Fever, cough, anorexia, diarrhea	Alert, wasting, thrush, skin lesions, hepatomegaly	NC, 1; RBC, 4	0.5	3.5/4.2	No prescription; MS change	Died 72 h after MS change	
6	28	M	2	Fever, MS change	Decreasing GCS, jaundice, hepatosplenomegaly	NC, 6; RBC, 1	0.6	2.8/4.3	No prescription	Died 24 h after admission	
7	37	M	2	Fever, MS change	Decreasing GCS, skin lesions, splenomegaly	NC, 1; RBC, 1	0.7	3.7/7	No prescription	Died 48 h after admission	
8	31	M	14	Fever, nausea, fatigue, abdominal pain, anorexia	Alert, thrush, skin lesions, diffuse abdominal tenderness	NC, 80 (20% N, 80% L); RBC, 1	1.2	1.5/3	No prescription; MS change and convulsion	Died 48 h after MS change	
9	25	M	30	Headache, pus in both ears, hearing loss	Alert, cranial nerve 7 palsy, bilateral deafness, pus in both ear canals	NC, 1; RBC, 1	0.8	3.2/3.8	AmB for 9 days; MS change	Died after 9 days of AmB; suspected brain abscess	
10	25	F	10	Fever, fatigue, skin lesions	Alert, wasting, hepatosplenomegaly, skin lesions	NC, 260 (82% N, 18% L); RBC, 580	1.5	2.8/4	AmB for 14 days; MS change	Improved, discharged home with Itra 400 mg/day after 17 days of hospitalization	

11	22	M	21	Fever, headache, diarrhea	Alert, hepatosplenomegaly	NC, 1640 (94% N, 6% L); RBC, 12	0.4	2/3.2	AmB for 14 days, followed by fluconazole 450 mg/day	Improved; discharged home after 21 days of hospitalization; concurrent cryptococcal meningitis
12	24	M	7	Fatigue, anorexia, dysphagia	Alert, wasting, skin lesions	NC, 80 (4% N, 96% L); RBC, 5100	1.78	3.6/5.4	Itra 400 mg/day for 1 day; MS change	Worsening MS at discharge 72 h after admission
13 ^b	23	M	60	Fever, fatigue, weight loss, diarrhea	Alert, wasting, skin lesions, hepatosplenomegaly	NC, 1; RBC, 0	0.2	2.7/7.7	Itra 400 mg/day for 7 days; MS change	Worsening MS at time of transfer to TB hospital for concurrent pulmonary TB
14	33	M	120	Fever, cough, vomiting, weight loss	Alert, thrush, hepatosplenomegaly	NC, 2; RBC, 1	0.6	3/4	AmB for 14 days, then Itra 400 mg/day for 57 days; MS change	Improved; discharge after 80 days of hospitalization
15	41	M	1	Fever, MS change	GCS 8, cervical lymphadenopathy, hepatosplenomegaly	NC, 2; RBC, 1	0.7	3.8/5.5	No prescription	Worsening MS at discharge 48 h after admission
16	24	F	2	Fever, MS change, cough	Decreasing GCS, wasting, hepatomegaly	NC, 1; RBC, 1	0.3	3.6/4.2	AmB for 24 h	Worsening MS at discharge 72 h after admission
17	31	M	30	Fever, fatigue, cough, anorexia	Alert, wasting, hepatomegaly	NC, 4; RBC, 0	0.3	1.8/2	No prescription; MS change	Died 24 h after MS change
18	31	M	6	Fever, 72 h of MS change	Decreasing GCS, skin lesions	NC, 3; RBC, 1	0.4	2.4/2.6	Itra for 24 h	Died 48 h after admission
19	29	M	4	Fever, MS change, abdominal pain	Decreasing GCS, thrush, cervical lymphadenopathy, wasting	NC, 1; RBC, 1	0.7	2.6/3.2	No prescription	Family requested discharge home 72 h after admission
20	30	M	Unknown	Fever, MS change	GCS 7, thrush, hepatosplenomegaly, skin lesions	NC, 8; RBC, 16,000	0.9	1.6/3.9	No prescription	Died 24 h after admission
21 ^b	60	F	2	Fever, MS change	GCS 10, wasting	...	0.3	1.9/3.6	No prescription	Family requested discharge home 24 h after admission

NOTE. AmB, amphotericin B; Itra, itraconazole; GCS, Glasgow Coma Scale; GNR, gram-negative rod; L, leukocyte; MS, mental status; N, neutrophil; NC, nucleated cell; RBC, red blood cell; TB, tuberculosis.

^a Normal protein level, <0.45 g/dL.

^b Patient 13 CSF lactate level, 6.1 mmol/L. Patient 21 CSF lactate level, 6.3 mmol/L.



Figure 1. Umbilicated skin lesions characteristic of penicilliosis.

rotropic potential of invasive *Penicillium* species, and *P. marneffeii*, the most invasive of *Penicillium* species, is likely not an exception.

CNS infections are medical emergencies, and early empirical therapy can prevent irreversible neuronal damage and save lives. In HIV-infected patients with CD4 count <100 cells/ μ L who present with a subacute febrile syndrome, including changes in mental status, and who live or have traveled to endemic regions, disease due to *P. marneffeii* should be considered, along with viral encephalitis, tuberculosis, and cryptococcal meningoen- cephalitis. Given the long culture incubation time and high disease mortality, heightened clinical suspicion and prompt em- pirical treatment with a CNS-penetrating antifungal drug, such as amphotericin B, are critical in the management of these patients.

Acknowledgments

Financial support. Fogarty International Clinical Research Fellowship (T.L.) and Wellcome Trust UK Major Overseas Programmes (T.L., J.F., and J.N.D.).

Potential conflicts of interest. All authors: no conflicts.

References

1. Vanittanakom N, Cooper CR, Fisher MC, Sirisanthana T. *Penicillium marneffeii* infection and recent advances in the epidemiology and molecular biology aspects. *Clin Microbiol Rev* **2006**; 19(1):95–110.
2. Imwidthaya P. Update of penicilliosis marneffeii in Thailand. *Mycopathologia* **1994**; 127:135–137.
3. Hilmarsdottir I, Coutellier A, Elbaz J, et al. A French case of laboratory-acquired disseminated *Penicillium marneffeii* infection in a patient with AIDS. *Clin Infect Dis* **1994**; 19:357–358.
4. Duong TA. Infection due to *P. marneffeii*, an emerging pathogen: review of 155 reported cases. *Clin Infect Dis* **1996**; 23:125–130.
5. Supparatpinyo K, Khamwan C, Baosoung V, Nelson KE, Sirisanthana T. Disseminated *Penicillium marneffeii* infection in southeast asia. *Lancet* **1994**; 344(8915):110–113.
6. Noritomi DT, Bud GL, Beer I, Da Silva AS, De Cleva R, Gama-Rodrigues JJ. Multiple brain abscesses due to *Penicillium* spp infection. *Rev Inst Med Trop Sao Paulo* **2005**; 47(3):167–170.
7. Le T, Hong Chau TT, Kim Cuc NT, et al. AIDS-associated *Cryptococcus neoformans* and *Penicillium marneffeii* coinfection: a therapeutic dilemma in resource-limited settings. *Clin Infect Dis* **2010**; 51(9):e65–e68.
8. Huang S, Harris LS. Acute disseminated penicilliosis. *Am J Clin Pathol* **1963**; 39:167–174.
9. Kantarcioglu AS, Apaydin H, Yucel A, et al. Central nervous system infection due to *Penicillium chrysogenum*. *Mycoses* **2004**; 47:242–248.
10. Liratsopoulos G, Ellis M, Nerringer R, Denning DW. Invasive infection due to *Penicillium* species other than *P. marneffeii*. *J Infect* **2002**; 45: 184–195.