

THE GENERALITY OF CLOSED G_2 SOLITONS

ROBERT L. BRYANT

ABSTRACT. The local generality of the space of solitons for the Laplacian flow of closed G_2 -structures is analyzed, and it is shown that the germs of such structures depend, up to diffeomorphism, on 16 functions of 6 variables (in the sense of É. Cartan). The method is to construct a natural exterior differential system whose integral manifolds describe such solitons and to show that it is involutive in Cartan's sense, so that Cartan-Kähler theory can be applied.

Meanwhile, it turns out that, for the more special case of *gradient* solitons, the natural exterior differential system is *not* involutive, and the generality of these structures remains a mystery.

Dedicated, with much admiration and gratitude for his many years of advice, encouragement, and friendship, to H. Blaine Lawson, Jr. on the occasion of his 80th birthday.

CONTENTS

1. Introduction	1
1.1. G_2 -structures	2
1.2. Relation with holonomy	2
1.3. Solitons	3
2. Basic Identities	4
2.1. Expression in differential forms	4
3. Formulation as an exterior differential system	5
3.1. A differential ideal	5
3.2. Involutivity	7
4. The gradient case	9
4.1. Formulation	9
4.2. Non-involutivity	10
References	10

1. INTRODUCTION

For the necessary background on G_2 -structures and the notation and conventions that this article uses, the reader may consult [2] and, especially, [3].

Date: April 30, 2022.

1991 Mathematics Subject Classification. 53E99, 53C29.

Key words and phrases. G_2 -structures, solitons.

Thanks to the Simons Foundation for its support via the Simons Collaboration Grant “Special Holonomy in Geometry, Analysis, and Physics”.

1.1. G_2 -structures. A (smooth) G_2 -structure on a 7-manifold M is a (smooth) 3-form $\phi \in \Omega^3(M)$ that is *definite* in the sense that, for any nonzero tangent vector $v \in T_x M$, the 7-form $(v \lrcorner \phi) \wedge (v \lrcorner \phi) \wedge \phi \in \Lambda^7(T_x^* M)$ is also nonzero. The space of definite 3-forms on M^7 will be denoted $\Omega_+^3(M)$.

Given a $\sigma \in \Omega_+^3(M)$, there is a unique metric g_σ and orientation $*_\sigma$ (i.e., the Hodge star operator defined by g_σ and the orientation) such that

$$(1.1) \quad (X \lrcorner \sigma) \wedge (Y \lrcorner \sigma) \wedge \sigma = 6\langle X, Y \rangle_{g_\sigma} *_\sigma 1$$

for all vector fields X and Y on M .

It is not difficult to prove the first-order 6-form identity [2]

$$(1.2) \quad (*_\sigma d(*_\sigma \sigma)) \wedge *_\sigma \sigma + (*_\sigma d\sigma) \wedge \sigma = 0.$$

In particular, the equations $d\sigma = d(*_\sigma \sigma) = 0$ represent only $35 + 21 - 7 = 49$ first-order equations on a G_2 -structure σ . (It is easy to see that these 49 equations are independent, i.e., there are no further algebraic relations between σ , $d\sigma$ and $*_\sigma d(*_\sigma \sigma)$ beyond those implied by (1.2).)

1.2. Relation with holonomy. Fernández and Gray [7] proved that, if σ and $*_\sigma \sigma$ are both closed forms, then σ is g_σ -parallel. This, in particular, implies that the holonomy of g_σ is isomorphic to a subgroup of $G_2 \subset \text{SO}(7)$. Conversely, if the holonomy of a metric g on M^7 is contained in G_2 , then there exists a definite 3-form σ (unique up to replacement by $-\sigma$ if the holonomy of g is isomorphic to G_2) satisfying $d\sigma = d(*_\sigma \sigma) = 0$ and such that $g = g_\sigma$. This characterization has been essential in all constructions of metrics on 7-manifolds with holonomy G_2 , indeed, in even proving their existence.

While it is easy to write down a local formula for the ‘generic’ closed G_2 -structure σ , the full equations $d\sigma = d*_\sigma \sigma = 0$ are highly nonlinear (as well as being overdetermined). In [2], it was shown how to interpret these equations as an involutive exterior differential system (see [4]) and analyze their local generality in the sense of Élie Cartan. In particular, it was shown that the ‘general’ solution, up to diffeomorphism, depends on six functions of six variables.

One way of expressing this in terms that are, perhaps, more familiar is the following: Suppose that one considers the space \mathcal{J}^k of k -jets of germs of solutions of $d\sigma = d*_\sigma \sigma = 0$ on a neighborhood of $0 \in \mathbb{R}^7$ with the property that the standard coordinates on \mathbb{R}^7 are geodesic normal coordinates for g_σ at 0 and the value of σ at 0 takes a standard form¹. Let $\delta_k = \dim \mathcal{J}^k - \dim \mathcal{J}^{k-1}$ and consider the Poincaré series $P(t) = \delta_0 + \delta_1 t + \delta_2 t^2 + \dots$. Then

$$(1.3) \quad P(t) = t^2 \left(\frac{14}{(1-t)^2} + \frac{21}{(1-t)^3} + \frac{21}{(1-t)^4} + \frac{15}{(1-t)^5} + \frac{6}{(1-t)^6} \right).$$

In particular, the ‘leading term’ contributing to the growth of δ_k , namely $6t^2/(1-t)^6$, is the same, up to a degree shift, as the ‘leading term’ describing the growth of k -jets of 6 functions of 6 variables.

¹This nearly removes the diffeomorphism ambiguity. In order to completely remove it, one would have to quotient by the natural action of G_2 on these jet spaces.

1.3. Solitons. D. Joyce pioneered the approach to constructing compact examples of metrics with holonomy G_2 by starting with a ‘soft’ construction of a closed G_2 -structure σ on certain 7-manifolds with the property that $d(*_\sigma\sigma)$ had sufficiently small norm (in comparison with other geometric features of g_σ) and showing that, in such situations, one could ‘perturb’ the given closed G_2 -structure σ to a nearby one that was also co-closed. (See the fundamental reference [12] and the original papers cited therein for details.)

The many successes of R. Hamilton’s program for using the Ricci flow in Riemannian geometry inspired the idea of canonically ‘improving’ a closed G_2 -structure σ_0 on a compact 7-manifold M by considering the so-called ‘Laplacian flow’

$$(1.4) \quad \frac{d\sigma}{dt} = \Delta_{g_\sigma} \sigma$$

with σ_0 as initial condition. In particular, a fixed point of this flow, i.e., a closed G_2 -structure on M that satisfies $\Delta_{g_\sigma} \sigma = 0$, would satisfy $d\sigma = d*_\sigma\sigma = 0$ and hence lead to a metric g_σ with holonomy contained in G_2 . It has been shown [6] that the closed G_2 -Laplacian flow on a compact 7-manifold has short-time existence and uniqueness.

Meanwhile, Hitchin [10, 11] observed that the above Laplacian flow is (up to a constant factor) the gradient flow of the *volume functional*

$$\mathcal{V}(\sigma) = \int_M \sigma \wedge *_\sigma\sigma$$

on the space $\mathcal{Z}(M, \gamma) \subset \Omega_+^3(M)$ of closed G_2 -structures in a fixed deRham cohomology class $\gamma \in H_{dR}^3(M)$. Hitchin has shown that any critical points of \mathcal{V} in $\mathcal{Z}(M, \gamma) \subset \Omega_+^3(M)$ must be local maxima.

However, there need not be any critical points even if the flow exists for all time, as the example in Remark 18 of [3] shows.

It is not clear just what sort of singularities one might encounter in the Laplacian flow, and one strategy for gaining some intuition, based on the strategy that worked in the case of Ricci flow, is to study the so-called ‘soliton’ solutions.

Definition 1. Let λ be a fixed real constant. A pair (σ, X) , where σ is a closed G_2 -structure on M^7 and X is a vector field on M^7 , is a λ -soliton for the G_2 -Laplacian flow if

$$(1.5) \quad \Delta_\sigma \sigma = \lambda \sigma + \mathcal{L}_X \sigma.$$

If, in addition, $X = \nabla^{g_\sigma} f$ for some function f on M , the pair (σ, X) will be said to be a *gradient λ -soliton*.

It is not difficult to show that, if (σ, X) is a λ -soliton for the G_2 -Laplacian flow on M and $\Phi_\tau : M \rightarrow M$ is the time- τ flow of X , then the time-dependent G_2 -structure

$$(1.6) \quad \phi(t) = e^{\lambda t} \Phi_{f(t, \lambda)}^* \sigma$$

where $f(t, 0) = t$ and $f(t, \lambda) = (1 - e^{-\lambda t})/\lambda$ when $\lambda \neq 0$, satisfies the initial condition $\phi(0) = \sigma$ and the Laplacian flow equation

$$(1.7) \quad \frac{d\phi}{dt} = \Delta_{g_\phi} \phi.$$

Many explicit examples of such λ -solitons are now known. Most of the known examples are constructed using some version of ‘dimensional reduction’, i.e., assuming some sort of symmetry, thereby reducing the problem to geometric data on a lower dimensional manifold, see [1, 8, 9, 13, 14].

The goal of this article is to investigate the local generality, in É. Cartan’s sense, of the space of λ -solitons. A straightforward count of equations reveals that, locally, this is 70 equations for the 42 unknowns in (σ, X) , so the local generality of the solutions of these equations is not at all clear. Moreover, the system is degenerate (in particular, every direction is characteristic) because it is evidently invariant under diffeomorphisms.

2. BASIC IDENTITIES

In this section, I will assume that (σ, X) is a λ -soliton for the G_2 -Laplacian flow equation on M^7 and explain how to express this condition naturally in terms of an exterior differential system.

2.1. Expression in differential forms. Because of the fundamental identity (1.2), the closedness of σ implies that $(*_\sigma d(*_\sigma \sigma)) \wedge *_\sigma \sigma = 0$, so there is a 2-form τ that satisfies

$$(2.1) \quad d(*_\sigma \sigma) = \tau \wedge \sigma = -*_\sigma \tau.$$

(See §4.6 of [3].) Moreover, τ satisfies

$$(2.2) \quad d\tau = \Delta_\sigma \sigma = \lambda \sigma + \mathcal{L}_X \sigma = \lambda \sigma + d(X \lrcorner \sigma),$$

which can be rewritten as

$$(2.3) \quad d(\tau - X \lrcorner \sigma) = \lambda \sigma.$$

As explained in §2.6 of [3], every 2-form $\beta \in \Omega^2(M)$ can be written uniquely as a sum $\beta = \beta_\tau + \beta_{14}$ where $*_\sigma \beta_{14} = -\beta_{14} \wedge \sigma$ while $*_\sigma \beta_\tau = \frac{1}{2} \beta_\tau \wedge \sigma$. In what follows, set $\beta = \tau - X \lrcorner \sigma$, so that $\beta_{14} = \tau$ and $\beta_\tau = -X \lrcorner \sigma$. Then (2.3) can be rewritten as the 3-form equation

$$(2.4) \quad 0 = d\beta - \lambda \sigma.$$

We already have the 4-form equation

$$(2.5) \quad 0 = d\sigma$$

since σ is closed. Using the above algebraic identities, (2.1) can be written as a 5-form equation

$$(2.6) \quad 0 = d(*_\sigma \sigma) - \tau \wedge \sigma = d(*_\sigma \sigma) + \frac{2}{3} *_\sigma \beta - \frac{1}{3} \beta \wedge \sigma.$$

Finally, taking the exterior derivative of this last equation and using the equations found so far, we have the 6-form equation

$$(2.7) \quad 0 = d(*_\sigma \beta).$$

The virtue of the equations (2.4–7) is that they only involve the constant λ , the definite 3-form σ (and its algebraically associated Hodge star operator), and the 2-form β . In fact, this is enough to recover the structure of a λ -soliton for the closed G_2 -Laplacian flow.

Proposition 1. *Let λ be a constant, and let $\sigma \in \Omega_+^3(M)$ and $\beta \in \Omega^2(M)$ satisfy equations (2.4–7). Then for the unique vector field X on M such that $\beta_7 = -X \lrcorner \sigma$, the pair (σ, X) is a λ -soliton for the closed G_2 -Laplacian flow. Conversely, if (σ, X) is a λ -soliton for the closed G_2 -Laplacian flow, then setting $\beta = -*_\sigma d(*_\sigma \sigma) - X \lrcorner \sigma$ defines a pair (σ, β) that satisfies (2.4–7).*

Proof. This is a matter of unwinding the definitions. We have $d\sigma = 0$, and then equation (2.6) implies that $\tau = \beta_{14}$ and that $d(*_\sigma \sigma) = \tau \wedge \sigma$, so that $d\tau = \Delta_\sigma \sigma$. Then (2.4) becomes (2.3). The converse follows in the same way. \square

Remark 1. The reader may wonder why (2.7) was included, since it is a consequence of the other three equations just by applying the exterior derivative to (2.6). Moreover, when λ is nonzero, (2.5) follows from (2.4), so it might seem that (2.5) could be omitted as well, at least when λ is nonzero.

The reason is that I want to make explicit *all* of the first-order equations satisfied by the pair (σ, β) . The four equations (2.4–7) are the vanishing of a 3-form, a 4-form, a 5-form, and a 6-form, which is potentially

$$35 + 35 + 21 + 7 = 98$$

independent first order equations. However, the identity (1.2), shows that these 98 equations are *not* independent; they ‘overlap’ by at least 7 equations.

In fact, the overlap is *exactly* the 7 equations of the fundamental identity (1.2) since it is easy to see that, for a generic $\sigma \in \Omega_+^3(M)$ and $\beta \in \Omega^2(M)$, the only relation among the forms

$$d\beta, d\sigma, d(*_\sigma \sigma), d(*_\sigma \beta)$$

is the fundamental identity relating the middle two. Thus, the equations (2.4–7) constitute exactly 91 independent (quasi-linear) first order equations on the pair (σ, β) , a fact to which I will return.

Indeed, I am going to argue that this system of 91 equations is *involution* in É. Cartan’s sense, so that the Cartan-Kähler Theorem can be applied to prove local existence of solutions and describe their ‘degree of generality’. By contrast, the ‘determined’ system of $35 + 21 = 56$ first-order equations for (σ, β) represented by combining (2.4) and (2.6) is certainly *not* involutive.

3. FORMULATION AS AN EXTERIOR DIFFERENTIAL SYSTEM

I now want to explain how the above system of 91 first-order partial differential equations for the pair (σ, β) , which is a section of a smooth bundle of fiber rank $35 + 21 = 56$ over M^7 , can be interpreted as defining an exterior differential system for such pairs.

3.1. A differential ideal. Let $\Lambda_+^3(T^*M) \subset \Lambda^3(T^*M)$ denote the open subset of the bundle of 3-forms on M^7 whose (local) sections are the (local) G_2 -structures on M . Similarly, let $\Lambda^2(T^*M)$ denote the bundle of 2-forms on M .

Let $X \subset \Lambda^3(T^*M) \oplus \Lambda^2(T^*M)$ denote the set of pairs (s, b) with $s \in \Lambda_+^3(T_m^*M)$ and $b \in \Lambda^2(T_m^*M)$ for some $m \in M$. This X is open in $\Lambda^3(T^*M) \oplus \Lambda^2(T^*M)$ and is a smooth fiber bundle over M with a submersive base projection $\pi : X \rightarrow M$, with $\pi(s, b) = m$. Let $\pi^* : \Lambda^k(T^*M) \rightarrow \Lambda^k(T^*X)$ denote the induced ‘pullback’ mapping.

There exist natural ‘tautological’ forms on X that are defined as follows: For each $(s, b) \in X$, let $\mathfrak{s}_{(s,b)} = \pi^*(s)$ and let $\mathfrak{b}_{(s,b)} = \pi^*(b)$. Then \mathfrak{s} (respectively, \mathfrak{b}) is a smooth 3-form (respectively, 2-form) on X with the ‘reproducing’ property that, for any section $(\sigma, \beta) : M \rightarrow X$, there holds $(\sigma, \beta)^*(\mathfrak{s}) = \sigma$ and $(\sigma, \beta)^*(\mathfrak{b}) = \beta$. Moreover, because exterior derivative commutes with pullback, the identities $(\sigma, \beta)^*(d\mathfrak{s}) = d\sigma$ and $(\sigma, \beta)^*(d\mathfrak{b}) = d\beta$ hold as well.

I will need an extension of this construction: Since, for any $s \in \Lambda_+^3(T_m^*M)$, there is a well-defined Hodge star operator $*_s : \Lambda^p(T_m^*M) \rightarrow \Lambda^{7-p}(T_m^*M)$, there also exists a smooth 4-form \mathfrak{t} (respectively, 5-form, \mathfrak{p}) on X with the property that $\mathfrak{t}_{(s,b)} = \pi^*(*_s s)$ (respectively, $\mathfrak{p}_{(s,b)} = \pi^*(*_s b)$). These two forms have the ‘reproducing’ property that, for any section $(\sigma, \beta) : M \rightarrow X$, there holds $(\sigma, \beta)^*(\mathfrak{t}) = *_\sigma \sigma$ and $(\sigma, \beta)^*(\mathfrak{p}) = *_\sigma \beta$.

Finally, there exists a smooth 7-form ω on X with the property that $\omega_{(s,b)} = \pi^*(*_s 1)$ that will be used below, so I introduce it now.

Note that all of the forms \mathfrak{s} , \mathfrak{t} , \mathfrak{b} , \mathfrak{p} , and ω are π -semibasic, i.e., their interior product with any π -vertical vector field on X vanishes identically.

Now, by construction, for any constant λ and any section $(\sigma, \beta) : M \rightarrow X$, one has

$$(3.1) \quad \begin{aligned} (\sigma, \beta)^*(d\mathfrak{b} - \lambda \mathfrak{s}) &= d\beta - \lambda \sigma \\ (\sigma, \beta)^*(d\mathfrak{s}) &= d\sigma \\ (\sigma, \beta)^*(d\mathfrak{t} + \frac{2}{3} \mathfrak{p} - \frac{1}{3} \mathfrak{b} \wedge \mathfrak{s}) &= d(*_\sigma \sigma) + \frac{2}{3} *_\sigma \beta - \frac{1}{3} \beta \wedge \sigma \\ (\sigma, \beta)^*(d\mathfrak{p}) &= d(*_\sigma \beta) \end{aligned}$$

This motivates defining the differential forms $\Upsilon_i \in \Omega^i(X)$,

$$(3.2) \quad \begin{aligned} \Upsilon_3 &= d\mathfrak{b} - \lambda \mathfrak{s}, \\ \Upsilon_4 &= d\mathfrak{s}, \\ \Upsilon_5 &= d\mathfrak{t} + \frac{2}{3} \mathfrak{p} - \frac{1}{3} \mathfrak{b} \wedge \mathfrak{s}, \\ \Upsilon_6 &= d\mathfrak{p}. \end{aligned}$$

Note the identities

$$(3.3) \quad \begin{aligned} d\Upsilon_3 &= -\lambda \Upsilon_4, \\ d\Upsilon_4 &= 0, \\ d\Upsilon_5 &= \frac{2}{3} \Upsilon_6 - \frac{1}{3} \Upsilon_3 \wedge \mathfrak{s} - \frac{1}{3} \mathfrak{b} \wedge \Upsilon_4, \\ d\Upsilon_6 &= 0, \end{aligned}$$

which imply that the algebraic ideal \mathcal{I}_λ in $\Omega^*(X)$ generated by $\Upsilon_3, \Upsilon_4, \Upsilon_5, \Upsilon_6$ is differentially closed. Consequently, \mathcal{I}_λ is an exterior differential system on X .

The interest in \mathcal{I}_λ is explained by the following result.

Proposition 2. *A section $(\sigma, \beta) : M \rightarrow X$ is an integral manifold of \mathcal{I}_λ if and only if it comes from a λ -soliton for the closed G_2 -Laplacian flow. In particular, a 7-dimensional integral manifold of \mathcal{I}_λ that is transverse to the fibers of $\pi : X \rightarrow M$ is locally the graph of a section $(\sigma, \beta) : M \rightarrow X$ that comes from a λ -soliton for the closed G_2 -Laplacian flow.*

Proof. It has been shown that a pair (σ, β) comes from a λ -soliton for the closed G_2 -Laplacian flow if and only if the pair satisfies the equations (2.4–7). Since this vanishing is exactly the vanishing of the righthand sides of (3.1), the definitions

given in (3.2) and the definition of \mathcal{I}_λ show that this vanishing is equivalent to the section (σ, β) being a 7-dimensional integral manifold (necessarily transverse to the π -fibers) of \mathcal{I}_λ . The rest of the statement follows immediately. \square

3.2. Involutivity. I can now state the main result of this article.

Theorem 1. *The exterior differential system \mathcal{I}_λ with independence condition ω is involutive, with Cartan characters*

$$(s_0, s_1, s_2, s_3, s_4, s_5, s_6, s_7) = (0, 0, 1, 3, 7, 15, 23, 7).$$

Proof. The proof will be by Cartan's Test for involutivity (see, for example, [4]).

The first thing to check for each $(s, b) \in X$ is the dimension of the space of 7-dimensional integral elements $E \subset T_{(s,b)}X$ of \mathcal{I}_λ on which ω is non-vanishing. I claim that this dimension is $301 = 392 - 91$.

To see this, choose a splitting $T_{(s,b)}X = \ker \pi'(s, b) + W$, where $\pi'(s, b) : W \rightarrow T_{\pi(s,b)}M$ is an isomorphism. Then an integral element $E \subset T_{(s,b)}X$ on which ω is non-vanishing can be thought of as the graph of a linear map $L : W \rightarrow \ker \pi'(s, b)$, i.e., it can be identified with an element of the vector space $\text{Hom}(W, \ker \pi'(s, b))$. Because the four tautological forms \mathfrak{s} , \mathfrak{b} , \mathfrak{t} , and \mathfrak{p} are π -semi-basic, it follows that the induced mapping of $\text{Hom}(W, \ker \pi'(s, b))$ into the 98-dimensional vector space

$$\Lambda^3(W^*) \oplus \Lambda^4(W^*) \oplus \Lambda^5(W^*) \oplus \Lambda^6(W^*)$$

induced by evaluating $(\Upsilon_3, \Upsilon_4, \Upsilon_5, \Upsilon_6)$ on the graph of $L \in \text{Hom}(W, \ker \pi'(s, b))$ is an affine mapping. Moreover, its image lies in the 91-dimensional subspace cut out by the fundamental identity. Finally, looking at the 'leading terms' of $(\Upsilon_3, \Upsilon_4, \Upsilon_5, \Upsilon_6)$, i.e., $(d\mathfrak{b}, d\mathfrak{s}, d\mathfrak{t}, d\mathfrak{p})$ and considering their interpretation as

$$(d\beta, d\sigma, d(*_\sigma\sigma), d(*_\sigma\beta)),$$

one sees that the leading order part is a linear mapping that is surjective onto this 91-dimensional subspace.

Consequently, the set $\mathcal{V}(\mathcal{I}_\lambda, \omega)$ of admissible integral elements of \mathcal{I}_λ at $(s, b) \in X$ is a (nonempty) affine subspace of $\text{Hom}(W, \ker \pi'(s, b))$ of dimension $S = 56 \times 7 - 91 = 392 - 91 = 301$. Indeed $\mathcal{V}(\mathcal{I}_\lambda, \omega)$ is a smooth submanifold of $\text{Gr}(7, TX)$, the Grassmann bundle of 7-dimensional subspaces of the tangent spaces to X .

It remains to compute the Cartan characters and apply Cartan's Test. Now that it has been established that the space of admissible integral elements of \mathcal{I}_λ at each point is nonempty, the Cartan characters can be computed by considering just the leading terms listed above.

First, recall that it was established in [2] that the Cartan characters of the (involutive) equation $D(\sigma) = (d\sigma, d(*_\sigma\sigma)) = (0, 0)$ relative to any flag are

$$(s_0, s_1, s_2, s_3, s_4, s_5, s_6, s_7) = (0, 0, 0, 1, 4, 10, 13, 7),$$

Meanwhile, once σ is fixed, the characters of the involutive first-order system $D(\beta) = (d\beta, d(*_\sigma\beta)) = (0, 0)$ for β a 2-form on a manifold with a specified G_2 -structure σ are

$$(s_0, s_1, s_2, s_3, s_4, s_5, s_6, s_7) = (0, 0, 1, 2, 3, 5, 10, 0).$$

It follows that there exists an integral flag of the combined system with characters

$$(s_0, s_1, s_2, s_3, s_4, s_5, s_6, s_7) = (0, 0, 1, 3, 7, 15, 23, 7).$$

It follows that $\sum_{k=0}^7 k s_k = 301 = S$, the dimension of the space of integral elements of \mathcal{I}_λ at (s, b) . Thus, equality holds in Cartan's Test, which implies that such a flag is Cartan-regular. Consequently, the system is involutive, as was to be shown. \square

Remark 2. Since M is assumed to be smooth, it also carries a real-analytic structure (in fact, many, but all of them are equivalent), and X will naturally inherit a real-analytic structure once one is chosen on M . Because of the nature of their construction, the forms $\Upsilon_3, \Upsilon_4, \Upsilon_5, \Upsilon_6$ will be real-analytic with respect to the real-analytic structure on X . Hence \mathcal{I}_λ is real-analytic with respect to any real-analytic structure on X .

Corollary 1. *For any $(s, b) \in X$ with $\pi(s, b) = m \in M$, there exists a λ -soliton (σ, β) for the closed G_2 -Laplacian flow on an m -neighborhood $U \subset M$ such that $(\sigma_m, \beta_m) = (s, b)$. In fact, for any 7-dimensional integral element $E \subset T_{(s,b)}X$ of \mathcal{I}_λ on which ω is non-vanishing, there exists such a $(\sigma, \beta) : U \rightarrow X$ such that $(\sigma, \beta)'(s, b)(T_m M) = E$.*

Proof. In light of Theorem 1 and Remark 2, Corollary 1 follows immediately from the Cartan-Kähler Theorem. \square

Remark 3 (Generality in Harmonic Coordinates). Since the last nonzero character of \mathcal{I}_λ is $s_7 = 7$, the naïve statement is that local λ -solitons for the closed G_2 -Laplacian flow depend on ‘seven functions of seven variables’ in Cartan’s terminology². However, this reflects the fact that the equations are invariant under the full diffeomorphism group in dimension 7, which itself is described (locally) by specifying seven functions of seven variables.

There are a number of methods of, at least partially, ‘normalizing’ the local solutions by choosing special coordinates. The simplest one perhaps, is to ask how general the local solutions are when presented in harmonic coordinates for the associated metric g_σ . One way to see this is to specialize to the case $M = \mathbb{R}^7$ with standard coordinates $u = (u^i)$ and impose the conditions $d(*_\sigma(du^i)) = 0$. This can be added to the exterior differential system by defining the seven 6-forms

$$(3.4) \quad \Phi^i = \pi^*(*_s(du^i)) \in \Lambda^6(T_{(s,b)}^*X)$$

and then defining the extended ideal \mathcal{I}_λ^+ generated by \mathcal{I}_λ and the seven closed 7-forms $d\Phi^i$. Following the same argument as that used in Theorem 1, one can show that the system \mathcal{I}_λ^+ on X is also involutive, but now with characters

$$(3.5) \quad (s_0, s_1, s_2, s_3, s_4, s_5, s_6, s_7) = (0, 0, 1, 3, 7, 15, 30, 0).$$

One can interpret the local integrals of \mathcal{I}_λ^+ as λ -solitons for the closed G_2 -Laplacian flow that are presented in harmonic coordinates. In Cartan’s terminology, such solitons depend locally on $s_6 = 30$ functions of six variables. However, the choice of a local harmonic function, given σ , depends on 2 functions of six variables, so a choice of a harmonic coordinate system depends on $2 \times 7 = 14$ functions of six variables. Thus, one can argue that local solitons (σ, β) depend on $16 = 30 - 14$ functions of six variables, in the sense that the rate of growth of the dimensions of the k -jets of solutions up to diffeomorphism is the same as the rate of growth of 16 functions of 6 variables.

²The Cartan characters s_i indicate how much essential choice there is in building up a solution to the equations via a sequence of (possibly underdetermined) Cauchy problems.

Remark 4 (Analyticity). Another reason for considering the presentation of λ -solitons (σ, β) in harmonic coordinates is that it makes clear the regularity of such structures. Because the last nonzero character of \mathcal{I}_λ^+ is $s_6 = 30$, it follows [5, Chapter V] that the complex characteristic variety $\Xi_E \subset \mathbb{P}((E \otimes \mathbb{C})^*) \simeq \mathbb{C}\mathbb{P}^6$ is a subvariety of (complex) dimension 5 for every integral element E . Because of the G_2 -invariance of the system and the fact that G_2 acts transitively on (real) 2-planes in \mathbb{R}^7 , the only possibility for the support of this characteristic variety for a given solution (σ, β) on a neighborhood of a point $u \in \mathbb{R}^7$ is the set of complex co-vectors in $\mathbb{P}((T_u^*\mathbb{R}^7) \otimes \mathbb{C})$ that are null for g_{σ_u} . In particular, the complex characteristic variety has no real points, implying that the associated PDE system is (overdetermined) elliptic. Consequently, λ -solitons that are $C^{1,\alpha}$ in harmonic coordinates (for some $\alpha > 0$) must be real-analytic in those coordinates [15].

4. THE GRADIENT CASE

In this final section, I will discuss what is known about the special case that the λ -soliton (σ, X) is a *gradient* soliton, i.e., $X = \nabla^{g_\sigma} f$ for some function f . This case is interesting partly because, in the case of solitons for the Ricci flow, particularly in dimension 3, the gradient solitons are the most important.

Remark 5 (Ricci solitons in dimension 3). Just for comparison, it might be interesting to note that the generality analysis for general Ricci solitons in dimension 3, i.e., pairs (g, X) of a metric g and a vector field X on a manifold M^3 that satisfy

$$(4.1) \quad \text{Ric}(g) = \lambda g + \mathcal{L}_X g,$$

shows that, up to local diffeomorphism, the local solutions of this system depend on $s_2 = 6$ functions of two variables. If one adds the ‘gradient condition’ that $X = \nabla^g f$ for some function f , then one sees that, up to local diffeomorphism, the local solutions satisfying this extra condition depend on $s_2 = 2$ functions of two variables. In both cases, the associated exterior differential system is involutive. See [4, §5.5] for the gradient case.

Many explicit examples of gradient λ -solitons are known, as can be seen by looking at the examples constructed in the articles [1, 8, 9, 13, 14]. However, as of this writing, it is not known what their generality is in the sense of É. Cartan. I will conclude this article by explaining where the difficulty lies.

4.1. Formulation. The gradient λ -soliton case, where $X = \nabla^{g_\sigma} f$ for some function f , is locally equivalent to requiring that X^\flat , the 1-form that is g_σ -dual to the vector field X , be closed.

Starting with the algebraic identity of 5-forms

$$(4.2) \quad (X \lrcorner \sigma) \wedge \sigma = 2X^\flat \wedge *_\sigma \sigma$$

and applying the exterior derivative to the left hand side of (4.2) gives

$$(4.3) \quad d((X \lrcorner \sigma) \wedge \sigma) = d(X \lrcorner \sigma) \wedge \sigma = (d\tau - \lambda\sigma) \wedge \sigma = 0,$$

since $d(*_\sigma \sigma) = \tau \wedge \sigma$ and $d\sigma = 0$ implies $d\tau \wedge \sigma = 0$. Meanwhile applying the exterior derivative to the right hand side of (4.2) (which is closed because of (4.3)) gives

$$(4.4) \quad 0 = 2d(X^\flat) \wedge *_\sigma \sigma - 2X^\flat \wedge \tau \wedge \sigma.$$

In particular, if $d(X^\flat) = 0$, which is the condition for a gradient λ -soliton, it follows that $X^\flat \wedge \tau \wedge \sigma = 0$, which is an algebraic relation on X and τ that is equivalent to

$$(4.5) \quad X \lrcorner \tau = 0,$$

i.e., the vanishing of a 1-form that is bilinear in X and τ .

4.2. Non-involutivity. The condition $d(X^\flat) = 0$ is expressed in terms of β as

$$(4.6) \quad d(*_\sigma(\beta \wedge *_\sigma \sigma)) = 0.$$

Thus, it is natural to study the gradient case by augmenting the differential ideal \mathcal{I}_λ by adding to it the closed 2-form dz , where z is the 1-form on X defined as

$$z_{(s,b)} = \pi^*(*_s(b \wedge *_s s)).$$

Unfortunately, as (4.5) shows, this augmented differential ideal does not have any admissible integral elements at any $(s, b) \in X$ for which the 6-form $(*_s(b \wedge *_s s)) \wedge b \wedge s$ is nonzero. Hence, the augmented ideal is not involutive.

The standard ‘next step’ in such a case is to pull back the augmented differential ideal to the (nonsmooth) subvariety $Z \subset X$ on which $z \wedge b \wedge s = 0$ and check that exterior differential system on Z for involutivity. Unfortunately, calculation shows that that exterior differential system is not involutive either, even on the smooth locus of Z .

Two successive applications of the standard prolongation procedure to this system does not yield an involutive exterior differential system, and the calculations become increasingly complicated. So far, no method has been found that leads to an involutive system whose solutions correspond to the gradient λ -solitons.

As a result, the generality of the gradient λ -solitons remains unknown as of this writing. All we have is a collection of the various examples already mentioned to show that gradient λ -solitons do, indeed, exist.

REFERENCES

- [1] G. Ball, *Quadratic closed G_2 -structures*, arXiv:2006.14155 4, 9
- [2] R. Bryant, *Metrics with exceptional holonomy*, *Annals of Mathematics* (2), **126** (1987), 525–576. 1, 2, 7
- [3] ———, *Some remarks on G_2 -structures*, in *Proceedings of Gökova Geometry-Topology Conference 2005*, edited by S. Akbulut, T. Onder, and R.J. Stern. Gökova Geometry/Topology Conference (GGT), Gökova, 2006. See also arXiv:math/0305124 1, 3, 4
- [4] ———, *Notes on exterior differential systems*, arXiv:1405.3116. 2, 7, 9
- [5] R. Bryant, S.-s. Chern, R. Gardner, H. Goldschmidt, and P. Griffiths, *Exterior Differential Systems*, *Mathematical Sciences Research Institute Publications* **18** (1991). 9
- [6] R. Bryant and F. Xu, *Laplacian flow for closed G_2 -structures: short time behavior*, arXiv:1101.2004 3
- [7] M. Fernández and A. Gray, *Riemannian manifolds with structure group G_2* , *Ann. Mat. Pura Appl.* (4) **132** (1982), 19–45. 2
- [8] U. Fowdar, *S^1 -invariant Laplacian flow*, to appear in the *Journal of Geometric Analysis*, arXiv:2007.05130 4, 9
- [9] M. Haskins and J. Nordström, *Cohomogeneity-one solitons in Laplacian flow: local, smoothly-closing and steady solitons*, arXiv:2112.09095 4, 9
- [10] N. Hitchin, *The geometry of three-forms in six and seven dimensions*, arXiv:math/0010054. 3
- [11] N. Hitchin, *Stable forms and special metrics*, in *Global differential geometry: the mathematical legacy of Alfred Gray* (Bilbao, 2000), 70–89, *Contemp. Math.* **288**, Amer. Math. Soc., Providence, RI, 2001. 3

- [12] D. Joyce, *Compact manifolds with special holonomy*, Oxford Mathematical Monographs, Oxford University Press, Oxford, 2000. MR 2001k:53093 3
- [13] J. Lauret, *Laplacian flow of homogeneous G_2 -structures and its solitons*, Proc. London Math. Soc. **114** (2017), 527–560. 4, 9
- [14] J. Lotay, *Geometric flows of G_2 -structures*, To appear in a forthcoming volume of the Fields Institute Communications entitled *Lectures and Surveys on G_2 manifolds and related topics*. Also, see arXiv:1810.13417. 4, 9
- [15] M. Taylor, *Regularity of solutions to nonlinear overdetermined elliptic PDE*, <https://mtaylor.web.unc.edu/notes>, Item 5.15. 9

DUKE UNIVERSITY MATHEMATICS DEPARTMENT, PO BOX 90320, DURHAM, NC 27708-0320
E-mail address: bryant@math.duke.edu
URL: <http://www.math.duke.edu/~bryant>