

## Dear Author

Here are the proofs of your article.

- You can submit your corrections **online**, via **e-mail** or by **fax**.
- For **online** submission please insert your corrections in the online correction form. Always indicate the line number to which the correction refers.
- You can also insert your corrections in the proof PDF and **email** the annotated PDF.
- For **fax** submission, please ensure that your corrections are clearly legible. Use a fine black pen and write the correction in the margin, not too close to the edge of the page.
- Remember to note the **journal title**, **article number**, and **your name** when sending your response via e-mail or fax.
- **Check** the metadata sheet to make sure that the header information, especially author names and the corresponding affiliations are correctly shown.
- **Check** the questions that may have arisen during copy editing and insert your answers/corrections.
- **Check** that the text is complete and that all figures, tables and their legends are included. Also check the accuracy of special characters, equations, and electronic supplementary material if applicable. If necessary refer to the *Edited manuscript*.
- The publication of inaccurate data such as dosages and units can have serious consequences. Please take particular care that all such details are correct.
- Please **do not** make changes that involve only matters of style. We have generally introduced forms that follow the journal's style.
- Substantial changes in content, e.g., new results, corrected values, title and authorship are not allowed without the approval of the responsible editor. In such a case, please contact the Editorial Office and return his/her consent together with the proof.
- If we do not receive your corrections **within 48 hours**, we will send you a reminder.
- Your article will be published **Online First** approximately one week after receipt of your corrected proofs. This is the **official first publication** citable with the DOI. **Further changes are, therefore, not possible.**
- The **printed version** will follow in a forthcoming issue.

### Please note

After online publication, subscribers (personal/institutional) to this journal will have access to the complete article via the DOI using the URL:

If you would like to know when your article has been published online, take advantage of our free alert service. For registration and further information, go to:

<http://www.link.springer.com>.

Due to the electronic nature of the procedure, the manuscript and the original figures will only be returned to you on special request. When you return your corrections, please inform us, if you would like to have these documents returned.

**Metadata of the article that will be visualized in OnlineFirst**

---

**Please note: Images will appear in color online but will be printed in black and white.**

---

1	Article Title	<b>Dominant Splice Site Mutations in <i>PIK3R1</i> Cause Hyper IgM Syndrome, Lymphadenopathy and Short Stature</b>
2	Article Sub- Title	
3	Article Copyright - Year	<b>Springer Science+Business Media New York 2016 (This will be the copyright line in the final PDF)</b>
4	Journal Name	Journal of Clinical Immunology
5		Family Name <b>Buckley</b>
6		Particle
7		Given Name <b>Rebecca H.</b>
8		Suffix
9		Organization Duke University Medical Center
10	Corresponding Author	Division Department of Pediatrics, Division of Allergy and Immunology
11		Address 363 Jones Building, Box 2898, Durham 27710, NC, USA
12		Organization Duke University Medical Center
13		Division Department of Immunology
14		Address Durham 27710, NC, USA
15		e-mail buckl003@mc.duke.edu
16		Family Name <b>Petrovski</b>
17		Particle
18		Given Name <b>Slavé</b>
19		Suffix
20	Author	Organization Columbia University
21		Division Institute for Genomic Medicine
22		Address New York 10032, NY, USA
23		e-mail
24		Family Name <b>Parrott</b>
25		Particle
26		Given Name <b>Roberta E.</b>
27	Author	Suffix
28		Organization Duke University Medical Center
29		Division Department of Pediatrics, Division of Allergy and Immunology

30		Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
31		e-mail	
32		Family Name	<b>Roberts</b>
33		Particle	
34		Given Name	<b>Joseph L.</b>
35		Suffix	
36	Author	Organization	Duke University Medical Center
37		Division	Department of Pediatrics, Division of Allergy and Immunology
38		Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
39		e-mail	
40		Family Name	<b>Huang</b>
41		Particle	
42		Given Name	<b>Hongxiang</b>
43		Suffix	
44		Organization	Duke University Medical Center
45	Author	Division	Department of Pediatrics, Division of Allergy and Immunology
46		Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
47		Organization	Southern Medical University
48		Division	Department of Oncology, Nanfang Hospital
49		Address	Guangzhou 510515, Guangdong, China
50		e-mail	
51		Family Name	<b>Yang</b>
52		Particle	
53		Given Name	<b>Jialong</b>
54		Suffix	
55	Author	Organization	Duke University Medical Center
56		Division	Department of Pediatrics, Division of Allergy and Immunology
57		Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
58		e-mail	
59		Family Name	<b>Gorentla</b>
60		Particle	
61		Given Name	<b>Balachandra</b>
62	Author	Suffix	
63		Organization	Duke University Medical Center
64		Division	Department of Pediatrics, Division of Allergy and Immunology

65	Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
66	e-mail	
<hr/>		
67	Family Name	<b>Mousallem</b>
68	Particle	
69	Given Name	<b>Talal</b>
70	Suffix	
71	Organization	Duke University Medical Center
72	Author Division	Department of Pediatrics, Division of Allergy and Immunology
73	Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
74	Organization	Wake Forest University School of Medicine
75	Division	Departments of Internal Medicine and Pediatrics
76	Address	Winston-Salem, NC, USA
77	e-mail	
<hr/>		
78	Family Name	<b>Wang</b>
79	Particle	
80	Given Name	<b>Endi</b>
81	Suffix	
82	Author Organization	Duke University Medical Center
83	Division	Department of Pathology
84	Address	Durham 27710, NC, USA
85	e-mail	
<hr/>		
86	Family Name	<b>Armstrong</b>
87	Particle	
88	Given Name	<b>Martin</b>
89	Suffix	
90	Author Organization	UCB NewMedicines
91	Division	
92	Address	Slough, UK
93	e-mail	
<hr/>		
94	Family Name	<b>McHale</b>
95	Particle	
96	Given Name	<b>Duncan</b>
97	Suffix	
98	Author Organization	UCB NewMedicines
99	Division	
100	Address	Slough, UK
101	e-mail	

102		Family Name	<b>MacIver</b>
103		Particle	
104		Given Name	<b>Nancie J.</b>
105		Suffix	
106		Organization	Duke University Medical Center
107	Author	Division	Department of Pediatrics, Division of Endocrinology
108		Address	Durham 27710, NC, USA
109		Organization	Duke University Medical Center
110		Division	Department of Immunology
111		Address	Durham 27710, NC, USA
112		e-mail	
<hr/>			
113		Family Name	<b>Goldstein</b>
114		Particle	
115		Given Name	<b>David B.</b>
116		Suffix	
117	Author	Organization	Columbia University
118		Division	Institute for Genomic Medicine
119		Address	New York 10032, NY, USA
120		e-mail	
<hr/>			
121		Family Name	<b>Zhong</b>
122		Particle	
123		Given Name	<b>Xiao-Ping</b>
124		Suffix	
125		Organization	Duke University Medical Center
126	Author	Division	Department of Pediatrics, Division of Allergy and Immunology
127		Address	363 Jones Building, Box 2898, Durham 27710, NC, USA
128		Organization	Duke University Medical Center
129		Division	Department of Immunology
130		Address	Durham 27710, NC, USA
131		e-mail	
<hr/>			
132		Received	12 February 2016
133	Schedule	Revised	
134		Accepted	31 March 2016
<hr/>			
135	Abstract	<p>The purpose of this research was to use next generation sequencing to identify mutations in patients with primary immunodeficiency diseases whose pathogenic gene mutations had not been identified. Remarkably, four unrelated patients were found by next generation sequencing to have the same heterozygous mutation in an essential donor splice site of <i>PIK3RI</i> (NM_181523.2:c.1425 + 1G &gt; A) found in three prior reports. All four had the</p>	

Hyper IgM syndrome, lymphadenopathy and short stature, and one also had SHORT syndrome. They were investigated with in vitro immune studies, RT-PCR, and immunoblotting studies of the mutation's effect on mTOR pathway signaling. All patients had very low percentages of memory B cells and class-switched memory B cells and reduced numbers of naïve CD4+ and CD8+ T cells. RT-PCR confirmed the presence of both an abnormal 273 base-pair (bp) size and a normal 399 bp size band in the patient and only the normal band was present in the parents. Following anti-CD40 stimulation, patient's EBV-B cells displayed higher levels of S6 phosphorylation (mTOR complex 1 dependent event), Akt phosphorylation at serine 473 (mTOR complex 2 dependent event), and Akt phosphorylation at threonine 308 (PI3K/PDK1 dependent event) than controls, suggesting elevated mTOR signaling downstream of CD40. These observations suggest that amino acids 435–474 in PIK3R1 are important for its stability and also its ability to restrain PI3K activity. Deletion of Exon 11 leads to constitutive activation of PI3K signaling. This is the first report of this mutation and immunologic abnormalities in SHORT syndrome.

---

136	Keywords separated by ' - '	PIK3R1 splice site mutations - Hyper IgM syndrome - lymphadenopathy - short stature - SHORT syndrome - mTOR pathway - next generation sequencing
<hr/>		
137	Foot note information	The online version of this article (doi:10.1007/s10875-016-0281-6) contains supplementary material, which is available to authorized users.

## Electronic supplementary material

### ESM 1

Figure S1: Sanger sequencing traces of DNA sequence around the *PIK3R1* donor splice site for each patient (1–4) and their parents (M = Mother, F = Father). Both forward (+) and reverse (–) directions are provided. Patients are heterozygous for the variant. The variant was not observed in the unaffected parents. Father of Patient 4 was unavailable. (JPEG 2560 kb)

# Dominant Splice Site Mutations in *PIK3R1* Cause Hyper IgM Syndrome, Lymphadenopathy and Short Stature

Slavé Petrovski<sup>1</sup> · Roberta E. Parrott<sup>2</sup> · Joseph L. Roberts<sup>2</sup> · Hongxiang Huang<sup>2,3</sup> · Jialong Yang<sup>2</sup> · Balachandra Gorentla<sup>2</sup> · Talal Mousallem<sup>2,4</sup> · Endi Wang<sup>5</sup> · Martin Armstrong<sup>6</sup> · Duncan McHale<sup>6</sup> · Nancie J. MacIver<sup>7,8</sup> · David B. Goldstein<sup>1</sup> · Xiao-Ping Zhong<sup>2,8</sup> · Rebecca H. Buckley<sup>2,8</sup>

Received: 12 February 2016 / Accepted: 31 March 2016  
© Springer Science+Business Media New York 2016

**Abstract** The purpose of this research was to use next generation sequencing to identify mutations in patients with primary immunodeficiency diseases whose pathogenic gene mutations had not been identified. Remarkably, four unrelated patients were found by next generation sequencing to have the same heterozygous mutation in an essential donor splice site of *PIK3R1* (NM\_181523.2:c.1425 + 1G > A) found in three prior reports. All four had the Hyper IgM syndrome, lymphadenopathy and short stature, and one also had SHORT syndrome. They were investigated with in vitro immune studies, RT-PCR, and immunoblotting studies of the mutation's effect on mTOR pathway signaling. All patients

had very low percentages of memory B cells and class-switched memory B cells and reduced numbers of naïve CD4+ and CD8+ T cells. RT-PCR confirmed the presence of both an abnormal 273 base-pair (bp) size and a normal 399 bp size band in the patient and only the normal band was present in the parents. Following anti-CD40 stimulation, patient's EBV-B cells displayed higher levels of S6 phosphorylation (mTOR complex 1 dependent event), Akt phosphorylation at serine 473 (mTOR complex 2 dependent event), and Akt phosphorylation at threonine 308 (PI3K/PDK1 dependent event) than controls, suggesting elevated mTOR signaling downstream of CD40. These observations suggest that amino acids 435–474 in *PIK3R1* are important for its stability and also its ability to restrain PI3K activity. Deletion of Exon 11 leads to constitutive activation of PI3K signaling. This is the first report of this mutation and immunologic abnormalities in SHORT syndrome.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10875-016-0281-6) contains supplementary material, which is available to authorized users.

✉ Rebecca H. Buckley  
buckl003@mc.duke.edu

Q1

<sup>1</sup> Institute for Genomic Medicine, Columbia University, New York, NY 10032, USA

Q2

<sup>2</sup> Department of Pediatrics, Division of Allergy and Immunology, Duke University Medical Center, 363 Jones Building, Box 2898, Durham, NC 27710, USA

<sup>3</sup> Department of Oncology, Nanfang Hospital, Southern Medical University, Guangzhou, Guangdong 510515, China

<sup>4</sup> Departments of Internal Medicine and Pediatrics, Wake Forest University School of Medicine, Winston-Salem, NC, USA

<sup>5</sup> Department of Pathology, Duke University Medical Center, Durham, NC 27710, USA

<sup>6</sup> UCB NewMedicines, Slough, UK

<sup>7</sup> Department of Pediatrics, Division of Endocrinology, Duke University Medical Center, Durham, NC 27710, USA

<sup>8</sup> Department of Immunology, Duke University Medical Center, Durham, NC 27710, USA

**Keywords** *PIK3R1* splice site mutations · Hyper IgM syndrome · lymphadenopathy · short stature · SHORT syndrome · mTOR pathway · next generation sequencing

## Introduction 45 Q3

Many critical aspects of immune cell development, differentiation and function are controlled by phosphoinositide 3 kinases (PI3Ks) [1]. Information is accruing about the effects of human gene mutations affecting these molecules. Hyperactivation of the PI3K signaling pathway due to heterozygous gain-of-function mutations in the gene encoding *PIK3CD* has been found by several groups to result in defects in immune function [2–5]. The resulting clinical problems included respiratory infections, Epstein Barr virus and/or cytomegalovirus infections, antibody deficiency, lymphadenopathy and lymphoma

56 susceptibility. More recently, dominant mutations in the  
57 gene encoding PIK3R1, the p85 $\alpha$  regulatory subunit for  
58 PIK3CD, were also found to result in constitutive hyperac-  
59 tivation of that pathway and immunodeficiency [6–8].  
60 Currently, the dominant *PIK3R1* mutations linked to con-  
61 stitutive hyperactivation have been restricted to mutations  
62 of essential donor splice sites in intron 11, resulting in the  
63 exclusion of exon 11 (*p85 $\alpha$  <sup>$\Delta$ 434–475</sup>*) [6, 7, 9]. Prior to these  
64 reports, a homozygous loss-of-function genotype in exon 6  
65 of *PIK3R1* was reported in a female patient characterized  
66 by agammaglobulinemia and absent B cells [10]. In addi-  
67 tion, other dominant *de novo* mutations in exon 14 of  
68 *PIK3R1* have been described among patients with SHORT  
69 syndrome characterized by short stature, hyperextensibility  
70 of joints, delayed bone age, hernias, low body mass index  
71 and a progeroid appearance [11–16]. No results of immune  
72 studies were recorded in the SHORT syndrome patients in  
73 the existing reports.

74 The purpose of this research was to use next generation  
75 sequencing to identify mutations in patients with primary im-  
76 munodeficiency diseases whose pathogenic gene mutations  
77 had not been identified. Four unrelated patients were found  
78 by next generation sequencing to have the same *de novo* het-  
79 erozygous mutation in an essential donor splice site of  
80 *PIK3R1* (NM\_181523.2:c.1425 + 1G > A) found in three  
81 prior reports [6–8]. All four had the Hyper IgM syndrome,  
82 lymphadenopathy and short stature, and one also had a clinical  
83 diagnosis of SHORT syndrome.

## 84 Patient, Materials and Methods

### 85 Patients

86 The patients from four unrelated families were referred to the  
87 Immunology Clinic at Duke University Medical Center  
88 (DUMC). All studies were performed with the approval of  
89 the DUMC Institutional Review Board and with the written  
90 informed consent of the patients' parents.

91 **Patient 1** A 2.5 year old Caucasian male born to  
92 nonconsanguineous healthy parents began having recurrent  
93 otitis, sinusitis and dacryocystitis in early infancy. He also  
94 had problems with intermittent hypoglycemia. On exami-  
95 nation, he was found to be symmetrically small with height  
96 and weight below the 5th percentile. He had enlarged tonsils  
97 and large cervical, axillary and inguinal lymph nodes.  
98 Family history was negative for immunodeficiency. On im-  
99 mune evaluation, he was found to have an IgG of 78, an IgA  
100 of 0, an IgM of 155 and an IgE of <1 and was started on  
101 monthly intravenous immunoglobulin infusions (Table 1).  
102 He is currently alive but with recurrent respiratory infec-  
103 tions and lymphadenopathy.

**Patient 2** A 2.4 year old Caucasian female was born to  
104 nonconsanguineous healthy parents and was healthy until  
105 she began having recurrent otitis and pneumonia at 2.3 years  
106 of age. On examination, she was found to be symmetrically  
107 small with height and weight below the 10th percentile. She  
108 had large tonsils and massive cervical adenopathy and an el-  
109 evated serum IgM of 209 mg/dl with very low levels of all of  
110 the other immunoglobulins (Table 1). She was started on  
111 monthly intravenous immunoglobulin infusions and is cur-  
112 rently alive but with continued lymphadenopathy. 113

**Patient 3** A 6 year old Caucasian male was born to  
114 nonconsanguineous healthy parents and began having recurrent  
115 otitis and dacryocystitis in infancy, followed by pneumonia on  
116 three occasions at age 4 years. On examination he was found to  
117 be symmetrically small, with height and weight below the 5th  
118 percentiles. Patient 3's height was less than the fifth percentile  
119 for all visits; his mean parental height would predict that he  
120 should be at the 25th percentile for height as an adult. He had  
121 large tonsils, enlarged cervical, axillary and inguinal lymph  
122 nodes and splenomegaly. Immune evaluation at age 4 years  
123 revealed that he had undetectable serum IgA, IgG and IgE but  
124 an IgM of 529 mg/dl (Table 1). He is currently alive and receiv-  
125 ing monthly IVIG with splenic peliosis but is otherwise well. 126

**Patient 4** A 5 year old Caucasian female was born to  
127 nonconsanguineous healthy parents and began having recur-  
128 rent pneumonias at age 3 years. She also had numerous bouts  
129 of otitis and had two sets of tympanostomy tubes. She has  
130 congenital dacryostenosis. She had large tonsils and adenoids  
131 and they were removed. Immune evaluation at age 5 years  
132 revealed undetectable serum IgA and IgE, an IgG of 68 mg/  
133 dl and an IgM of 197 mg/dl (Table 1). At age 12, she was  
134 diagnosed independently by the Genetics Division as having  
135 SHORT syndrome after ruling out multiple other genetic  
136 causes of short stature through whole exome sequencing in a  
137 commercial laboratory where the only abnormality detected  
138 was the *PIK3R1* mutation. Patient 4's height was at the 0.01  
139 percentile for all visits and her mean parental height would  
140 predict that she should be at the 25th percentile as an adult.  
141 She has a triangular facial shape with a prominent forehead,  
142 deep set eyes, thin nostrils, a low-hanging columella, a  
143 downturned small mouth, a small chin, a conductive hearing  
144 loss, hyperextensibility of the joints, delayed eruption of the  
145 secondary teeth, lack of subcutaneous fat and a learning dis-  
146 ability. She is currently alive and receiving monthly IVIG. 147

### 148 Immunologic Phenotype Analysis

149 Flow cytometry of peripheral blood leucocytes was performed  
150 with labeled Abs to CD3 $\epsilon$ , CD4, CD8, CD10, CD14, CD16,  
151 CD20, CD22, CD24, CD25, CD27, CD38, CD45, CD45RA,  
152 CD45RO, CD56, CD57, IgD, IgM, CD62L, CD197, CD279, 152



**Q5/Q4 t1.1 Table 1**

t1.2 Patient	1	2	3	4	
t1.3 Gender	Male	Female	Male	Female	
t1.4 Age at Presentation (yr)	1 7/12	2 5/12	2	5 1/2	
t1.5 Growth Percentile	<b>&lt;5th</b>	<b>&lt;10th</b>	<b>5th</b>	<b>&lt;&lt;&lt;5th</b>	
t1.6 Viral Infections	–	–	RSV, H1N1 Parainfluenza	–	
t1.7 Lymphoproliferation	<b>3+</b>	<b>3+</b>	<b>3+</b>	<b>2+</b>	
t1.8 Autoimmunity	–	–	–	–	
t1.9 Allergy	–	–	–	–	
t1.10 Respiratory Infections	Ear, Sinus	Throat, Lung	Eye, Ear, Lung	Eye, Ear, Lung	
t1.11 Other Features	–	–	–	Abnormal facies <sup>a</sup>	
t1.12 Ig Replacement	+	+	+		<b>Normal Ranges</b>
t1.13 IgG <sup>b</sup> mg/dl (Age in yr)	<b>78</b> (1 7/12)	<b>47</b> (2 5/12)	<b>&lt;60</b> (4 1/12)	<b>68</b> (5 ½)	391–1047
t1.14 IgA <sup>b</sup> mg/dl (Age in yr)	<b>6</b> (1 7/12)	<b>8</b> (2 5/12)	<b>&lt;16</b> (4 1/12)	<b>0</b> (5 ½)	15–95
t1.15 IgM <sup>b</sup> mg/dl (Age in yr)	<b>155</b> (1 7/12)	<b>209</b> (2 5/12)	<b>529</b> (4 1/12)	<b>197</b> (5 ½)	49–202
t1.16 IgE <sup>b</sup> I.U./ml (Age in yr)	<b>&lt;10</b> (1 7/12)	<b>8</b> (2 5/12)	<b>&lt;10</b> (4 1/12)	<b>1</b> (5 ½)	0–150
t1.17 Lymphocyte Studies (Yr)	13 1/4	10 3/4	10 1/3	13 1/12	
t1.18 Absolute Lymphocyte Ct	1280	1476	<b>770</b>	1254	1000–4800
t1.19 CD3+ T cells/μl	891	1231	708	1157	631–4142
t1.20 CD3+ CD4 + T cells/μl	<b>321</b>	344	<b>324</b>	658	340–2746
t1.21 CD3+ CD8 + T cells/μl	392	773	181	433	154–1776
t1.22 Naïve <sup>c</sup> CD4 + T cells (%)	<b>14.2</b>	<b>31.1</b>	<b>11.1</b>	<b>14.2</b>	<sup>i</sup> 35.6–72.5
t1.23 Naïve <sup>d</sup> CD8 + T cells (%)	<b>14.2</b>	<b>5.4</b>	41.2	39.4	<sup>i</sup> 36.1–74.2
t1.24 <sup>e</sup> CD3 + CD8 + PD-1+/μl	56	<b>325</b>	20	22	12–141
t1.25 CD19+ B cells/μl	182	<b>38</b>	<b>16</b>	<b>29</b>	51–821
t1.26 <sup>f</sup> Transitional B cells (%)	<b>52.7 % (102 abs)</b>	4.02 % ( <b>0 abs</b> )	<b>48.8 % (5 abs)</b>	14.2 % ( <b>4 abs</b> )	<sup>j</sup> 2.9%–23.8% (12–35 abs)
t1.27 <sup>g</sup> Memory B cells (%)	3.3 % (6 abs)	11.9 % ( <b>1 abs</b> )	4.4 % ( <b>0 abs</b> )	<b>1.2 % (0 abs)</b>	1.4–27.6 % (4–168abs)
t1.28 <sup>h</sup> Switched Memory B (%)	8.5 % (16 abs)	11.9 % ( <b>1 abs</b> )	3.7 % ( <b>0 abs</b> )	<b>2.4 % (1 abs)</b>	3.0–32.9 % (13–105abs)
t1.29 CD3-CD56+ NK cells/μl	175	155	16	36	12–864
t1.30 PHA Stimulated Cells (cpm)	<b>64,853</b> (1483)	144,073 (2812)	<b>115,456</b> (981)	137,276 (1914)	117,512–235,792
t1.31 Anti-CD3 Stimulated (cpm)	<b>53,366</b> (1483)	133,166 (2812)	<b>33,415</b> (981)	<b>27,782</b> (1528)	59,808–186,006
t1.32 Candida Stimulated (cpm)	<b>1415</b> (888)	41,566 (1450)	<b>6149</b> (4076)	<b>495</b> (545)	6460–60,028

Abnormal values are in bold font

<sup>a</sup> Triangular shape, deep-set eyes, thin nostrils, low-hanging columella, downturned small mouth, small chin and prominent forehead

<sup>b</sup> Normal ranges for immunoglobulins are for 2 year old Caucasians

<sup>c</sup> Naïve CD4+ T cells are CCR7 + CD45RA+/CD4+/CD3+

<sup>d</sup> Naïve CD8+ T cells are CCR7 + CD45RA+/CD8+/CD3+

<sup>e</sup> Senescent CD8+ T cells are CD57 + (PD-1)+/CD8+/CD3+

<sup>f</sup> Transitional B cells are CD24++/CD38++/IgD + CD27–/CD19+

<sup>g</sup> Unswitched memory B cells are IgD + CD27+/CD19+

<sup>h</sup> Class switched memory B cells are CD27 + IgD–/CD19+

<sup>i</sup> Reference range provided by Dr. Alan Kirk of Department of Surgery, Duke University Medical Center

<sup>j</sup> Reference range from R. van Gent et al.: Clin. Immunol. 133: 95–107, 2009

153 TCRαβ, and TCRγδ. Samples were collected on a BD  
 154 FACSCanto II cytometer and data analyzed using  
 155 FACSDiva software. Lymphocyte proliferation was assessed  
 156 by measuring [<sup>3</sup>H]thymidine incorporation into mononuclear  
 157 cells following culture with optimal concentrations of the indicated  
 158 stimuli as previously described [17].


**Next Generation Sequencing, Alignment and Variant Calling** 159  
 160

Next generation sequencing was carried out on the HiSeq2000 161  
 within the Genomic Analysis Facility in the Center for Human 162  
 Genome Variation (Duke University). DNA samples from 163

164	Patients 1 and 2 were analyzed using whole exome sequenc-	pH 7.4). To determine cell expansion, $2 \times 10^4$ of EBV-	215
165	ing and from Patient 3 by whole genome sequencing. For the	transformed B-cell lines were plated in a 96-well plate in	216
166	exome sequenced samples, sequencing libraries were pre-	0.2 mL media. Cell numbers were counted after cultured for	217
167	pared from primary DNA extracted from leukocytes of pa-	24, 48 and 72 h.	218
168	tients using either the KAPA Biosystem's (patient 1) or		
169	Illumina TruSeq (patient 2) library preparation kit follow-	<b>Immunoblot Analysis</b>	219
170	ing the manufacturer's protocol. For patient 1 and his par-		
171	ents, the Nimblegen SeqCap EZ V3.0 Enrichment kit (Roche	EBV-transformed B cells were rested in PBS for 30' at 37 °C	220
172	NimbleGen, Madison, WI) was used to selectively amplify	followed by stimulation with an anti-CD40 antibody for 10 or	221
173	the coding regions of the genome according to the manufac-	30 min or with human insulin (Sigma-Aldrich) for the indicat-	222
174	turer's protocol. For patient 2, the 65-Mb Illumina TruSeq	ed times and subsequently lysed in 1 % Nonidet P-40 lysis	223
175	Exome Enrichment Kit (Illumina, San Diego, CA) was used	buffer containing freshly added protease and phosphatase in-	224
176	to selectively amplify the coding regions of the genome ac-	hibitor cocktails. Cell lysates were subjected to immunoblot-	225
177	ording to the manufacturer's protocol. Patient 3 was whole-	ting analysis using anti-p85 $\alpha$ , phosphor-S6, S6 protein,	226
178	genome sequenced. Whole exome sequencing was performed	phosphor-Akt serine 473, and $\beta$ -actin antibodies according	227
179	initially at a commercial CLIA-certified laboratory for patient	to published protocols [19].	228
180	4, and the mutation was confirmed by the authors.		
181	Alignment of the sequenced DNA fragments to the Human	<b>RT-PCR and DNA Sequencing</b>	229
182	Reference Genome (NCBI Build 37) was performed using the		
183	Burrows–Wheeler Alignment Tool (BWA) (version 0.5.10).	Total RNAs from PBMCs lysed in the Trizol reagent were	230
184	The reference sequence used is identical to the 1000 Genomes	isolated according to the manufacturer's protocol. The first	231
185	Phase II reference and it consists of chromosomes 1–22, X, Y,	strand cDNA was made using the iScript Select cDNA	232
186	MT, unplaced and unlocalized contigs, the human herpesvirus	Synthesis Kit (Bio-Rad). PIK3R1 cDNA was amplified using	233
187	4 type 1 (NC_007605), and decoy sequences (hs37d5) derived	primers huPIK3R1F 5'-TGGGAAATATGGCTTCTCTGA-3	234
188	from HuRef, Human Bac and Fosmid clones and NA12878.	and huPIK3R1R 5'-TCTTTCTCATTGCCTTCACG-3'	235
189	After alignments were produced for each individual sepa-	aligning to exon 10 and 12, respectively. PCR products were	236
190	rately using BWA, variant and genotype calling was per-	separated and visualized by agarose gel electrophoresis and	237
191	formed using the Genome Analysis Tool Kit (GATK, version	further sequenced to identify aberrant splicing.	238
192	1.6–11-g3b2fab9). SnpEff (version 3.3) was used to annotate		
193	the variants according to Ensembl (version 73) and consensus	<b>Glucose Uptake</b>	239
194	coding sequencing (CCDS release 14) and limited analyses to		
195	protein-coding or essential splice site (2 base pairs flank-	To measure glucose uptake, cells were washed and resuspend-	240
196	ing an exon) mutations.	ed in Krebs-Ringer-HEPES (at pH 7.4, 136 mM NaCl, 4.7 ml	241
197	The sequence data for patients 1–3 were analyzed using	KCl, 1.25 mM CaCl <sub>2</sub> , 1.25 mM MgSO <sub>4</sub> , and 10 mM HEPES).	242
198	established protocols that identify qualifying variants forming	2-Deoxy-D-H <sup>3</sup> glucose (2 $\mu$ Ci/reaction) was added, and the	243
199	genotypes not observed in an in-house control database of	cells were incubated for 10 min at 37 °C. The reactions were	244
200	2357 samples or 60,706 sequenced samples made available	quenched by the addition of ice-cold 200 $\mu$ M phloretin	245
201	by the Exome Aggregation Consortium (ExAC), Cambridge,	(Calbiochem, Gibbstown, NJ) followed by centrifugation	246
202	MA (URL: <a href="http://exac.broadinstitute.org">http://exac.broadinstitute.org</a> (accessed Jan.,	through an oil layer (1:1 Dow Corning 550 Silicon fluid from	247
203	2015, release 0.3). The <i>PIK3R1</i> essential splice site <i>de novo</i>	Motion Industries, Birmingham, AL; and dinonyl phthalate	248
204	mutations were independently validated with Sanger	from Sigma-Aldrich). The cell pellets were washed and solu-	249
205	sequencing for each trio.	bilized in 1 M NaOH, and radioactivity was measured using a	250
206		scintillation counter.	251
207	<b>Establishment of EBV Cell Lines and Cell Growth</b>		
208	Cyclosporine A, B95.8 culture EBV supernatant, and media	<b>Results</b>	252
209	(RPMI 1640 + PenStrep + L-glu + Hepes + 20%FBS) were		
210	added to 10 million peripheral blood leucocytes [18]. EBV-	<b>Patients</b> All four patients have had recurrent upper and/or	253
211	immortalized B-cell lines from patients and controls were cul-	lower respiratory infections from early childhood, but only	254
212	tured in RPMI 1640 medium (Sigma-Aldrich) supplemented	one had a history of viral infections and only with viral agents	255
213	with 20 % FBS (HyClone), 100 U/mL penicillin G, 100 U/mL	that cause upper respiratory illnesses (Table 1). They have not	256
214	streptomycin, 292 $\mu$ g/mL of L-glutamine and 10 mM HEPES	had recurrent fungal infections nor have they had opportunist-	257
	(N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid;	tic infections. Three have had recurrent bacterial	258

259 dacryocystitis, and one has had intermittent hypoglycemia.  
 260 All four are symmetrically small: three had heights and  
 261 weights below the 5th percentile and patient 2's height was  
 262 below the 10th percentile. Three of the patients have had pe-  
 263 ripheral lymphadenopathy and the fourth had very large tonsils  
 264 and adenoids that were removed. The lymph nodes are  
 265 generally very large and are often soft and movable. Three  
 266 patients have had lymph nodes excised and no lymph node  
 267 has been found to be malignant. Lymph node histology in the  
 268 three for whom it is available revealed hyperplastic lymphoid  
 269 follicles with attenuated mantle zones (Fig. 1). Only one has  
 270 splenomegaly and that is complicated by peliosis. None of the  
 271 patients has autoimmunity or allergy. Only one patient is lym-  
 272 phopenic. Two of them have low numbers of B cells, and all

273 have very low percentages of memory B cells and class- 273  
 274 switched memory B cells. One of them has an elevated per- 274  
 275 centage and number of transitional B cells. They all have 275  
 276 normal numbers of natural killer cells, although one has only 276  
 277 16 such cells. Their T cell counts are all within the normal 277  
 278 range, but two of them have inverted CD4:CD8 ratios. All of 278  
 279 them have low percentages of naïve CD4+ T cells, and two 279  
 280 have low percentages of naïve CD8+ T cells. One of the pa- 280  
 281 tients has elevated CD8 senescent T cells (Table 1). T cell 281  
 282 functional studies in the four patients suggest some T cell 282  
 283 impairment in that two had low responses to PHA, three had 283  
 284 low responses to immobilized anti-CD3 and three had no re- 284  
 285 sponse to candida (Table 1). 285




**Fig. 1** Histopathology of cervical lymph node biopsy of Patient 1. **a** A low magnification demonstrates reactive follicular hyperplasia with expanded germinal centers and attenuated mantle zones, as indicated by the arrows. Note the polarity of well-defined germinal centers. H&E stain,  $\times 40$ . **b** Immunohistochemical stain for CD20 highlights well defined germinal centers with diminished mantle zones or absence of mantle zones, as indicated by the arrows. Note an apparently naked germinal center without mantle zone in upper left, as indicated by the arrowhead, and increased B-cells in the interfollicular area. Anti-CD20 stain,  $\times 40$

**Mutation** Whole exome sequencing of patient 1 and his parents identified a putative *de novo* mutation in an essential donor splice site of *PIK3R1* (NM\_181523.2:c.1425 + 1G > A) (Supplementary Fig. 1). Among our set of 50 patients who have undergone next generation sequencing for immunodeficiency of unknown molecular type, we identified three other patients, patients 2, 3 and 4, with the exact same *de novo* *PIK3R1* essential donor splice site mutation (NM\_181523.2:c.1425 + 1G > A) (Supplementary Fig. 1). Variation at this essential splice site was not observed among 2357 samples sequenced for various projects at the Institute for Genomic Medicine, Columbia University (formerly the Center for Human Genome Variation, Duke University); nor was it observed among 59,413 samples among the ExAC consortia dataset that had at least 10-fold coverage at this precise site. For patients 1–3, the parents were also Sanger sequenced for the mutation and it was confirmed that the mutation arose *de novo* in those patients. For patient 4, who was clinically whole exome sequenced initially by a commercial CLIA-certified laboratory, the variant was Sanger confirmed in the child and confirmed to be absent in mother, but her father was unavailable (Supplementary Fig. 1). It is likely these *de novo* mutations arose in the germline cells; however, we have not ruled out the possibility that the mutations may be post-zygotic—limited to the hematopoietic system. In our patient population, we have not yet observed instances of the other two described *PIK3R1* essential splice variants at the same site: NM\_181523.2:c.1425 + 1G > C or NM\_181523.2:c.1425 + 1G > T [6, 7].

**Detection of Aberrant Splicing of Mutant mRNA**

To determine if the mutation results in skipping of exon 11, we amplified *PIK3R1* cDNA from total RNAs from patient 1 and his parents PBMCs using primers corresponding to exon 10 and 12 of the gene (transcript NM\_181523.2). Both parents generated a single normal 399 base-pair (bp) band. The patient sample, however, produced both a 399 bp band and a unique 273 bp band (Fig. 2). Further sequencing of the PCR products



**Fig. 2** a–c Aberrant splicing in *PIK3R1* mRNA in patient 1. **a** Detection of mutant form of *PIK3R1* mRNA by RT-PCR. **b** Sequence of normal and mutant allele cDNA. **c** Chromatographs showing aberrant exon 10 to 12 splicing in patient 1’s mutant *PIK3R1* allele. **d** Detection of p85α protein in lysates from EBV-immortalized B cells from patient 1 and his parents. In the patient sample, the intensity of the band corresponding to p85α is decreased and there is a weak band (blue arrow) that could be vaguely see right beneath the p85α band, which is likely the exon 11 deletion mutant of p85α. The band indicated with a red arrow is only seen in patient,

which is likely a degraded product of the mutant p85α. The identities of the other two bands indicated by \*, whether they are non-specific proteins detected by the anti-p85α antibody or degradation products of p85α, are unknown at present. **e** Assessment of S6 and Akt phosphorylation in EBV-B cells following anti-CD40 stimulation. EBV-B cells from patient 1 and her parents were rested in PBS at 37°C for 30’ and then stimulated with an anti-CD40 antibody (10 μg/ml) for 10 or 30 min. Cell lysates were subjected to immunoblotting analysis with the indicated antibodies

324 revealed that the 273 bp band was produced by direct exon 10  
325 to 12 splicing.

### 326 **Effects of $p85\alpha^{\Delta 434-475}$ Mutation on Signaling**

327 To examine how  $p85\alpha^{\Delta 434-475}$  mutation might affect  $p85\alpha$ ,  
328 we generated EBV-immortalized B cell lines from patient 1  
329 and his parents. As shown in Fig. 2d, patient EBV-B cells  
330 expressed lower level of  $p85\alpha$  compared with parent control.  
331 In addition, a small protein band of about 30 kDa was only  
332 observed in the patient's B cells but not parents. Thus, the  
333 mutant protein might be less stable than the WT protein. An  
334 important downstream event of PIK3 signaling is the activa-  
335 tion of mTOR. Activated mTOR phosphorylates and activates  
336 S6 K1, leading to subsequent phosphorylation of the ribosomal  
337 protein S6. After resting in PBS at 37 °C for 30 min, EVB-B  
338 cells from both parents of patient 1 did not contain obvious S6  
339 phosphorylation (mTOR complex 1 dependent event), Akt  
340 phosphorylation at serine 473 (mTOR complex 2 dependent  
341 event), and Akt phosphorylation at threonine 308  
342 (PI3K/PDK1 dependent event). However, a low level but no-  
343 ticeable phosphorylation of these molecules was observed in  
344 patient 1's EBV-B cells, suggesting constitutive activation of  
345 PI3K-mTOR signaling (Fig. 2e). Following anti-CD40 stim-  
346 ulation, patient 1's EBV-B cells displayed a higher level of S6  
347 and Akt phosphorylation than controls, suggesting elevated  
348 PI3K/PDK1 and mTOR signaling downstream of CD40.  
349 The same was true when they were stimulated with insulin  
350 for both patient 1 (top panel, Fig. 3a) and patient 4 (bottom  
351 panel, Fig. 3a). Moreover, patient EBV-B cells had increased  
352 uptake of  $^3\text{H}$ -2-deoxy-glucose following 1 h of stimulation  
353 with insulin, and were more responsive to insulin than the  
354 parent/control EBV-B cells (Fig. 3b). Altogether, these obser-  
355 vations suggest that amino acids 435–474 in PIK3R1 are im-  
356 portant for its stability and also restrain PI3K activity. Its de-  
357 letion may lead to constitutive activation of PI3K/Akt/mTOR  
358 signaling, which is consistent with recent reports of patients  
359 with similar mutations [6–8].

### 360 **Discussion**


361 Including the 12 patients in the three earlier reports [6–8] all  
362 sixteen known cases of *PIK3R1* immunodeficiency have  
363 had mutations at this essential donor splice site  
364 (NM\_181523.2:c.1425 + 1G > A/C/T). Despite this appar-  
365 ent limited allelic heterogeneity for *PIK3R1* immunodefi-  
366 ciency, the findings in the four patients presented here con-  
367 tinue to highlight the heterogeneity of the clinical and im-  
368 munologic abnormalities of the patients who have  
369  $p85\alpha^{\Delta 434-475}$  *PIK3R1* essential splice mutations [6–8].  
370 The clinical features of the four patients reported here are  
371 uniform and similar to some but not all of those described

372 by Deau et al. [6] and Lucas et al. [7]. This could possibly  
373 be due to the younger ages of our patients when compared  
374 with those of Lucas et al. [7], but the ages at presentation of  
375 our patients were similar to three of the four patients de-  
376 scribed by Deau et al. [6] and all of the patients reported by  
377 Lougaris et al. [8] All four of our patients had short stature;  
378 poor growth was noted in the patients reported by Lucas  
379 et al. [7] and Lougaris et al. [8] but in only one of those  
380 reported by Deau et al. [6] Lymphadenopathy and/or ton-  
381 sillar hypertrophy were noted in all four of our patients and  
382 in the four reported by Lucas et al. [7], and in three of the  
383 four reported by Lougaris et al. [8] but in only one of those  
384 reported by Deau et al. [6] None of our patients has had a  
385 diagnosed malignancy as yet and none has had problems  
386 with CMV or EBV. Our patients were each diagnosed clin-  
387 ically as having the Hyper IgM syndrome, but none of them  
388 had any of the known gene mutations reported in the Hyper  
389 IgM syndrome. Patient 1 had been found to have a common  
390 CD40L polymorphism, but his CD4+ T cells displayed  
391 functional CD40L after activation [20]. Patient 4, with her  
392 diagnosis of SHORT syndrome, appears to be a unique case  
393 compared to other patients with this same mutation. While  
394 there may exist another mutation in Patient 4 that makes the  
395 presentation unique, it was not found in chromosomal mi-  
396 croarray studies, fragile X studies, whole exome sequence  
397 analysis in the commercial lab setting, subsequent Sanger  
398 sequencing of *PIK3R1* exon 14, or in deletion testing of the  
399 mitochondrial genome.

400 *PIK3R1* is an established pleiotropic gene. In addition to  
401 immunodeficiency, SHORT syndrome has been reported to be  
402 caused by dominant *de novo* mutations in exon 14 of *PIK3R1*  
403 [11–16]. Immunologic studies were not reported on in those  
404 earlier SHORT syndrome publications. Thus, our Patient 4  
405 with SHORT syndrome having the same mutation at this es-  
406 sential donor splice site (NM\_181523.2:c.1425 + 1G > A/C/  
407 T) in intron 11 as well as the same immunologic findings as  
408 the other patients with *PIK3R1* immunodeficiency, provides  
409 an important link within *PIK3R1* mediated disease. The com-  
410 mercial lab reported Patient 4 as negative for additional  
411 *PIK3R1*, or other SHORT syndrome disease gene, variants.  
412 We also subsequently Sanger sequenced *PIK3R1* exon 14 in  
413 our research lab setting and confirmed that both Patient 4 and  
414 her mother were not carriers of variants in exon 14. We per-  
415 formed glucose uptake studies because some but not all pa-  
416 tients with SHORT syndrome have been insulin resistant.  
417 However, in keeping with the fact than none of the four pa-  
418 tients reported here were clinically insulin resistant, the glu-  
419 cose uptake studies were normal (Fig. 3).

420 Our data suggest that  $p85\alpha^{\Delta 434-475}$  *PIK3R1* mutations lead  
421 to elevated PI3K activity. The immunodeficiency observed in  
422 our patients is similar to those with gain-of-function mutations  
423 in *PIK3CD*, suggesting that this *PIK3R1* mutation might  
424 cause elevated PIK3CD activity [21]. At present, it is still

**Fig. 3 a** Assessment of S6 phosphorylation in EBV-B cells following anti-CD40 stimulation. EBV-B cells from patients 1 and 4 and their mothers were rested in PBS at 37 °C for 30' and then either unstimulated or stimulated with insulin (1 μM) for the indicated minutes. Cell lysates were subjected to immunoblotting analysis with the indicated antibodies. **b** Assessment of glucose uptake before and after insulin stimulation: EBV-transformed B cells from Patients 1, 2 and 4 their parents (controls) were rested in PBS for 30 min followed by culture in the presence or absence of 1–2 μM insulin for 30–60 min. Glucose uptake was determined



425 unclear how the p85α<sup>Δ434-475</sup> mutation causes increased PI3K  
 426 activity. Normal p85α stabilizes the p110 catalytic subunit but  
 427 inhibits its lipid kinase activity in unstimulated cells. Under  
 428 stimulating conditions, p85α recruits p110 to activated recep-  
 429 tors or adaptor molecules via its SH2 domain, leading to acti-  
 430 vation of its kinase activity [22, 23]. Because the inter-SH2  
 431 domain of p85α is involved in p85α - p110 heterodimerization  
 432 [24], it is possible that the p85α<sup>Δ434-475</sup> mutant inefficiently  
 433 inhibits its cognate partner p110 (p110δ in the hematopoietic  
 434 cells and p110α in muscle and other tissues), leading to elevat-  
 435 ed PI3K signaling. Additionally, p85α<sup>Δ434-475</sup> was under-  
 436 represented compared with WT p85α in patients' samples, indi-  
 437 cating that amino acids absent in the mutant in the inter-SH2  
 438 domain are important for its stability.

439 The concomitant observation of SHORT syndrome in patient  
 440 4 is interesting in that it confirms *PIK3R1* as a hot spot of  
 441 mutation for this syndrome. It also suggests that the  
 442 p85α<sup>Δ434-475</sup> mutation may affect other catalytic PI3K sub-  
 443 units such as PIK3CA that are known to be important for  
 444 insulin receptor signaling [25]. It is important to note that  
 445 some SHORT syndrome patients with mutations in exon 14  
 446 [11–14] and a murine model mimicking one of the *PIK3R1*  
 447 mutations [26] displayed impaired PI3K signaling. Studies  
 448 have demonstrated that chronic overactivation of signal path-  
 449 ways can lead impairment of receptor induced signaling [27].  
 450 It is possible that chronic overactivation of the PI3K pathway  
 451 might trigger a negative feedback mechanism(s) that prevents  
 452 normal insulin receptor signaling in some tissues, although we

453 did not find abnormal signaling in patient 4's blood  
 454 lymphocytes.

455 The immunologic abnormalities in our patients were similar  
 456 in many but not all respects to those previously reported  
 457 [6–8]. Three of our patients have low numbers of B cells, and  
 458 all have very low percentages of memory B cells and class-  
 459 switched memory B cells. IgM was elevated in all four of our  
 460 patients, in three of the four patients reported by Deau et al.  
 461 [6], all of the patients reported by Lougaris et al. [8] but in only  
 462 one of the patients reported by Lucas et al. [7] Most of our  
 463 patients have low percentages of naïve CD4+ and CD8+ T  
 464 cells, but only one had elevated CD8 senescent T cells as  
 465 reported by Lucas et al. [7] T cell functional studies in our  
 466 patients suggest some T cell impairment but this was not  
 467 commented on in the three prior reports [6–8]. T cell dysfunc-  
 468 tion would not be unexpected, since PI3K signaling is re-  
 469 quired for the differentiation of T cells into subsets [1].

470 *PIK3R1* has been identified as a gene intolerant to func-  
 471 tional variation in the general human population. The Residual  
 472 Variation Intolerance Score (RVIS) for *PIK3R1* is -0.76,  
 473 which corresponds to a ranking of 13.5 % most intolerant  
 474 genes, genome wide (<http://igm.cumc.columbia.edu/GenicIntolerance/>).  
 475 The intolerance of a gene to functional  
 476 variation among the human population has been  
 477 demonstrated to be highly predictive of disease-causing genes,  
 478 particularly those causing disease through a dominant model  
 479 [28]. Existing mouse knock-out work has also shown that  
 480 disruptions to *Pik3r1* result in a wide range of B cell

481 phenotypes [29]. Moreover, *Pik3r1* is considered an ‘essential’ gene, with mouse knock-outs resulting in complete perinatal lethality and partial postnatal lethality [29]. We have previously shown that the bioinformatics signature of a disruptive *de novo* mutation in an intolerant gene that is also known to be essential is highly linked to pathogenicity among children with undiagnosed genetic disorders [30].

488 Available mTOR inhibitors as well as non-selective PI3K inhibitors approved for use in other conditions already exist. A range of next generation PI3K inhibitors, which are highly selective for the delta isoform, are under development. These drugs could offer a pharmacological solution for these patients where existing treatment options are very limited and include extreme therapies such as bone marrow transplantation. The availability of highly selective orally available PI3Kdelta inhibitors has the potential to normalize the pathological mechanism in these patients.

498 **Conclusions**

499 Together, these observations suggest that amino acids 435–474 in PIK3R1 are important for its stability and also its ability to restrain PI3K activity. Deletion of exon 11 leads to constitutive activation of PI3K signaling. The clinical and immunologic manifestations of patients with dominant splice site mutations in *PIK3R1* have been variable, with all having very low percentages of memory B cells and class-switched memory B cells and reduced percentages of naïve CD4+ and CD8+ T cells. The clinical manifestations in our patients were uniform, with all having the Hyper IgM syndrome, short stature and lymphadenopathy, except that one also had a clinical diagnosis of SHORT syndrome. This is the first report of immunologic abnormalities and this precise mutation in a patient ascertained for SHORT syndrome. The elucidation of the molecular mechanism in these patients results in the opportunity for new and potentially transformative therapeutic options.

515 **Acknowledgments** This study was supported by the following grants: Baxter Healthcare Grant BT13-21694 to R.H.B. and NIH Grant R01 AI101206 to X-P. Z. Partial funding for this study was provided by UCB Celltech, including salary support for D.B.G. M.A. and D.M. are employees of UCB. The other authors declare no conflict of interest. S.P. is a National Health and Medical Research Council (NHMRC) CJ Martin Fellow. We thank our patients and their parents for participating in this study. We also thank B Krueger, J Bridges, Q Wang, N Ren, S Gewalt, S Kisselev, Y-F Lu, K Cronin, and N Walley for excellent technical support. We would like to acknowledge the following individuals for the contributions of control samples: W. B. Gallentine, E.L. Heinzen, A.M. Husain, K. N. Linney, M. A. Mikati, R. A. Radtke, and S. R. Sinha; J.P. McEvoy, A. Need, J. Silver, and *M. Silver*; D. H. Murdock and The MURDOCK Study Community Registry and Biorepository; G. Cavalleri, N. Delanty, and C. Depondt; J. Burke, C. Hulette, and K. Welsh-Bohmer; J. Milner; J. Hoover-Fong, N. L. Sobreira and D. Valle; E. J. Holtzman; W. L. Lowe; P. Lugar; S. M. Palmer; Z. Farfel, D. Lancet, E. Elon Pras; A. Poduri; M. Hauser; D. Marchuk; D. Koltai Attix, O. Chiba-Falek; E. T. Cirulli, V.

Dixon and J. McEvoy; K. Schmader, S. McDonald, H. K. White, M. Yanamadala, and the Carol Woods and Croasdale Retirement Communities; R. Gbadegesin and M. Winn; D. Daskalakis; Q. Zhao; A. Holden and E. Behr; R. Brown; and S. Kerns and H. Oster. The collection of control samples was funded in part by Bryan ADRC NIA P30 AG028377, the Ellison Medical Foundation New Scholar award AG-NS-0441-08, an award from SAIC-Frederick, Inc. (M11-074), funding from Biogen Idec, NIMH awards RC2MH089915, R01MH097971, R01MH099216, and K01MH098126, the Epi4K Gene Discovery in Epilepsy study (NINDS U01-NS077303) and the Epilepsy Genome/Phenome Project (EPGP - NINDS U01-NS053998), and the Center for HIV/AIDS Vaccine Immunology (“CHAVI”) under a grant from the National Institute of Allergy and Infectious Diseases, National Institutes of Health (U01AIO67854), NIAID (R01AIO79088 and R01AI101206). Finally, the authors would also like to thank the Exome Aggregation Consortium and the groups that provided exome variant data for comparison. A full list of contributing groups can be found at <http://exac.broadinstitute.org/about>.

**References**

1. Okkenhaug K. Signaling by the phosphoinositide 3-kinase family in immune cells. *Annu Rev Immunol.* 2013;31:675–704.

2. Angulo I, Vadas O, Garcon F, Banham-Hall E, Plagnol V, Leahy TR, et al. Phosphoinositide 3-kinase delta gene mutation predisposes to respiratory infection and airway damage. *Science.* 2013;342(6160):866–71.

3. Lucas CL, Kuehn HS, Zhao F, Niemela JE, Deenick EK, Palendira U, et al. Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human immunodeficiency. *Nat Immunol.* 2014;15(1):88–97.

4. Kracker S, Curtis J, Ibrahim MA, Sediva A, Salisbury J, Campr V, et al. Occurrence of B-cell lymphomas in patients with activated phosphoinositide 3-kinase delta syndrome. *J Allergy Clin Immunol.* 2014;134(1):233–6.

5. Crank MC, Grossman JK, Moir S, Pittaluga S, Buckner CM, Kardava L, et al. Mutations in PIK3CD can cause hyper IgM syndrome (HIGM) associated with increased cancer susceptibility. *J Clin Immunol.* 2014;34(3):272–6.

6. Deau MC, Heurtier L, Frange P, Suarez F, Bole-Feysot C, Nitschke P, et al. A human immunodeficiency caused by mutations in the PIK3R1 gene. *J Clin Invest.* 2014;124(9):3923–8.

7. Lucas CL, Zhang Y, Venida A, Wang Y, Hughes J, McElwee J, et al. Heterozygous splice mutation in PIK3R1 causes human immunodeficiency with lymphoproliferation due to dominant activation of PI3K. *J Exp Med.* 2014;211(13):2537–47.

8. Lougaris V, Faletta F, Lanzi G, Vozzi D, Marcuzzi A, Valencic E, et al. Altered germinal center reaction and abnormal B cell peripheral maturation in PI3KR1-mutated patients presenting with HIGM-like phenotype. *Clin Immunol.* 2015;159(1):33–6.

9. Deau MC, Heurtier L, Frange P, Suarez F, Bole-Feysot C, Nitschke P, et al. A human immunodeficiency caused by mutations in the PIK3R1 gene. *J Clin Invest.* 2015;125(4):1764–5.

10. Conley ME, Dobbs AK, Quintana AM, Bosompem A, Wang YD, Coustan-Smith E, et al. Agammaglobulinemia and absent B lineage cells in a patient lacking the p85alpha subunit of PI3K. *J Exp Med.* 2012;209(3):463–70.

11. Dymnt DA, Smith AC, Alcantara D, Schwartzentruber JA, Basel-Vanagaite L, Curry CJ, et al. Mutations in PIK3R1 cause SHORT syndrome. *Am J Hum Genet.* 2013;93(1):158–66.

12. Chudasama KK, Winnay J, Johansson S, Claudi T, Konig R, Haldorsen I, et al. SHORT syndrome with partial lipodystrophy

595 due to impaired phosphatidylinositol 3 kinase signaling. *Am J Hum* 625  
 596 *Genet.* 2013;93(1):150–7. 626

597 13. Thauvin-Robinet C, Auclair M, Duplomb L, Caron-Debarle M, 627  
 598 Avila M, St-Onge J, et al. PIK3R1 mutations cause syndromic 628  
 599 insulin resistance with lipoatrophy. *Am J Hum Genet.* 2013;93(1): 629  
 600 141–9. 630

601 14. Schroeder C, Riess A, Bonin M, Bauer P, Riess O, Dobler- 631  
 602 Neumann M, et al. PIK3R1 mutations in SHORT syndrome. *Clin* 632  
 603 *Genet.* 2014;86(3):292–4. 633

604 15. Barcena C, Quesada V, De Sandre-Giovannoli A, Puente DA, 634  
 605 Fernandez-Toral J, Sigaudy S, et al. Exome sequencing identifies 635  
 606 a novel mutation in PIK3R1 as the cause of SHORT syndrome. 636  
 607 *BMC Med Genet.* 2014;15:51. 637

608 16. Chung BK, Gibson WT. Autosomal dominant PIK3R1 mutations 638  
 609 cause SHORT syndrome. *Clin Genet.* 2014;85(3):228–9. 639

610 17. Buckley RH, Schiff SE, Sampson HA, Schiff RI, Markert ML, 640  
 611 Knutsen AP, et al. Development of immunity in human severe 641  
 612 primary T cell deficiency following haploidentical bone marrow 642  
 613 stem cell transplantation. *J Immunol.* 1986;136:2398–407. 643

614 18. Sugden B, Mark W. Clonal transformation of adult human leuko- 644  
 615 cytes by Epstein-Barr virus. *J Virol.* 1977;23(3):503–8. 645

616 19. Gorentla BK, Wan CK, Zhong XP. Negative regulation of mTOR 646  
 617 activation by diacylglycerol kinases. *Blood.* 2011;117(15):4022– 647  
 618 31. 648

619 20. Seyama K, Nonoyama S, Gangsaas I, Hollenbaugh D, Pabst HF, 649  
 620 Aruffo A, et al. Mutations of the CD40 ligand gene and its effect on 650  
 621 CD40 ligand expression in patients with X-linked hyper IgM syn- 651  
 622 drome. *Blood.* 1998;92(7):2421–34. 652

623 21. Elgizouli M, Lowe DM, Speckmann C, Schubert D, Hulsdunker J, 653  
 624 Eskandarian Z, et al. Activating PI3Kdelta mutations in a cohort of 654  
 625 669 patients with primary immunodeficiency. *Clin Exp Immunol.* 625  
 626 2015;6. 626

627 22. Burke JE, Williams RL. Synergy in activating class I PI3Ks. *Trends* 627  
 628 *Biochem Sci.* 2015;40(2):88–100. 628

629 23. Geering B, Cutillas PR, Vanhaesebroeck B. Regulation of class IA 629  
 630 PI3Ks: is there a role for monomeric PI3K subunits? *Biochem Soc* 630  
 631 *Trans.* 2007;35(Pt 2):199–203. 631

632 24. Chiu YH, Lee JY, Cantley LC. BRD7, a tumor suppressor, interacts 632  
 633 with p85alpha and regulates PI3K activity. *Mol Cell.* 2014;54(1): 633  
 634 193–202. 634

635 25. Cheng Z, Tseng Y, White MF. Insulin signaling meets mitochondria 635  
 636 in metabolism. *Trends Endocrinol Metab.* 2010;21(10):589–98. 636

637 26. Winnay JN, Solheim MH, Dirice E, Sakaguchi M, Noh HL, Kang 637  
 638 HJ, et al. PI3-kinase mutation linked to insulin and growth factor 638  
 639 resistance in vivo. *J Clin Invest.* 2016. doi:10.1172/JC184005. 639

640 27. Yang J, Zhang P, Krishna S, Wang J, Lin X, Huang H, et al. 640  
 641 Unexpected positive control of NFkappaB and miR-155 by 641  
 642 DGKalpha and zeta ensures effector and memory CD8+ T Cell 642  
 643 differentiation. *Oncotarget.* 2016. doi:10.18632/oncotarget.8164. 643

644 28. Petrovski S, Wang Q, Heinzen EL, Allen AS, Goldstein DB. Genic 644  
 645 intolerance to functional variation and the interpretation of personal 645  
 646 genomes. *PLoS Genet.* 2013;9(8):e1003709. 646

647 29. Blake JA, Bult CJ, Eppig JT, Kadin JA, Richardson JE. The mouse 647  
 648 genome database: integration of and access to knowledge about the 648  
 649 laboratory mouse. *Nucleic Acids Res.* 2014;42(Database issue): 649  
 650 D810–7. 650

651 30. Zhu X, Petrovski S, Xie P, Ruzzo EK, Lu YF, McSweeney KM, 651  
 652 et al. Whole-exome sequencing in undiagnosed genetic diseases: 652  
 653 interpreting 119 trios. *Genet Med.* 2015;17:774–81. 653Q6



## AUTHOR QUERIES

### **AUTHOR PLEASE ANSWER ALL QUERIES.**

- Q1. Please check if the affiliations are presented correctly.
- Q2. Please check captured corresponding affiliation if correct.
- Q3. Please check if the section headings are assigned to appropriate levels.
- Q4. Please check table entries, notes and other relevant details for table 1 if captured and presented correctly.
- Q5. Please provide descriptive legend (caption) for Table 1.
- Q6. Please advise if the provided volume and page numbers are correct.

UNCORRECTED PROOF