

## Give Me Some Room to Breathe! Can Targeting SPHK2 Reduce Airway Smooth Muscle Thickening in Asthma?

During the pathobiology of asthma, airway smooth muscle cells (ASMCs) play prominent roles in all aspects of the disease, including airway inflammation, hyperresponsiveness, and remodeling (1). Remodeling includes irreversible airway structural changes, leading to airway wall thickening, decreased airway luminal diameter, and fixed airway obstruction (2). Persistent airflow obstruction, which contributes to asthma severity, is associated with increased airway wall thickness and ASM area (3, 4). Furthermore, excessive airway narrowing is associated with airway hyperresponsiveness (AHR) in asthma (5). ASMC hyperplasia (6), hypertrophy (7), and migration (8) have all been implicated in human asthma as potential mechanisms of increased ASM mass. Studies of the beneficial effects of standard asthma therapies on airway wall thickening have yielded contradictory results (9–12), which may be related to corticosteroid resistance in subsets of patients and to reduction of airway inflammation rather than reversal of established airway remodeling. Thus, identification of pharmacological agents that can halt the progression of or even reverse ASMC accumulation in asthma is a key therapeutic goal.

S1P (sphingosine-1-phosphate) is a potent stimulator of ASMC proliferation, contraction, and airway hyperreactivity in the context of asthma and allergic airways disease (13–15). Sphingosine is a natural substrate of SPHK1 and SPHK2 (sphingosine kinases 1 and 2), which catalyze the intracellular formation of S1P. Transport of S1P outside of the cell renders it cell impermeant, and its signaling effects on cell activity are mediated by binding of S1P to several cell membrane G-protein-coupled receptors (S1P1–5) (16). Much research interest in recent years has focused on the ability of sphingosine analogs to abrogate disease progression through either receptor-mediated or -independent mechanisms. Prophylactic administration of the drug FTY720 (fingolimod), a synthetic sphingosine analog that has been approved for clinical treatment of multiple sclerosis, has been shown to reduce AHR and ASM mass in rodent models of allergic airways disease (17, 18). Thus, the sphingosine pathway represents a potential molecular target for development of new therapies to prevent airway remodeling in asthma.

In this issue of the *Journal*, Blais-Lecours and colleagues (pp. 35–42) define the mechanism of action of a synthetic sphingosine analog and SPHK2 substrate, AAL-R ([*(R)*-2-amino-4-(4-heptyloxyphenyl)-2-methylbutanol]), to inhibit ASMC proliferation (19). Previous work by this group demonstrated that intratracheal administration of AAL-R halted airway immune cell accumulation and AHR in a mouse model of allergic airway disease with established house dust mite allergen-induced airway inflammation (20). Building on those studies, the authors further showed that AAL-R reversed AHR and *in vivo* ASM thickening in a model of chronic house dust mite allergen challenge with established airway remodeling. Interestingly, *in vitro* studies of

human ASMCs in complete media showed that at low concentrations (1  $\mu$ M), AAL-R did not induce apoptosis but diminished cell proliferation and metabolic activity (21). The current well-designed study expands on these findings by delineating the mechanism whereby AAL-R inhibits ASMC growth. The authors show that AAL-R perturbed cellular metabolism such that oxygen use by the cells was reduced and metabolic activity was shifted to glycolysis. These metabolism-altering effects of AAL-R were associated with high levels of SPHK2 in Ki-67<sup>+</sup> (proliferating) ASMCs and in those cells after mitogenic stimuli. Other pulmonary stromal cells that express relatively lower levels of SPHK2, such as airway epithelial cells and arterial endothelial cells, were not as susceptible to the oxidoreductive effects of AAL-R as the ASMCs were. Finally, the authors show that SPHK2-induced phosphorylation was required for the AAL-R-mediated reduction of cellular metabolism in proliferating ASMCs, but S1P1 and S1P3 receptor signaling was not required for this effect.

Currently, the only approved therapy that effectively reduces ASM mass in asthma is bronchial thermoplasty, which is indicated for patients with severe asthma who have persistent symptoms that are refractive to other therapies (22). For the majority of patients with asthma, innovative pharmacological options are still needed to address airway wall thickening and hyperreactivity. Although Blais-Lecours and colleagues used normal ASMCs and stromal cell lines for most of their experiments, their results provide preliminary evidence that suggests that ASMCs derived from patients with asthma are susceptible to the metabolic perturbations induced by AAL-R. Much more work needs to be done to test the efficacy of sphingosine analogs and other therapies that modulate SPHK2 expression or activity to inhibit or reverse ASMC accumulation in asthmatic airways. Nevertheless, this work and other recent reports of pharmacology-mediated reductions of ASM mass in human asthma (23) reveal molecular pathways that may hold the key to alleviating airway remodeling in asthma. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Jennifer L. Ingram, Ph.D.  
Department of Medicine  
Department of Surgery  
and  
Department of Pathology  
Duke University Medical Center  
Durham, North Carolina

ORCID ID: 0000-0002-5269-8864 (J.L.I.).

Ⓞ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0 (<http://creativecommons.org/licenses/by-nc-nd/4.0/>). For commercial usage and reprints, please contact Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Originally Published in Press as DOI: 10.1165/rcmb.2019-0240ED on July 12, 2019

## References

- Ojaku CA, Yoo EJ, Panettieri RA Jr. Transforming growth factor  $\beta$ 1 function in airway remodeling and hyperresponsiveness: the missing link? *Am J Respir Cell Mol Biol* 2017;56:432–442.
- Pascoe CD, Seow CY, Hackett TL, Paré PD, Donovan GM. Heterogeneity of airway wall dimensions in humans: a critical determinant of lung function in asthmatics and nonasthmatics. *Am J Physiol Lung Cell Mol Physiol* 2017;312:L425–L431.
- Kaminska M, Foley S, Maghni K, Storness-Bliss C, Coxson H, Ghezzo H, *et al.* Airway remodeling in subjects with severe asthma with or without chronic persistent airflow obstruction. *J Allergy Clin Immunol* 2009;124:45–51, e1–4.
- Ferreira DS, Carvalho-Pinto RM, Gregório MG, Annoni R, Teles AM, Buttignol M, *et al.* Airway pathology in severe asthma is related to airflow obstruction but not symptom control. *Allergy* 2018;73:635–643.
- Paré PD, McParland BE, Seow CY. Structural basis for exaggerated airway narrowing. *Can J Physiol Pharmacol* 2007;85:653–658.
- Woodruff PG, Dolganov GM, Ferrando RE, Donnelly S, Hays SR, Solberg OD, *et al.* Hyperplasia of smooth muscle in mild to moderate asthma without changes in cell size or gene expression. *Am J Respir Crit Care Med* 2004;169:1001–1006.
- Ebina M, Takahashi T, Chiba T, Motomiya M. Cellular hypertrophy and hyperplasia of airway smooth muscles underlying bronchial asthma: a 3-D morphometric study. *Am Rev Respir Dis* 1993;148:720–726.
- Salter B, Pray C, Radford K, Martin JG, Nair P. Regulation of human airway smooth muscle cell migration and relevance to asthma. *Respir Res* 2017;18:156.
- Kistemaker LE, Bos IS, Menzen MH, Maarsingh H, Meurs H, Gosens R. Combination therapy of tiotropium and ciclesonide attenuates airway inflammation and remodeling in a guinea pig model of chronic asthma. *Respir Res* 2016;17:13.
- Niimi A, Matsumoto H, Amitani R, Nakano Y, Sakai H, Takemura M, *et al.* Effect of short-term treatment with inhaled corticosteroid on airway wall thickening in asthma. *Am J Med* 2004;116:725–731.
- Debelleix S, Siao-Him Fa V, Begueret H, Berger P, Kamaev A, Ousova O, *et al.* Montelukast reverses airway remodeling in actively sensitized young mice. *Pediatr Pulmonol* 2018;53:701–709.
- Gao JM, Cai F, Peng M, Ma Y, Wang B. Montelukast improves air trapping, not airway remodeling, in patients with moderate-to-severe asthma: a pilot study. *Chin Med J (Engl)* 2013;126:2229–2234.
- Roviezzo F, Di Lorenzo A, Bucci M, Brancaleone V, Vellecco V, De Nardo M, *et al.* Sphingosine-1-phosphate/sphingosine kinase pathway is involved in mouse airway hyperresponsiveness. *Am J Respir Cell Mol Biol* 2007;36:757–762.
- Miller M, Tam AB, Mueller JL, Rosenthal P, Beppu A, Gordillo R, *et al.* Cutting edge: targeting epithelial ORMDL3 increases, rather than reduces, airway responsiveness and is associated with increased sphingosine-1-phosphate. *J Immunol* 2017;198:3017–3022.
- Liu L, Zhai C, Pan Y, Zhu Y, Shi W, Wang J, *et al.* Sphingosine-1-phosphate induces airway smooth muscle cell proliferation, migration, and contraction by modulating Hippo signaling effector YAP. *Am J Physiol Lung Cell Mol Physiol* 2018;315:L609–L621.
- Natarajan V, Dudek SM, Jacobson JR, Moreno-Vinasco L, Huang LS, Abassi T, *et al.* Sphingosine-1-phosphate, FTY720, and sphingosine-1-phosphate receptors in the pathobiology of acute lung injury. *Am J Respir Cell Mol Biol* 2013;49:6–17.
- Karmouty-Quintana H, Siddiqui S, Hassan M, Tsuchiya K, Risse PA, Xicota-Vila L, *et al.* Treatment with a sphingosine-1-phosphate analog inhibits airway remodeling following repeated allergen exposure. *Am J Physiol Lung Cell Mol Physiol* 2012;302:L736–L745.
- Oyeniran C, Sturgill JL, Hait NC, Huang WC, Avni D, Maceyka M, *et al.* Aberrant ORM (yeast)-like protein isoform 3 (ORMDL3) expression dysregulates ceramide homeostasis in cells and ceramide exacerbates allergic asthma in mice. *J Allergy Clin Immunol* 2015;136:1035–46.e6.
- Blais-Lecours P, Laouafa S, Arias-Reyes C, Santos WL, Joseph V, Burgess JK, *et al.* Metabolic adaptation of airway smooth muscle cells to an SPHK2 substrate precedes cytotostasis. *Am J Respir Cell Mol Biol* 2020;62:35–42.
- Gendron D, Lemay A-M, Tremblay C, Lai LJA, Langlois A, Bernatchez É, *et al.* Treatment with a sphingosine analog after the inception of house dust mite-induced airway inflammation alleviates key features of experimental asthma. *Respir Res* 2015;16:7.
- Gendron DR, Lecours PB, Lemay A-M, Beaulieu M-J, Huppé C-A, Lee-Gosselin A, *et al.* A phosphorylatable sphingosine analog induces airway smooth muscle cytotostasis and reverses airway hyperresponsiveness in experimental asthma. *Front Pharmacol* 2017;8:78.
- Chakir J, Haj-Salem I, Gras D, Joubert P, Beaudoin É-L, Biardel S, *et al.* Effects of bronchial thermoplasty on airway smooth muscle and collagen deposition in asthma. *Ann Am Thorac Soc* 2015;12:1612–1618.
- Saunders R, Kaul H, Berair R, Gonem S, Sinqapuri A, Sutcliffe AJ, *et al.* DP2 antagonism reduces airway smooth muscle mass in asthma by decreasing eosinophilia and myofibroblast recruitment. *Sci Transl Med* 2019;11:eaao6451.