

QUALITY AND PATIENT SAFETY

A systematic review and consensus definitions for standardised end-points in perioperative medicine: pulmonary complications

T. E. F. Abbott^{1,2}, A. J. Fowler¹, P. Pelosi³, M. Gama de Abreu⁴, A. M. Møller⁵, J. Canet⁶, B. Creagh-Brown^{7,8}, M. Mythen², T. Gin⁹, M. M. Lalu¹⁰, E. Futier¹¹, M. P. Grocott^{12,13}, M. J. Schultz^{14,15}, R. M. Pearse^{1,*}, and the StEP-COMPAC Group[#]

¹Queen Mary University of London, London, UK, ²Department of Anaesthesia and Critical Care, University College London Hospital, London, UK, ³Department of Surgical Sciences and Integrated Diagnostics, San Martino Policlinico Hospital, IRCCS for Oncology, University of Genoa, Genoa, Italy, ⁴Department of Anesthesiology and Intensive Care Therapy, University Hospital Carl Gustav Carus, Technische Universität Dresden, Dresden, Germany, ⁵Department of Anaesthesiology and Intensive Care Medicine, Herlev University Hospital, Herlev, Denmark, ⁶Department of Anaesthesiology, Hospital Universitari Germans Trias i Pujol, Barcelona, Spain, ⁷Intensive Care Unit, Royal Surrey County Hospital, Guildford, UK, ⁸Surrey Perioperative Anaesthesia Critical Care Collaborative Research Group (SPACeR), Faculty of Health and Medical Sciences, University of Surrey, Guildford, UK, ⁹Department of Anaesthesia and Intensive Care, The Chinese University of Hong Kong, Hong Kong, China, ¹⁰Department of Anesthesiology and Pain Medicine, Ottawa Hospital, University of Ottawa, Clinical Epidemiology and Regenerative Medicine Programs, Ottawa Hospital Research Institute, Ottawa, ON, Canada, ¹¹Department of Perioperative Medicine, Anaesthesiology and Critical Care, Estaing Hospital, University Hospital of Clermont-Ferrand, Clermont-Ferrand, France, ¹²Integrative Physiology and Critical Illness Group, Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, UK, ¹³Critical Care Research Group, Southampton NIHR Biomedical Research Centre, University Hospital Southampton, Southampton, UK, ¹⁴Department of Intensive Care, Academic Medical Center, Amsterdam, The Netherlands and ¹⁵Mahidol Oxford Tropical Medicine Research Unit, Mahidol University, Bangkok, Thailand

*Corresponding author. E-mail: r.pearse@qmul.ac.uk

[#] Members listed in the [Supplementary Material](#).



This article is accompanied by an editorial in a forthcoming issue: Shades of grey: embracing uncertainty in the exercise room by R J T Wilson, *Br J Anesth* 2018;120, doi: [10.1016/j.bja.2018.03.016](https://doi.org/10.1016/j.bja.2018.03.016).

Editorial decision: February 12, 2018; Accepted: February 12, 2018

© 2018 British Journal of Anaesthesia. Published by Elsevier Ltd. All rights reserved.
For Permissions, please email: permissions@elsevier.com

Abstract

Background: There is a need for robust, clearly defined, patient-relevant outcome measures for use in randomised trials in perioperative medicine. Our objective was to establish standard outcome measures for postoperative pulmonary complications research.

Methods: A systematic literature search was conducted using MEDLINE, Web of Science, SciELO, and the Korean Journal Database. Definitions were extracted from included manuscripts. We then conducted a three-stage Delphi consensus process to select the optimal outcome measures in terms of methodological quality and overall suitability for perioperative trials.

Results: From 2358 records, the full texts of 81 manuscripts were retrieved, of which 45 met the inclusion criteria. We identified three main categories of outcome measure specific to perioperative pulmonary outcomes: (i) composite outcome measures of multiple pulmonary outcomes (27 definitions); (ii) pneumonia (12 definitions); and (iii) respiratory failure (six definitions). These were rated by the group according to suitability for routine use. The majority of definitions were given a low score, and many were imprecise, difficult to apply consistently, or both, in large patient populations. A small number of highly rated definitions were identified as appropriate for widespread use. The group then recommended four outcome measures for future use, including one new definition.

Conclusions: A large number of postoperative pulmonary outcome measures have been used, but most are poorly defined. Our four recommended outcome measures include a new definition of postoperative pulmonary complications, incorporating an assessment of severity. These definitions will meet the needs of most clinical effectiveness trials of treatments to improve postoperative pulmonary outcomes.

Keywords: outcome assessment (healthcare)/standards; perioperative care/methods

Each year, more than 300 million patients undergo surgery worldwide.^{1,2} Estimates of attributable mortality vary from 1% to 4%, however some studies suggest that more than one in five deaths occur in a small group of high-risk patients.^{1,3–6} Even when complications are successfully treated, these are still associated with reduced long-term survival.⁷ Some of the most common postoperative complications affect the respiratory tract, with incidences ranging from 9% to 40%, depending on the definition used.^{8–11} A number of factors promote pulmonary complications amongst patients undergoing major surgery. Anaesthesia can lead to reduced functional residual capacity, hypoxaemia, and impaired central respiratory drive, while surgical manipulation can restrict ventilation, damage respiratory muscles, and cause atelectasis.¹² When combined with pre-existing respiratory disease and postoperative pain, the risk of pneumonia, respiratory failure, and death is increased.¹⁰ There is a need for large, high quality clinical trials to improve the treatment of patients at risk of postoperative pulmonary complications.

The impact of any large clinical trial is critically dependent on the use of well-defined outcome measures, which must be patient-centred, reliable, valid, and potentially modifiable by the trial intervention.^{13,14} Inconsistent reporting of outcomes across trials investigating similar clinical problems hinders the interpretation of new research findings in the context of existing research evidence, and prevents the use of pooled data in systematic reviews.¹⁴ Many definitions are similar but not sufficiently so to allow robust comparison. There is a recognised need for a standardised list of clearly defined clinical outcome measures for use in large, pragmatic clinical trials in this field.¹⁵ This would help investigators to improve the design of clinical trials through the use of recognised reference standards. This should lead to a stronger evidence base to inform clinical practice and improve short- and long-term outcomes for patients after surgery.

The Standardized Endpoints for Perioperative Medicine (StEP) collaboration was established to evaluate the literature and create standards for the definition and use of outcome measures in clinical effectiveness research in perioperative medicine, addressing the need for greater precision and consistency in defining outcomes for perioperative clinical trials.¹⁵ As part of this initiative, we performed a systematic literature review of clinical trials and observational studies of pulmonary outcomes in perioperative care, followed by a Delphi process to develop recommendations for standardised outcomes.

Methods

Members of the group were recruited to better understand and improve the pulmonary morbidity, which occurs during and after surgery. We performed a systematic literature review to identify studies with relevant clinical outcome measures that should be considered for future use in clinical trials. The group then evaluated these outcomes to decide which should be considered as evidence in a wider evaluation of clinical outcome measures in all areas of perioperative medicine.

Search strategy

MEDLINE, Web of Science, SciELO, and the Korean Journal Database were searched electronically from the year 2005 to present using the Web of Science (Thomson Reuters) platform. The search strategy was: TS=((Pneumonia[Title/Abstract] OR Respiratory[Title/Abstract] OR Pulmonary[Title/Abstract] OR 'Pulmonary Complications'[title/abstract] OR 'Respiratory complications'[title/abstract] OR Hypoxia[Title/Abstract] OR 'Lower Respiratory Tract'[Title/Abstract] OR Re-Intubation [Title/Abstract] OR 'Prolonged Ventilation'[Title/Abstract] OR 'Respiratory Failure'[Title/Abstract] OR 'Pulmonary Oedema'[Title/Abstract] OR 'Respiratory Distress Syndrome'[Title/Abstract] OR 'postoperative pulmonary complications'[title/abstract]

abstract] OR 'Acute Lung Injury'[Title/Abstract]) OR ('Lung Injury' OR 'Pneumonia' OR 'Pulmonary Edema' or 'Respiratory Distress Syndrome' OR Pneumothorax OR 'Respiratory Aspiration'[MeSH Terms])) AND TS=('Hospital Mortality' OR 'Mortality' OR 'Morbidity' OR 'Prognosis' OR 'Perioperative care' OR 'Intraoperative Complications' OR 'Fatal Outcome' OR 'Postoperative Complications' OR 'Outcome Assessment Health Care' OR 'Outcome and Process Assessment Health Care' OR 'Treatment Outcome') AND TS=(Mortality OR Morbidity OR Outcome OR 'Postoperative Complications' OR 'Prognosis' OR 'Perioperative Complications' OR 'Intraoperative Complications') AND TS=(('Surgical Procedures' OR 'Operative' OR 'Surgery' OR 'General Surgery' OR 'Operation') AND 'Postoperative Complications'). Search results were limited to those that were observational trials, randomised controlled trials, and meta-analyses conducted in adult subjects. The bibliography of evaluable studies and other selected papers were searched manually for potentially relevant citations.

Study selection criteria

Records were held within a record management system (Mendeley, London, UK) and duplicates were removed. The title and abstract of remaining records were then reviewed independently by two researchers (T.E.F.A. and A.J.F.). The full text paper of records identified as potentially relevant were downloaded and further assessed. Full texts were selected for inclusion if the study design was that of an observational study with prospective patient recruitment, a randomised controlled trial, or a meta-analysis and reported a respiratory outcome as the primary outcome measure. Articles were excluded if they included paediatric participants (<18 yr of age), had a primary outcome unrelated to the respiratory system, or did not provide a primary outcome definition. Differences in opinion were resolved by referral to a third researcher.

Data extraction

Data were extracted from the selected articles by two researchers acting independently. Data was extracted into a database (Excel 2007, Microsoft, Redmond, WA, USA) and included lead author name, year of publication, number of participants (and papers for meta-analyses), primary outcome measure, and definition of primary outcome measure.

Delphi process

The results of this systematic review informed a Delphi consensus process conducted during 2017 to obtain input and consensus from a group of medical and other health researchers with experience in anaesthesia and perioperative medicine trials.^{16,17} The StEP Working Group consisted of experienced perioperative trialists and other investigators from various countries and was overseen by a steering committee (see [Supplementary Material](#)). Each Delphi round was coordinated by the Health Services Research Centre of the Royal College of Anaesthetists in the UK or the Research Unit of the Department of Anaesthesia and Perioperative Medicine at the Alfred Hospital in Melbourne, Australia. The item scores and number of respondents were recorded for each of the above Delphi rounds in an Excel spreadsheet, and then converted to an SPSS database for calculation of final median and range scores, and consensus rates.

Stage 1: literature search to develop preliminary list of trial endpoints and definitions

Publications identified in the literature search were used to create a preliminary list of outcome measures and their definitions.

Stage 2: formal rating of the recommendations (Delphi round one)

We extracted the outcome measures from manuscripts identified in our literature search, and then created summary tables in different categories, allowing easy comparison of outcome measure definitions. We circulated these tables within the Pulmonary StEP theme group and asked members to submit their ratings via an online form. Manuscripts were first evaluated by using a 'traffic light' scale (green/amber/red) for methodological quality, and overall suitability as endpoints in the context of perioperative trials. This rating encompassed validity, reliability, ease of use, and frequency of use in the perioperative literature. Where two or more alternative definitions were very similar, we promoted use of the more precise and detailed alternative(s). Definitions that had been superseded or were clearly out of date were categorised as red.

Stage 3: Delphi round two

The second Delphi round included participants from the entire StEP Working Group ($n=75$). Participants were asked to score each of the items listed using a scale of 1–9, with 1–3 labelled 'Not that important or invalid', 4–6 labelled 'Important but requires revision', and 7–9 labelled 'Critical for inclusion'.^{18,19} Participants were given the option to select 'unsure' if they were unable to offer an opinion as to which category to apply to the item. Participants were invited to suggest any other endpoints, or definitions, or modifications to existing definitions that they believed should be included. A reminder email was sent to ensure prompt completion of the survey. The final numbers of respondents and item completions were recorded. The number of participants who scored the item and the median, and inter-quartile range of scores were quantified. Members of the Pulmonary StEP theme group were then invited to discuss the results via email. Any items not rated as critical (i.e. 70th centile score <7) but still with a median score of 7 or greater were retained for consideration in the second round. Lower-rated endpoint items identified for removal could be retained if they were considered as critical by any group member for the second round. Items with a median score of ≤ 3 were not retained.

Stage 4: Delphi round three and final recommendations

Delphi round three included members of the Pulmonary StEP theme group ($n=14$). The summary results of the above process were provided to these participants, inviting further comments. If responses to this final stage suggested a need for further modification to endpoint definitions, then this was resolved by the authors via email discussion.

Results

Study selection

We identified 2366 records from the electronic search strategy, and a further 13 records through other sources. After removal

Table 1 Alternative definitions of ‘postoperative pulmonary complications’ identified in the literature search. Includes composite respiratory outcomes recorded as: ‘postoperative pulmonary complications’, ‘major pulmonary complications’, ‘major respiratory complications’, ‘postoperative respiratory complications’, ‘postoperative respiratory outcomes’, ‘pulmonary complications’ or ‘respiratory complications’

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Agostini and colleagues ²⁴	Postoperative pulmonary complications	Four or more of: atelectasis or consolidation on chest X-ray; white cell count $>11.2 \times 10^9$ litre ⁻¹ or administration of respiratory antibiotics after operation (in addition to prophylactic antibiotics); temperature $>38^\circ\text{C}$; signs of infection on sputum microbiology; purulent sputum differing from preoperative status; oxygen saturations $<90\%$ on room air; physician diagnosis of pneumonia; or prolonged high dependency unit (HDU) stay or readmission to HDU or intensive care unit (ICU) for respiratory complications.	Red	—	—
Barrera and colleagues ²⁵	Postoperative pulmonary complications	Any of: respiratory failure requiring intensive care admission, intubation, or both; pneumonia (new pulmonary infiltrate with fever treated with i.v. antibiotics); atelectasis requiring bronchoscopy (need determined by the surgical team); pulmonary embolism (diagnosed by computed tomography scan and treated); and need for supplemental oxygen at hospital discharge.	Red	—	—
Bellinetti and colleagues ²⁶	Postoperative respiratory complications	Any of: pneumonia (core temperature more than 38°C , radiological signs of pulmonary consolidation, and productive cough); atelectasis with evident clinical repercussions; bronchospasm; prolonged mechanical ventilation; pleural effusion or pneumothorax; surgical re-intervention caused by inadequate lung re-expansion; or death related or not to pulmonary disease.	Amber	—	—
Canet and colleagues ²⁷	Postoperative pulmonary complications	Any of: respiratory infection (antibiotics for a suspected respiratory infection and at least one of: new or changed sputum, new or changed lung opacities, fever, leucocyte count $>12\,000\ \mu\text{l}^{-1}$), respiratory failure (postoperative $\text{PaO}_2 <60$ mm Hg on room air, a ratio of PaO_2 to inspired oxygen fraction <300 , or pulse oximetry $<90\%$ and requiring oxygen therapy), pleural effusion (chest X-ray demonstrating blunting of the costo-phrenic angle, loss of the sharp silhouette of the ipsilateral hemi-diaphragm when upright, displacement of adjacent anatomical structures, or a hazy opacity in one hemithorax with preserved vascular shadows when supine), atelectasis (lung opacification with a shift of the mediastinum, hilum or hemi-diaphragm, and compensatory over inflation in the adjacent non-atelectatic lung), pneumothorax (air in the pleural space with no vascular bed surrounding the visceral pleura), bronchospasm (newly detected expiratory wheezing), or aspiration pneumonitis (acute lung injury after the inhalation of regurgitated gastric contents).	Green	Median score 7 44/75 scored ≥ 7	Median score 7 6/10 scored ≥ 7
Casado and colleagues ²⁸	Respiratory complications	Any of: pneumonia confirmed by a positive culture result (tracheal aspiration or blood sample) or chest X-ray; acute lung injury ($\text{PaO}_2/\text{FiO}_2 <300$), or adult distress respiratory syndrome (ARDS) ($\text{PaO}_2/\text{FiO}_2 <200$ regardless of the level of PEEP, pulmonary capillary wedge pressure <18 mm Hg or bilateral infiltrates consistent with pulmonary oedema without clinical evidence of left ventricular failure) according to the American-European Consensus Conference (1994).	Red	—	—
Dronkers and colleagues ²⁹	Postoperative pulmonary complications	Operationalised as atelectasis, which is considered a ‘precursor’ of more clinically relevant postoperative pulmonary complications. A	Red	—	—

Continued

Table 1 Continued

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Futier and colleagues ³⁰	Major pulmonary complications	blinded radiologist evaluated radiographs of the lung base for the presence of atelectasis. Pneumonia defined according to standard criteria or the need for invasive or non-invasive ventilation for acute respiratory failure.	Red	—	—
Galvao Serejo and colleagues ³¹	Postoperative pulmonary complications	Any of: pneumonia (presence of new or progressive pulmonary infiltration on chest X-ray associated with at least two of: purulent trachea-bronchial secretion, elevated temperature >38.3°C, or increased blood leucocytes (>25% of base count); atelectasis (pulmonary atelectasis on chest X-ray associated with acute respiratory symptoms without fulfilling the criteria for pneumonia); pleural effusion (excessive fluid in the pleural space, detected by clinical examination and chest radiograph, requiring percutaneous intervention); acute respiratory failure (acute deficiency of gas exchange requiring mechanical ventilation). If patients with pneumonia developed respiratory failure they would be included only in the group 'acute respiratory failure'.	Amber	—	—
Grant and colleagues ³²	Postoperative pulmonary complications	Any of: acute lung injury, pulmonary infection, or atelectasis, as defined by the individual studies included in the meta-analysis.	Red	—	—
Guimaraes and colleagues ³³	Pulmonary complications	Any of: (i) atelectasis (radiographic, tomographic, or bronchoscopic diagnosis in patients whose clinical signs were acute respiratory symptoms such as dyspnoea, cough, wheeze); (ii) respiratory failure (radiographical diagnosis in patients with signs of acute respiratory symptoms such as tracheobronchial purulent secretions, fever (>38°C), or increased white blood cell (WBC) count (>10 000 mm ⁻³); or (iii) tracheobronchial infection or pneumonia.	Red	—	—
Haines and colleagues ³⁴	Postoperative pulmonary complications	Four or more of: chest radiograph report of collapse/consolidation; oral temperature >38°C on more than one consecutive postoperative day; pulse-oximetry (SpO ₂) <90% on more than one consecutive postoperative day; production of yellow or green sputum different to preoperative assessment; infection on sputum culture report; unexplained white cell count >11×10 ⁹ litre ⁻¹ or antibiotic therapy for respiratory infection; new abnormal breath sounds on auscultation different to preoperative assessment; or physician diagnosis of postoperative pulmonary complication.	Red	—	—
Hemmes and colleagues ³⁵	Postoperative pulmonary complications	Any of: hypoxaemia, severe hypoxaemia, bronchospasm, suspected pulmonary infection, pulmonary infiltrate, aspiration pneumonitis, development of Acute Respiratory Distress Syndrome, atelectasis, pleural effusion, pulmonary oedema caused by cardiac failure, and pneumothorax.	Green	Median score 6 32/75 scored ≥7	Median score 6 5/10 scored ≥7
Hodari and colleagues ³⁶	Postoperative respiratory outcomes	Respiratory outcomes tracked in National Surgical Quality Improvement Program (NSQIP): re-intubation (placement of an tracheal tube and mechanical or assisted ventilation because of respiratory or cardiac failure manifested by severe respiratory distress, hypoxia, hypercarbia, or respiratory acidosis within 30 days of surgery); postoperative pneumonia [one of two criteria: Criterion 1: rales or dullness to percussion on physical examination of chest plus: (i) new onset of purulent sputum or change in character of sputum; (ii) organism isolated from blood culture; or (iii) isolation of pathogen from specimen obtained by trans-tracheal aspirate, bronchial brushing, or biopsy. Criterion	Green	Median score 7 38/75 scored ≥7	Median score 6 4/10 scored ≥7

Continued

Table 1 Continued

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
		2: chest X-ray shows new or progressive infiltrate, consolidation, cavitation, or pleural effusion and any of: (i) new onset of purulent sputum or change in character of sputum; (ii) organism isolated from blood culture; (iii) isolation of pathogen from specimen obtained by trans-tracheal aspirate, bronchial brushing, or biopsy; (iv) isolation of virus or detection of viral antigen in respiratory secretions; (v) diagnostic single antibody titre (immunoglobulin M) or four-fold increase in paired serum samples (immunoglobulin G) for pathogen; or (vi) histopathologic evidence of pneumonia]; and prolonged ventilation (ventilator-assisted respirations during postoperative hospitalisation >48 h).			
Hulzebos and colleagues ⁴⁴	Postoperative pulmonary complications	Grade 1: cough, or micro-atelectasis (abnormal lung findings and temperature >37.5°C without other cause) or normal/unavailable chest x-ray, or dyspnoea not because of other documented cause. Grade 2: productive cough, not because of other cause, or bronchospasm (new wheezing or pre-existent wheezing resulting in change therapy), or hypoxemia (alveolar-arterial gradient >29 and symptoms of dyspnoea or wheezing) or atelectasis (radiological confirmation plus either temperature >37.5°C or abnormal lung findings) or hypercarbia (transient, requiring treatment, such as naloxone or increased manual or mechanical ventilation) or adverse reaction to pulmonary medication. Grade 3: pleural effusion, resulting in thoracentesis, or suspected pneumonia (radiological evidence without bacteriological confirmation) or proved pneumonia (radiological evidence with pathological organism by Gram stain or culture) or pneumothorax or postoperative re-intubation or intubation (period of ventilator dependence not >48 h). Grade 4: ventilatory failure (ventilator dependence >48 h) or re-intubation with ventilator dependence.	Green	Median score 7 37/75 scored ≥7	Median score 6 4/10 scored ≥7
Ireland and colleagues ¹¹	Major respiratory complications	Significant atelectasis, pneumonia, significant hypoxia, tracheal re-intubation, ICU admission. Atelectasis was defined by the authors of the individual studies.	Red	—	—
Ladha and colleagues ³⁸	Postoperative respiratory complications	Any of: re-intubation, respiratory failure, pneumonia, and pulmonary oedema within 3 days of the procedure.	Red	—	—
Li and colleagues ³⁹	Postoperative pulmonary complications	Pulmonary infection (fever with positive sputum culture and infiltration on chest X-ray), atelectasis, or hypoxemia (SaO ₂ <90% for >30 min).	Red	—	—
Lunardi and colleagues ⁴⁰	Postoperative pulmonary complications	Any of: atelectasis with clinical consequences, hypoxemia with oxygen saturation <85% and need of supplemental oxygen, pneumonia, or acute respiratory failure.	Red	—	—
Mackay and colleagues ⁴¹	Postoperative pulmonary complications	Three or more of the following signs within the same day, in the first 14 days after surgery: Auscultation changes (decreased breath sounds, crackles, wheezes, bronchial breathing) that were additional to those found before surgery. Temperature >38°C. Chest X-ray changes consistent with collapse, consolidation, or atelectasis. Increase in amount, changed colour, or both, of sputum produced, compared with what the patient reports is usual for them.	Red	—	—
McAlister and colleagues ⁴²	Postoperative pulmonary complications	Any one of: respiratory failure requiring mechanical ventilation, pneumonia, atelectasis requiring bronchoscopic intervention, and	Red	—	—

Continued

Table 1 Continued

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Muehling and colleagues ⁴³	Postoperative pulmonary complications	pneumothorax or pleural effusion requiring percutaneous intervention. Any one of: atelectasis; pneumonia; or prolonged air leak >7 days. Atelectasis was diagnosed on chest X-ray or computed tomography scan; pneumonia was confirmed if antibiotic medication was required because of clinical and radiological signs of infection combined with elevated white cell count or raised C-reactive protein (CRP).	Red	—	—
Nascimento and colleagues ⁴⁵	Pulmonary complications	Any one of: atelectasis (radiographic, tomographic, or bronchoscopic diagnosis in patients whose clinical signs were acute respiratory symptoms such as dyspnoea, cough, wheeze); respiratory failure (radiographical diagnosis in patients with signs of acute respiratory symptoms such as tracheobronchial purulent secretions, fever (>38°C), or increased WBC count (>10 000 mm ⁻³); tracheobronchial infection or pneumonia.	Red	—	—
Paisani and colleagues ⁴⁶	Postoperative pulmonary complications	One or more of: pneumonia (presence of radiological evidence of pulmonary infiltration associated with at least two of: purulent sputum, temperature >38.0°C or leucocytosis >25% above baseline value); trachea-bronchitis (marked increase in sputum production or presence of purulent sputum when chest X-ray normal); atelectasis (radiological evidence of atelectasis associated with dyspnoea); acute respiratory failure (acute deficiency of gas exchange with necessity for invasive or non-invasive mechanical ventilation); bronchoconstriction (wheezing associated with dyspnoea, requiring bronchodilators or change in previous dosage of bronchodilator).	Amber	—	—
Scholes and colleagues ⁴⁷	Pulmonary complications	Four or more of: chest radiograph report of collapse/consolidation; oral temperature >38°C on more than one consecutive postoperative day; pulse-oximetry (SpO ₂) <90% on more than one consecutive postoperative day; production of yellow or green sputum different to preoperative assessment; infection on sputum culture report; unexplained white cell count >11×10 ⁹ litre ⁻¹ or antibiotic therapy for respiratory infection; new abnormal breath sounds on auscultation different to preoperative assessment; or physician diagnosis of postoperative pulmonary complication.	Red	—	—
Vidotto and colleagues ⁴⁸	Postoperative pulmonary complications	Any one of: acute respiratory infection (pneumonia or purulent trachea-bronchitis, where pneumonia was established by the presence of lung infiltration on chest X-ray associated with at least two of: purulent tracheobronchial secretion, temperature >38.3°C or a 25% increase in baseline blood leucocytes; and purulent trachea-bronchitis was diagnosed when tracheobronchial secretions increased in amount or changed in colour or purulence, associated with a normal chest X-ray; atelectasis (acute respiratory symptoms associated with radiological imaging); bronchospasm (wheezing detectable with a stethoscope associated with acute respiratory symptoms and the need for medication therapy).	Amber	—	—
Yang and colleagues ⁴⁹	Postoperative pulmonary complications	PaO ₂ /FiO ₂ >300 mm Hg, the presence of newly developed lung lesions (lung infiltration and atelectasis), or both, within 72 h of the operation.	Red	—	—
Zarbock and colleagues ⁵⁰	Postoperative pulmonary complications	Any one of: decreased pulmonary oxygen transfer ratio, nosocomial pneumonia, re-intubation rate, and readmission to the ICU after elective discharge to the general ward.	Red	—	—

Table 2 Definitions of pneumonia outcome measures

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Akutsu and colleagues ⁵¹	Postoperative pneumonia	All of: increase of sputum, opacity on chest X-ray, consolidation on computed tomography, increased temperature, increased WBC count, and CRP value in the serum.	Red	—	—
Cassidy and colleagues ⁵²	Postoperative pneumonia	At least one definitive chest X-ray, and at least one sign (fever, leucocytosis, or altered mental status with no other cause), and at least one microbiologic laboratory finding (positive cultures from blood, broncho-alveolar lavage, or pleural fluid) or at least two symptoms (purulent sputum, worsening cough, dyspnoea or tachypnoea, rales or rhonchi, or worsening gas exchange).	Amber	Median score 5 20/75 scored ≥ 7	Median score 5 3/10 scored ≥ 7
Markar and colleagues ⁵³	Postoperative pneumonia	Pyrexia, productive cough, raised white cell count, and localising signs on chest examination or chest radiography.	Red	—	—
Schussler and colleagues ⁵⁴ (2006)	Postoperative pneumonia	Fibreoptic bronchoscopy samples were obtained before antibiotic therapy in every patient with signs of pneumonia: abnormal radiographic findings (new or changing radiographic infiltrates that persisted after physiotherapy or broncho-aspiration), fever $>38^{\circ}\text{C}$, and one of: a new increase in CRP or WBC count over the last 24 h (with $\text{WBC} > 12 \times 10^9 \text{ litre}^{-1}$) or increased and modified expectorate, possibly purulent. Patients with positive plugged telescopic catheter sample [$>103 \text{ colony-forming unit (CFU ml}^{-1})$], protected specimen brush sample ($>103 \text{ CFU ml}^{-1}$), or positive blood culture represented the 'documented POP' group. If the significant cut-off values were not reached, but clinical and radiologic improvement occurred after the administration of antibiotics, patients were considered as having 'non-documented POP.'	Red	—	—
Schussler and colleagues ⁵⁵ (2008)	Postoperative pneumonia	Quantitative fibreoptic bronchoscopy aspiration, plugged telescopic catheter (PTC), or protected specimen brush (PSB) sampling in case of: abnormal radiographic findings (new or changing radiographic infiltrates that persisted after physiotherapy or fibreoptic bronchial aspiration), fever $>38^{\circ}\text{C}$ and one of: purulent secretions or an increase of $>30\%$ of the CRP or WBC count during the last 24 h (with count $>12 \times 10^9 \text{ litre}^{-1}$). Pneumonia was considered documented if bacteria were identified in blood culture or at the 48-h culture of the fibreoptic sample with thresholds: PTC or PSB at $103 \geq \text{CFU ml}^{-1}$, or bilateral quantitative endobronchial aspirate (QEBA) at $\geq 106 \text{ CFU ml}^{-1}$.	Red	—	—
Bagyi and colleagues ⁵⁶	Pneumonia	According to American Thoracic Society guidelines by X-ray (new or progressive radiographic infiltrate), new onset of fever $>38.0^{\circ}\text{C}$, symptoms of coughing with purulent sputum, chest pain and leucocytosis ($\text{WBC} > 10 \text{ g litre}^{-1}$).	Red	—	—
Centers for Disease Control definition ²⁰	Pneumonia	Two or more serial chest radiographs with at least one of the following (one radiograph is sufficient for patients with no underlying pulmonary or cardiac disease): (i) New or progressive and persistent infiltrates, (ii) consolidation, (iii) cavitation; at least one of the following: (a) fever ($>38^{\circ}\text{C}$) with no other recognised cause, (b) leucopenia (white cell count $< 4 \times 10^9 \text{ litre}^{-1}$) or leucocytosis (white cell count $> 12 \times 10^9 \text{ litre}^{-1}$), (c) for adults $>70 \text{ yr}$ old, altered mental status with no other recognised cause; And at least two of the following:	Green	Median score 8 56/75 scored ≥ 7	Median score 8 10/10 scored ≥ 7

Continued

Table 2 Continued

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Bouza and colleagues ⁵⁷	Ventilator associated pneumonia	(a) new onset of purulent sputum or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements, (b) new onset or worsening cough, or dyspnoea, or tachypnoea, (c) rales or bronchial breath sounds, (d) worsening gas exchange (hypoxaemia, increased oxygen requirement, increased ventilator demand). Mechanical ventilation for 48 h when the presence of new, progressive, or both, pulmonary infiltrates on a chest X-ray plus two or more of: temperature >38.5°C or <36°C; leucocytosis (>12×10 ⁹ cells litre ⁻¹); purulent tracheobronchial secretions; or a reduction in the PaO ₂ fraction of inspired oxygen ratio ~15% according to Centers for Disease Control and Prevention definitions. Patients with a clinical pulmonary infection score >6 were also considered to have pneumonia.	Red	—	—
Hortal and colleagues ³⁷	Ventilator associated pneumonia	New, progressive, or both, pulmonary infiltrates on chest radiograph in a patient ventilated more than 48 h plus two or more of: fever >38.5°C or hypothermia <36°C, leucocytosis >12×10 ⁹ litre ⁻¹ , purulent tracheobronchial secretions or a reduction of PaO ₂ /FiO ₂ >15% in the last 48 h according to the Centers for Disease Control definitions. We also included as pneumonia, those with a clinical pulmonary infection score >6. Early-onset pneumonia was defined as occurring within the first 4 days of hospitalisation and late-onset pneumonia as occurring on Day 5 or later on. The isolation of one or more microorganisms in significant bacterial count was required to confirm the diagnosis.	Red	—	—
Jiao and colleagues ⁵⁸	Ventilator associated pneumonia	Chest X-ray results, with at least two of: fever >38°C; a white blood cell count of >11 000 or <3000 µl ⁻¹ ; or the presence of purulent endotracheal secretions. Microbiological samples were obtained by fiberoptic bronchoalveolar lavage (BAL), and the growth of ≥10 ⁴ CFU ml ⁻¹ microorganism culture of BAL was considered to be positive.	Red	—	—
Nicolosi and colleagues ⁵⁹	Ventilator associated pneumonia	Within 48 h of intubation or 72 h after extubation, evidence of a new lung infiltrate on chest X-ray and at least two of: leucocytosis, fever, or purulent tracheobronchial secretion.	Red	—	—
Severgnini and colleagues ²²	Clinical pulmonary infection score	The modified clinical pulmonary infection score was calculated by a modified original score as described by Pelosi and colleagues. ⁶⁰	Amber	Median score 5 17/75 scored ≥7	Median score 4 2/10 scored ≥7

of duplicates, 2358 records remained. The full texts of 81 manuscripts were retrieved, of which 45 met the inclusion criteria (Supplementary Fig. S1 and Table S1). A summary of full texts excluded after review, with reason for exclusion, is shown in Supplementary Table S1. The definitions extracted from included studies are summarised in Table 1. We identified three main categories of outcome measure specific to pulmonary outcomes in perioperative medicine: (i) composite outcome measures including a number of pulmonary outcomes (27 definitions); (ii) pneumonia (12 definitions); and (iii) respiratory failure (six definitions). These are summarised in Supplementary Table S2, Tables 1–3, along with the rating

(red/amber/green) given by the group and a summary definition of each outcome measure.

Postoperative pulmonary complications

Despite a large number of candidate definitions, the group was unable to reach a consensus on the best definition of postoperative pulmonary complications. Many definitions used component definitions, which differed widely in severity and underlying biological mechanism (e.g. atelectasis and acute respiratory distress). After a discussion, and having sought advice from the STEP steering committee, we have

Table 3 Definitions of respiratory failure outcome measures. *Chest radiograph or computed tomography scan. †If altitude is higher than 1000 m, a correction factor should be calculated [$\text{PaO}_2:\text{FiO}_2 \times (\text{barometric pressure}/101 \text{ kPa})$]. ‡This may be delivered non-invasively in the mild Acute Respiratory Distress Syndrome group

Lead author	Outcome measure	Definition	Round 1	Round 2	Round 3
Perkins and colleagues ⁶¹	Acute lung injury	American-European Consensus Conference definition: acute onset of bilateral infiltrates on the chest radiograph and hypoxaemia ($\text{PaO}_2/\text{FiO}_2$ ratio of <300) in the absence of clinical evidence of left atrial hypertension	Red	—	—
Canadian Orthopedic Trauma ⁶²	Acute Respiratory Distress Syndrome	1994 American-European consensus definition: (i) acute onset, (ii) bilateral chest infiltrate, (iii) pulmonary artery occlusion pressure of 18 mm Hg or less and no evidence of left atrial hypertension, (iv) impaired oxygenation regardless of the level of PEEP, and (v) a $\text{PaO}_2/\text{FiO}_2$ ratio of <200.	Red	—	—
ARDS Definition Task Force ⁶³	Acute Respiratory Distress Syndrome	The Berlin definition of Respiratory Distress Syndrome: Timing. Within 1 week of a known clinical insult or new or worsening respiratory symptoms and Chest imaging.* Bilateral opacities not fully explained by effusions, lobar/lung collapse or nodules and Origin of oedema. Respiratory failure not fully explained by cardiac failure or fluid overload. Need objective assessment (e.g. echocardiography) to exclude hydrostatic oedema Oxygenation.† Mild: $\text{PaO}_2:\text{FiO}_2$ between 26.7 and 40.0 kPa (200–300 mm Hg) with PEEP or CPAP ≥ 5 cm H ₂ O.‡ Moderate: $\text{PaO}_2:\text{FiO}_2$ between 13.3 and 26.6 kPa (100–200 mm Hg) with PEEP ≥ 5 cm H ₂ O. Severe: $\text{PaO}_2:\text{FiO}_2$ 13.3 kPa (100 mm Hg) with PEEP ≥ 5 cm H ₂ O	Green	Median score 8 53/75 scored ≥ 7	Median score 7 6/10 scored ≥ 7
Camp and colleagues ⁶⁴	Early extubation	Removal of breathing turs after arrival to cardiovascular intensive care.	Red	—	—
Squadrone and colleagues ⁶⁵	Tracheal intubation	Intubation was performed when patients presented with one of: severe hypoxemia (arterial oxygen saturation <80% despite maximal FiO_2 ; respiratory acidosis (arterial pH <7.30 with carbon dioxide tension >50 mm Hg; signs of patient distress with accessory muscle recruitment and paradoxical abdominal or thoracic motion; hemodynamic instability (80–90 mm Hg increase or a 30–40 mm Hg decrease in systolic blood pressure relative to the baseline value or need for inotropic drugs for at least 2 h to maintain systolic blood pressure higher than 85 mm Hg or electrocardiogram evidence of ischemia or significant ventricular arrhythmias); need for sedation for major agitation; decreased alertness defined as a Glasgow Coma Score <9; or cardiac arrest.	Amber	Median score 5 26/75 scored ≥ 7	Median score 5 4/10 scored ≥ 7
Fernandez-Perez and colleagues ²¹	Respiratory failure	Need for mechanical ventilation for >48 h after operation or need for re-institution of mechanical or non-invasive ventilation after extubation	Amber	Median score 6 25/75 scored ≥ 7	Median score 5 1/10 scored ≥ 7

recommended a new definition which allows lower severity pulmonary events to be captured as part of the overall clinical outcome (Box 1).

Pneumonia

Only the US Centers for Disease Control definition was recommended.²⁰ Whilst there are numerous other well-written definitions, there seemed little value in departing from such a widely used international standard (Box 2).

Respiratory failure

The group recommended two definitions covering overlapping patient scenarios. The Berlin definition of Acute Respiratory Distress Syndrome was recommended, again because this is a widely used international standard, which has already been carefully developed in a consensus process. However, as many important episodes of postoperative respiratory failure are not severe enough to meet this definition, the group also recommended an adapted version of a definition reported by

Fernandez-Perez and colleagues,²¹ which describes re-institution of mechanical or non-invasive ventilation after extubation (Box 3).

Discussion

We have summarised and evaluated a range of definitions used to describe pulmonary outcomes in perioperative medicine research. Many definitions were imprecise, or difficult to apply consistently in large patient populations because of the requirement from resource intensive diagnostic tests (e.g. bronchoscopy). The composite outcome of postoperative pulmonary complications was widely used, but the group was concerned about the lack of equivalence of component outcomes and the differing biological mechanisms, which may cause these. We therefore propose a new definition for this outcome. Importantly, this new definition includes a measure of severity which we hope will significantly reduce the subjectivity in evaluating pulmonary outcomes, and ensure a more valid comparison between studies. The group also

Box 1

Recommended definition of postoperative pulmonary complications.

Postoperative pulmonary complications

Mechanism

Composite of respiratory diagnoses that share common pathophysiological mechanisms including pulmonary collapse and airway contamination:

- (i) atelectasis detected on computed tomography or chest radiograph,
- (ii) pneumonia using US Centers for Disease Control criteria,
- (iii) Acute Respiratory Distress Syndrome using Berlin consensus definition,
- (iv) pulmonary aspiration (clear clinical history AND radiological evidence).

Severity

None: planned use of supplemental oxygen or mechanical respiratory support as part of routine care, but not in response to a complication or deteriorating physiology. Therapies which are purely preventive or prophylactic for example high flow nasal oxygen or continuous positive airways pressure (CPAP) should be recorded as none
 Mild: therapeutic supplemental oxygen <0.6 FiO₂
 Moderate: therapeutic supplemental oxygen ≥ 0.6 FiO₂, requirement for high-flow nasal oxygen, or both
 Severe: unplanned non-invasive mechanical ventilation, CPAP, or invasive mechanical ventilation requiring tracheal intubation

Exclusions

Other diagnoses that do not share a common biological mechanism are best evaluated separately and only when clearly relevant to the treatment under investigation:

- (i) pulmonary embolism,
- (ii) pleural effusion,
- (iii) cardiogenic pulmonary oedema,
- (iv) pneumothorax,
- (v) bronchospasm.

Box 2

Recommended definition of postoperative pneumonia.

US Centers for Disease Control definition of pneumonia

Two or more serial chest radiographs with at least one of the following (one radiograph is sufficient for patients with no underlying pulmonary or cardiac disease):

(i) New or progressive and persistent infiltrates, (ii) consolidation, (iii) cavitation; AND at least one of the following:

- (a) fever ($>38^{\circ}\text{C}$) with no other recognised cause,
- (b) leucopaenia (white cell count $<4 \times 10^9$ litre⁻¹) or leucocytosis (white cell count $>12 \times 10^9$ litre⁻¹),
- (c) for adults >70 yr old, altered mental status with no other recognised cause;

AND at least two of the following:

- (a) new onset of purulent sputum or change in character of sputum, or increased respiratory secretions, or increased suctioning requirements,
- (b) new onset or worsening cough, or dyspnoea, or tachypnoea,
- (c) rales or bronchial breath sounds,
- (d) worsening gas exchange (hypoxaemia, increased oxygen requirement, increased ventilator demand).

recommended one outcome to describe pneumonia and two outcomes to describe respiratory failure.

The adverse events identified do not represent a comprehensive list of all that may occur. Instead, we have focused on those considered most important in a mixed population of surgical patients and relevant to perioperative care as opposed to specific technical complications of surgery or anaesthesia.

Box 3

Recommended definition of postoperative respiratory failure.

Berlin definition of Respiratory Distress Syndrome

Timing: within 1 week of a known clinical insult or new or worsening respiratory symptoms AND...

Chest imaging: bilateral opacities not fully explained by effusions, lobar/lung collapse or nodules AND...

Origin of oedema: respiratory failure not fully explained by cardiac failure or fluid overload (requires objective assessment, e.g. echocardiography, to exclude hydrostatic oedema), AND...

Oxygenation: mild PaO₂:FiO₂ between 26.7 and 40.0 kPa (200–300 mm Hg) with PEEP or CPAP ≥ 5 cm H₂O; moderate PaO₂:FiO₂ between 13.3 and 26.6 kPa (100–200 mm Hg) with PEEP ≥ 5 cm H₂O; severe PaO₂:FiO₂ ≤ 13.3 kPa (100 mm Hg) with PEEP ≥ 5 cm H₂O.

Mechanical ventilation

The need for need for tracheal re-intubation and mechanical ventilation after extubation, and within 30 days after surgery OR mechanical ventilation for more than 24 h after surgery. The inclusion of non-invasive ventilation may be considered on a study by study basis.

In some cases, we did not identify existing research that utilised outcomes that would be considered important. An example is pulmonary thromboembolism, which can only be reliably detected through screening of all participating patients in a given trial. Some outcomes may be more appropriate for use in smaller studies or early phase trials (e.g. clinical pulmonary infection score).²² As previously noted, the physiological changes that follow surgery and anaesthesia, including the systemic inflammatory response, may result in partial compliance with the criteria for a number of outcome measures, in particular atelectasis. We note that reports relating to perioperative medicine are not easily identified through a literature search because there is no standard approach to the use of Medical Subject Headings (MeSH) terms by authors of perioperative medicine research articles. There is an argument for creating a new MeSH heading of 'perioperative medicine' to ensure that trials in this field are more easily identified.

The strengths of this project include a detailed and extensive systematic literature search followed by a robust three stage Delphi process including a core group of experts in postoperative pulmonary complications and the wider StEP group, which includes many of the leading international experts in perioperative care research. However, this work is limited by the large number of alternative definitions which have been used in previous research. In particular, we identified a large number of composite outcomes for postoperative pulmonary complications which included a diverse range of adverse events which differed widely in frequency, severity, and biological mechanism. As a result, the group was unable to make a strong recommendation for this important outcome, and instead we have recommended a new definition which we believe is less prone to methodological bias.

Conclusions

We have summarised and evaluated a wide range of pulmonary outcome measures in perioperative medicine research, most of which were poorly defined. Our four recommended outcome measures include a new definition of postoperative pulmonary complications, incorporating an assessment of severity. Together, these definitions will meet the needs of most clinical effectiveness trials of treatments to improve postoperative pulmonary outcomes, although not every definition will be relevant to every trial. This work represents part of an international initiative to improve the definition and use of outcome measures in clinical effectiveness trials in perioperative medicine. These projects include the recent work by the European Society of Anaesthesiology and European Society of Intensive Care Medicine joint taskforce on perioperative outcome measures,²⁰ and the Core Outcome Measures in Perioperative and Anaesthetic Care (COMPAC) project which is part of the Core Outcome Measures in Effectiveness Trials (COMET) initiative.²³

Authors' contributions

Contributed to protocol design, data acquisition, analysis, and preparation of the manuscript: all authors

Declaration of interest

R.M.P. holds research grants, and has given lectures, performed consultancy work, or both, for Nestle Health Sciences,

BBraun, Medtronic, Glaxo Smithkline, Intersurgical, and Edwards Lifesciences, and is a member of the associate editorial board of the *British Journal of Anaesthesia*. M.G.A. has given lectures, performed consultancy, or both, for Drägerwerk AG, Lübeck, Germany, and Ventinova Ltd, Eindhoven, The Netherlands. M.G. is an elected council member of the Royal College of Anaesthetists. T.G. is a member of the Editorial Board of *Anesthesia and Analgesia*. E.F. reported receiving consulting fees from Edwards Lifesciences and Dräger; lecture fees from Dräger, GE Healthcare, Edwards Lifesciences, Fisher and Paykel Healthcare, Fresenius Kabi, and Getinge. J.C. is Associate Editor of *Anesthesiology*. M.G. is the National Specialty Lead for Anaesthesia, Perioperative Medicine and Pain within the UK National Institute of Health Research Clinical Research Network, an elected council member of the Royal College of Anaesthetists and serves on the board of Oxygen Control Systems Ltd, the Evidence Based Perioperative Medicine (EBPOM) social enterprise and the medical advisory board of Sphere Medical Ltd. M.G. has received honoraria for speaking, travel expenses, or both, from Edwards Lifesciences, Fresenius-Kabi, BOC Medical (Linde Group), Ely-Lilly Critical Care, and Cortex GmbH. M.G. is Executive Chair of the Xtreme-Everest Oxygen Research Consortium and joint Editor-in-Chief of *Perioperative Medicine*. M.M. is a paid consultant for Edwards Lifesciences, Deltex and Baxter; his chair at UCL is sponsored by Smiths Medical; Director Evidence Based Perioperative Medicine (EBPOM) Community Interest Company, director of Medinspire Ltd, Clinical Hydration Solutions, Clinical Fabric Solutions Ltd, Oxygen Control Systems Ltd, and a share-holder and scientific advisor for Medical Defense Technologies LLC. M.M. sits on the editorial boards of the *British Journal of Anaesthesia*, *Critical Care* and is Editor-in-Chief of *Perioperative Medicine*. All other authors declare no conflicts of interest.

Funding

The Standardized Endpoints for Perioperative Medicine (StEP) Collaborative is supported by an unrestricted grant from the *British Journal of Anaesthesia*. T.E.F.A. was supported by a Medical Research Council and *British Journal of Anaesthesia* Clinical Research Training Fellowship. R.M.P. is a National Institute for Health Research (UK) Professor.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.bja.2018.02.007>.

References

- Abbott TEF, Fowler AJ, Dobbs TD, Harrison EM, Gillies MA, Pearse RM. Frequency of surgical treatment and related hospital procedures in the UK: a national ecological study using hospital episode statistics. *Br J Anaesth* 2017; **119**: 249–57
- Weiser TG, Haynes AB, Molina G, et al. Estimate of the global volume of surgery in 2012: an assessment supporting improved health outcomes. *Lancet* 2015; **385**(Suppl 2): S11
- Pearse RM, Holt PJ, Grocott MP. Managing perioperative risk in patients undergoing elective non-cardiac surgery. *Br Med J* 2011; **343**: d5759

4. Pearse RM, Moreno RP, Bauer P, et al. Mortality after surgery in Europe: a 7 day cohort study. *Lancet* 2012; **380**: 1059–65
5. International Surgical Outcomes Study group. Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries. *Br J Anaesth* 2016; **117**: 601–9
6. Jhanji S, Thomas B, Ely A, Watson D, Hinds CJ, Pearse RM. Mortality and utilisation of critical care resources amongst high-risk surgical patients in a large NHS trust. *Anaesthesia* 2008; **63**: 695–700
7. Khuri SF, Henderson WG, DePalma RG, et al. Determinants of long-term survival after major surgery and the adverse effect of postoperative complications. *Ann Surg* 2005; **242**: 326–41
8. Ackland GL, Iqbal S, Paredes LG, et al. Individualised oxygen delivery targeted haemodynamic therapy in high-risk surgical patients: a multicentre, randomised, double-blind, controlled, mechanistic trial. *Lancet Respir Med* 2015; **3**: 33–41
9. Warner DO. Preventing postoperative pulmonary complications: the role of the anesthesiologist. *Anesthesiology* 2000; **92**: 1467–72
10. Arozullah AM, Daley J, Henderson WG, Khuri SF. Multifactorial risk index for predicting postoperative respiratory failure in men after major noncardiac surgery. The National Veterans Administration Surgical Quality Improvement Program. *Ann Surg* 2000; **232**: 242–53
11. Ireland CJ, Chapman TM, Mathew SF, Herbison GP, Zacharias M. Continuous positive airway pressure (CPAP) during the postoperative period for prevention of postoperative morbidity and mortality following major abdominal surgery. *Cochrane Database Syst Rev* 2014; **8**, CD008930-CD
12. Nunn JF, Payne JP. Hypoxaemia after general anaesthesia. *Lancet* 1962; **2**: 631–2
13. Koroshetz W. A core set of trial outcomes for every medical discipline? *Br Med J* 2015; **350**: h85
14. Tovey D. The impact of cochrane reviews. *Cochrane Database Syst Rev* 2011; **2011**. ED000007
15. Myles PS, Grocott MP, Boney O, Moonesinghe SR, Group CO-S. Standardizing end points in perioperative trials: towards a core and extended outcome set. *Br J Anaesth* 2016; **116**: 586–9
16. Diamond IR, Grant RC, Feldman BM, et al. Defining consensus: a systematic review recommends methodologic criteria for reporting of Delphi studies. *J Clin Epidemiol* 2014; **67**: 401–9
17. Sinha IP, Smyth RL, Williamson PR. Using the Delphi technique to determine which outcomes to measure in clinical trials: recommendations for the future based on a systematic review of existing studies. *PLoS Med* 2011; **8**. e1000393
18. Kirkham JJ, Gorst S, Altman DG, et al. COS-STAR: a reporting guideline for studies developing core outcome sets (protocol). *Trials* 2015; **16**: 373
19. Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 2011; **64**: 395–400
20. Jammer I, Wickboldt N, Sander M, et al. Standards for definitions and use of outcome measures for clinical effectiveness research in perioperative medicine: European Perioperative Clinical Outcome (EPCO) definitions: a statement from the ESA-ESICM joint taskforce on perioperative outcome measures. *Eur J Anaesth* 2015; **32**: 88–105
21. Fernandez-Perez ER, Sprung J, Afessa B, et al. Intraoperative ventilator settings and acute lung injury after elective surgery: a nested case control study. *Thorax* 2009; **64**: 121–7
22. Severgnini P, Selmo G, Lanza C, et al. Protective mechanical ventilation during general anaesthesia for open abdominal surgery improves postoperative pulmonary function. *Anesthesiology* 2013; **118**: 1307–21
23. Comet initiative. *Core Outcome Measures in Perioperative and Anaesthetic Care (COMPAC)*. 2016. Available from: <http://www.comet-initiative.org/studies/details/632> (accessed)
24. Agostini P, Cieslik H, Rathinam S, et al. Postoperative pulmonary complications following thoracic surgery: are there any modifiable risk factors? *Thorax* 2010; **65**: 815–8
25. Barrera R, Shi WJ, Amar D, et al. Smoking and timing of cessation—impact on pulmonary complications after thoracotomy. *Chest* 2005; **127**: 1977–83
26. Bellinetti LM, Thomson JC. Respiratory muscle evaluation in elective thoracotomies and laparotomies of the upper abdomen. *J Bras Pneumol* 2006; **32**: 99–105
27. Canet J, Gallart L, Gomar C, et al. Prediction of postoperative pulmonary complications in a population-based surgical cohort. *Anesthesiology* 2010; **113**: 1338–50
28. Casado D, Lopez F, Marti R. Perioperative fluid management and major respiratory complications in patients undergoing esophagectomy. *Dis Esophagus* 2010; **23**: 523–8
29. Dronkers J, Veldman A, Hoberg E, van der Waal C, van Meeteren N. Prevention of pulmonary complications after upper abdominal surgery by preoperative intensive inspiratory muscle training: a randomized controlled pilot study. *Clin Rehabil* 2008; **22**: 134–42
30. Futier E, Constantin JM, Paugam-Burtz C, et al. A trial of intraoperative low-tidal-volume ventilation in abdominal surgery. *New Engl J Med* 2013; **369**: 428–37
31. Galvao Serejo LG, da Silva Junior FP, Catunda Bastos JP, de Bruin GS, Salani Mota RM, Carvalhede de Bruin PF. Risk factors for pulmonary complications after emergency abdominal surgery. *Respir Med* 2007; **101**: 808–13
32. Grant MC, Yang D, Stone A, Wu CL, Wick EC. A meta-analysis of intraoperative ventilation strategies to prevent pulmonary complications: is low tidal volume alone sufficient to protect healthy lungs? *Ann Surg* 2016; **263**: 881–7
33. Guimaraes MMF, El Dib R, Smith AF, Matos D. Incentive spirometry for prevention of postoperative pulmonary complications in upper abdominal surgery. *Cochrane Database Syst Rev* 2009; **3**, CD006058
34. Haines KJ, Skinner EH, Berney S, Austin Hlth PS. Association of postoperative pulmonary complications with delayed mobilisation following major abdominal surgery: an observational cohort study. *Physiotherapy* 2013; **99**: 119–25
35. Hemmes SNT, Gama de Abreu M, Pelosi P, Schultz MJ, the PNIftCTNo, European Society of A. High versus low positive end-expiratory pressure during general anaesthesia for open abdominal surgery (PROVHILO trial): a multicentre randomised controlled trial. *Lancet* 2014; **384**: 495–503
36. Hodari A, Tsiouris A, Eichenhorn M, Horst M, Rubinfeld I. Exploring National Surgical Quality Improvement Program respiratory comorbidities: developing a predictive understanding of postoperative respiratory occurrences,

- Clavien 4 complications, and death. *J Surg Res* 2013; **183**: 663–7
37. Hortal J, Giannella M, Jesus Perez M, et al. Incidence and risk factors for ventilator-associated pneumonia after major heart surgery. *Intensive Care Med* 2009; **35**: 1518–25
 38. Ladha K, Vidal Melo MF, McLean DJ, et al. Intraoperative protective mechanical ventilation and risk of postoperative respiratory complications: hospital based registry study. *Br Med J* 2015; **351**: h3646
 39. Li Q, Yao G, Zhu X. High-dose ambroxol reduces pulmonary complications in patients with acute cervical spinal cord injury after surgery. *Neurocrit Care* 2012; **16**: 267–72
 40. Lunardi AC, Paisani DM, Marques da Silva CCB, Cano DP, Tanaka C, Carvalho GRF. Comparison of lung expansion techniques on thoracoabdominal mechanics and incidence of pulmonary complications after upper abdominal surgery: a randomized and controlled trial. *Chest* 2015; **148**: 1003–10
 41. Mackay MR, Ellis E, Johnston C. Randomised clinical trial of physiotherapy after open abdominal surgery in high risk patients. *Aust J Physiother* 2005; **51**: 151–9
 42. McAlister FA, Bertsch K, Man J, Bradley J, Jacka M. Incidence of and risk factors for pulmonary complications after nonthoracic surgery. *Am J Respir Crit Care Med* 2005; **171**: 514–7
 43. Muehling BM, Halter GL, Schelzig H, et al. Reduction of postoperative pulmonary complications after lung surgery using a fast track clinical pathway. *Eur J Cardiothorac Surg* 2008; **34**: 174–80
 44. Hulzebos EHJ, Helders PJM, Favie NJ, DeBrie RA, de la Riviere AB, Van Meeteren NLU. Preoperative intensive inspiratory muscle training to prevent postoperative pulmonary complications in high-risk patients undergoing CABG surgery – a randomised clinical trial. *JAMA* 2006; **296**: 1851–7
 45. Junior Pd Nascimento, Modolo NSP, Andrade S, et al. Incentive spirometry for prevention of postoperative pulmonary complications in upper abdominal surgery. *Cochrane Database Syst Rev* 2014; **2**, CD006058-CD
 46. Paisani DM, Fiore Jr JF, Lunardi AC, et al. Preoperative 6-min walking distance does not predict pulmonary complications in upper abdominal surgery. *Respirology* 2012; **17**: 1013–7
 47. Scholes RL, Browning L, Sztendur EM, Denehy L. Duration of anaesthesia, type of surgery, respiratory co-morbidity, predicted VO(2)max and smoking predict postoperative pulmonary complications after upper abdominal surgery: an observational study. *Aust J Physiother* 2009; **55**: 191–8
 48. Vidotto MC, Sogame LC, Gazzotti MR, Prandini M, Jardim JR. Implications of extubation failure and prolonged mechanical ventilation in the postoperative period following elective intracranial surgery. *Braz J Med Biol Res* 2011; **44**: 1291–8
 49. Yang M, Ahn HJ, Kim K, et al. Does a protective ventilation strategy reduce the risk of pulmonary complications after lung cancer surgery? A randomized controlled trial. *Chest* 2011; **139**: 530–7
 50. Zarbock A, Mueller E, Netzer S, Gabriel A, Feindt P, Kindgen-Milles D. Prophylactic nasal continuous positive airway pressure following cardiac surgery protects from postoperative pulmonary complications: a prospective, randomized, controlled trial in 500 patients. *Chest* 2009; **135**: 1252–9
 51. Akutsu Y, Matsubara H, Shuto K, et al. Pre-operative dental brushing can reduce the risk of postoperative pneumonia in esophageal cancer patients. *Surgery* 2010; **147**: 497–502
 52. Cassidy MR, Rosenkranz P, McCabe K, Rosen JE, McAneny DI. COUGH: reducing postoperative pulmonary complications with a multidisciplinary patient care program. *JAMA Surg* 2013; **148**: 740–5
 53. Markar SR, Walsh SR, Griffin K, Khandanpour N, Tang TY, Boyle JR. Assessment of a multifactorial risk index for predicting postoperative pneumonia after open abdominal aortic aneurysm repair. *Vascular* 2009; **17**: 36–9
 54. Schussler O, Alifano M, Dermine H, et al. Postoperative pneumonia after major lung resection. *Am J Respir Crit Care Med* 2006; **173**: 1161–9
 55. Schussler O, Dermine H, Alifano M, et al. Should we change antibiotic prophylaxis for lung surgery? Postoperative pneumonia is the critical issue. *Ann Thorac Surg* 2008; **86**: 1727–33
 56. Bagyi K, Haczkua A, Marton I, et al. Role of pathogenic oral flora in postoperative pneumonia following brain surgery. *BMC Infect Dis* 2009; **9**: 104
 57. Bouza E, Perez MJ, Munoz P, Rincon C, Barrio JM, Hortal J. Continuous aspiration of subglottic secretions in the prevention of ventilator-associated pneumonia in the postoperative period of major heart surgery. *Chest* 2008; **134**: 938–46
 58. Jiao J, Wang M, Zhang J, Shen K, Liao X, Zhou X. Procalcitonin as a diagnostic marker of ventilator-associated pneumonia in cardiac surgery patients. *Exp Ther Med* 2015; **9**: 1051–7
 59. Nicolosi LN, del Carmen Rubio M, Martinez CD, González NN, Cruz ME. Effect of oral hygiene and 0.12% chlorhexidine gluconate oral rinse in preventing ventilator-associated pneumonia after cardiovascular surgery. *Respir Care* 2014; **59**: 504–9
 60. Pelosi P, Barassi A, Severgnini P, et al. Prognostic role of clinical and laboratory criteria to identify early ventilator-associated pneumonia in brain injury. *Chest* 2008; **134**: 101–8
 61. Perkins GD, Gates S, Park D, et al. The beta agonist lung injury trial prevention. *Am J Respir Crit Care Med* 2014; **189**: 674–83
 62. Canadian Orthopaedic Trauma Society. Reamed versus unreamed intramedullary nailing of the femur: comparison of the rate of ARDS in multiple injured patients. *J Orthop Trauma* 2006; **20**: 384–7
 63. Force ADT, Ranieri VM, Rubenfeld GD, et al. Acute respiratory distress syndrome: the Berlin definition. *JAMA* 2012; **307**: 2526–33
 64. Camp SL, Stamou SC, Stiegel RM, et al. Quality improvement program increases early tracheal extubation rate and decreases pulmonary complications and resource utilization after cardiac surgery. *J Card Surg* 2009; **24**: 414–23
 65. Squadrone V, Coxa M, Cerutti E, et al. Continuous positive airway pressure for treatment of postoperative hypoxemia—a randomized controlled trial. *JAMA* 2005; **293**: 589–95