

Circulation

JOURNAL OF THE AMERICAN HEART ASSOCIATION



Mutation Type Is Not Clinically Useful in Predicting Prognosis in Hypertrophic Cardiomyopathy

Andrew P. Landstrom and Michael J. Ackerman

Circulation 2010;122:2441-2450

DOI: 10.1161/CIRCULATIONAHA.110.954446

Circulation is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75214

Copyright © 2010 American Heart Association. All rights reserved. Print ISSN: 0009-7322. Online ISSN: 1524-4539

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circ.ahajournals.org/cgi/content/full/122/23/2441>

Subscriptions: Information about subscribing to *Circulation* is online at
<http://circ.ahajournals.org/subscriptions/>

Permissions: Permissions & Rights Desk, Lippincott Williams & Wilkins, a division of Wolters Kluwer Health, 351 West Camden Street, Baltimore, MD 21202-2436. Phone: 410-528-4050. Fax: 410-528-8550. E-mail:
journalpermissions@lww.com

Reprints: Information about reprints can be found online at
<http://www.lww.com/reprints>

Is genotype clinically useful in predicting prognosis in hypertrophic cardiomyopathy?

Mutation Type Is Not Clinically Useful in Predicting Prognosis in Hypertrophic Cardiomyopathy

Andrew P. Landstrom, BS; Michael J. Ackerman, MD, PhD

Hypertrophic cardiomyopathy (HCM), or clinically unexplained hypertrophy of the heart, is a common genetic cardiovascular disorder marked by genetic and phenotypic heterogeneity. As the genetic mutations underlying the pathogenesis of this disease have been identified, investigators have attempted to link mutations to clearly defined alterations in survival in hopes of identifying prognostically relevant biomarkers of disease. While initial studies labeling particular *MYH7*-encoded beta myosin heavy chain and *TNNT2*-encoded cardiac troponin T mutations as “malignant” or “benign” raised hopes for mutation-specific risk stratification in HCM, a series of subsequent investigations identified mutations in families with contradictory disease phenotypes. Furthermore, subsequent proband-based cohort studies indicated that the clinical prognostic relevance of individual mutations labeled as “malignant” or “benign” in large referral centers is negligible. Herein, we seek to summarize the controversy and dispute the notion that mutation-specific risk stratification in HCM is possible at the present time. We provide evidence for clinicians and basic scientists alike to move beyond simple mutation descriptors to a more nuanced understanding of HCM mutations which fully captures the multi-factorial nature of HCM disease expression.

Response by Ho on p 2450

Over the last 2 decades, the genetic underpinnings of heritable cardiovascular disease have begun to be unveiled, starting with the discovery of rare pathogenic mutations that cause cardiomy-

opathies and cardiac channelopathies. The simple paradigm of Mendelian inheritance, while helpful in certain monogenic disease processes, is fundamentally incapable of explaining the entirety of how complex diseases express in the context of complex human physiology under the influence of a myriad of intrinsic and extrinsic variables. Our understanding of the intricate interplay between 1 such intrinsic variable, the genome, and the manifestation of disease was advanced significantly in the decoding of a handful of human’s genomes in February of 2001.^{1,2} Despite the decade of research these discoveries have initiated, clinicians and scientists alike are only beginning to appreciate the impact of this revolution on our understanding of health and disease. One disease, hypertrophic cardiomyopathy (HCM), typifies this struggle, as we seek to integrate the role of genetic alterations into the variation we see in this sudden-death-predisposing disease.

Defined by clinically unexplained hypertrophy of the ventricular walls and/or septum, HCM affects approximately 1 in 500 persons and is the most common inherited cardiovascular disease.³ HCM is the most common cause of sudden cardiac death (SCD) in young athletes and a significant cause of sudden death in the young in general.^{4,5} A heterogeneous disease, HCM demonstrates phenotypic variation in the degree of hypertrophy (none to extreme), fibrosis and myocyte disarray (none to extreme), left ventricular (LV) outflow tract obstruction (none to severe), morphological subtype (reverse curve, sigmoid, and apical-HCM, for example), associated symptoms (none to debilitating symptoms refrac-

The opinions expressed in this article are not necessarily those of the editors or of the American Heart Association.

From the Departments of Medicine, Pediatrics, and Molecular Pharmacology & Experimental Therapeutics, Divisions of Cardiovascular Diseases and Pediatric Cardiology, and the Windland Smith Rice Sudden Death Genomics Laboratory, Mayo Clinic, Rochester, Minn.

This article is Part II of a 2-part article. Part I appears on p 2430.

Correspondence to Michael J. Ackerman, MD, PhD, Windland Smith Rice Sudden Death Genomics Laboratory, Guggenheim 501, Mayo Clinic, Rochester, MN 55905. E-mail ackerman.michael@mayo.edu

(*Circulation*. 2010;122:2441-2450.)

© 2010 American Heart Association, Inc.

Circulation is available at <http://circ.ahajournals.org>

DOI: 10.1161/CIRCULATIONAHA.110.954446

tory to pharmacotherapy), and sudden death susceptibility (normal longevity to premature sudden death). The clinical progression is variable as well, with some patients remaining asymptomatic over their lifetime while others present during infancy with profound cardiac hypertrophy.

Genetic Basis of Hypertrophic Cardiomyopathy

The sentinel discovery of a genetic locus responsible for familial HCM was identified in 1989 by Jarcho and colleagues.⁶ Jarcho used linkage analysis of a large, multigenerational family to identify a portion of the long arm of chromosome 14, which cosegregated with incidence of disease. The following year, Geisterfer-Lowrance and investigators identified the first HCM-causative mutation in the *MYH7*-encoded beta myosin heavy chain.⁷ Over the past 20 years, HCM has been appreciated as principally an autosomal dominant disease with variable expressivity and penetrance, and hundreds of mutations found in dozens of genes encoding various sarcomeric/myofilament, Z-disc, and calcium (Ca^{2+})-handling proteins have been identified. These HCM-susceptibility genes are summarized in Table 1. In addition, the genetic basis of myocardial disease, so-called HCM phenocopies, which can masquerade as HCM, have also been elucidated (Table 2).

The majority of HCM is due to mutations in genes encoding the components of the cardiac sarcomere responsible for generating the molecular force of myocyte contraction. This sarcomeric basis of HCM is comprised of proteins of the thick myofilament (*MYH7*, *MYL2*-encoded regulatory myosin light chain, and *MYL3*-encoded essential myosin light chain^{8,9}), the intermediate myofilament (*MYBPC3*-encoded cardiac myosin binding protein C¹⁰), and the thin myofilament (*ACTC*-encoded actin,¹¹ *TPM1*-encoded α -tropomyosin,¹² *TNNT2*-encoded cardiac troponin T¹², *TNNI3*-encoded cardiac troponin I¹³, and *TNNC1*-encoded cardiac troponin C¹⁴). Finally, while a complete coding region interrogation through a large cohort of HCM probands has yet to be performed, a small number of mutations have been identified in the giant filament *TTN*-encoded titin. In this exceptionally large protein, which extends throughout half of the cardiac sarcomere, *TTN*-R740L has been identified in a single individual.¹⁵ In an independent study, Arimura and colleagues identified 2 additional mutations, *TTN*-R8500H and R8604Q, in a cohort of 384 HCM probands.¹⁶

Based on the replication and wide-spread acceptance of the role of these genes in the pathogenesis of HCM, this panel of genes, with the exception of *TTN*, has moved from the realm of research investigation to commercially/clinically available genetic tests for HCM. Depending on the cohort analyzed, the overall yield of HCM genetic testing, or its research equivalent, varies from 24% to 63% (24% Swedish,¹⁷ 34% German/Turkish,¹⁸ 38% United States,¹⁹ and 63% European²⁰ cohorts, respectively, among others). In comparison, the yield for the commercially available genetic test is purported to be 50% to 70% depending on the company. Although the frequency of mutations in these 9 genes depends on the cohort studied, mutations in *MYBPC3* and

Table 1. Summary of Hypertrophic Cardiomyopathy-Associated Genes

Gene	Locus	Protein	Frequency
Myofilament/Sarcomeric HCM			
Giant filament			
<i>TTN</i>	2q31	Titin	Rare
Thick filament			
<i>MYH7</i>	14q11.2-q12	β -Myosin heavy chain	25–40%
<i>MYH6</i>	14q11.2-q12	α -Myosin heavy chain	Rare
<i>MYL2</i>	12q23-q24.3	Regulatory myosin light chain	Rare
<i>MYL3</i>	3p21.2-p21.3	Essential myosin light chain	Rare
Intermediate filament			
<i>MYBPC3</i>	11p11.2	Cardiac myosin-binding protein C	25–40%
Thin filament			
<i>TNNT2</i>	1q32	Cardiac troponin T	3–5%
<i>TNNI3</i>	19p13.4	Cardiac troponin I	1–5%
<i>TPM1</i>	15q22.1	α -Tropomyosin	1–5%
<i>ACTC</i>	15q14	α -Cardiac actin	Rare
<i>TNNC1</i>	3p21.1	Cardiac troponin C	Rare
Z-disc HCM			
<i>ACTN2</i>	1q42-q43	α -Actinin 2	Rare
<i>ANKRD1</i>	10q23.31	Cardiac ankyrin repeat protein	Rare
<i>CSRFP3</i>	11p15.1	Muscle LIM protein	Rare
<i>LBD3</i>	10q22.2-q23.3	LIM binding domain 3	Rare
<i>MYOZ2</i>	4q26-q27	Myozenin 2	Rare
<i>TCAP</i>	17q12-q21.1	Telethonin	Rare
<i>VCL</i>	10q22.1-q23	Vinculin/metavinculin	Rare
Calcium-Handling HCM			
<i>CALR3</i>	19p13.11	Calreticulin 3	Rare
<i>CASQ2</i>	1p13.3-p11	Calsequestrin 2	Rare
<i>JPH2</i>	20q13.12	Junctophilin 2	Rare
<i>PLN</i>	6q22.1	Phospholamban	Rare
<i>RYR2</i>	1q42.1-q43	Ryanodine receptor 2	Rare

MYH7 comprise the majority of mutations among myofilament genes. Depending on the cohort, the prevalence of *MYBPC3*- and *MYH7*-HCM mutations varies among mutation-positive HCM probands, yet these 2 genes represent the most common genetic subtypes of HCM.^{17,18,20}

Although the clinically available genetic test represents a significant advance in the understanding of HCM pathogenesis, a significant proportion of the HCM population remains genotype-negative with no biomarker for, or mechanistic explanation of, their disease process. One possible explanation for the incomplete yield of mutation-positive HCM may

Table 2. Summary of HCM Phenocopy Genes

Syndrome	Gene	Locus	Protein
Barth syndrome/LVNC	<i>DTNA</i>	18q12	α -dystrobrevin
Barth syndrome/LVNC	<i>TAZ</i>	Xq28	Tafazzin (G4.5)
Danon's syndrome/WPW	<i>LAMP2</i>	Xq24	Lysosome-associated membrane protein 2
Fabry's disease	<i>GLA</i>	Xq22	α -galactosidase A
Forbes disease	<i>AGL</i>	1p21	Amylo-1,6-glucosidase
Friedrich's ataxia	<i>FXN</i>	9q13	Frataxin
Noonan's syndrome	<i>KRAS</i>	12p12.1	v-Ki-ras2 Kirsten rat sarcoma viral oncogene homolog
Noonan's syndrome	<i>SOS1</i>	2p22-p21	Son of sevenless homolog 1
Noonan's syndrome, LEOPARD syndrome	<i>PTPN11</i>	12q24.1	Protein tyrosine phosphatase, non-receptor type 11, SHP-2
Noonan's syndrome, LEOPARD syndrome	<i>RAF1</i>	3p25	V-RAF-1 murine leukemia viral oncogene homolog 1
Pompe's disease	<i>GAA</i>	17q25.2-q25.3	α -1,4-glucosidase deficiency
WPW	<i>PRKAG2</i>	7q35-q36.36	AMP-activated protein kinase

LVNC indicates left ventricular non-compaction; WPW, Wolff-Parkinson-White syndrome.

be due to limitations in the mutation detection methodology. As most genetic analyses utilize direct DNA sequencing of coding exons, possibly including a denaturing high-performance liquid chromatography intermediate platform, large genetic deletions or duplications as well as intronic or promoter variations are missed.²¹ Another potential explanation, which is certainly not mutually exclusive with the first, is that the disease causative pathogenic substrate lies elsewhere in the genome, thereby fostering the possibility for subsequent investigation into the causes of genotype-negative HCM to identify not only disease biomarkers but also novel cardiac physiology and disease pathophysiology.

The search for novel HCM genes moved away from principle myofilaments of the cardiac sarcomere to the genes that encode proteins comprising the adjacent cardiac Z-disc. Serving as mechanical integration site of the myofilaments, the network of Z-disc proteins anchor the thin filaments from adjacent cardiac sarcomeres and allow for transduction of sarcomere force generation. Furthermore, rather than being passive molecular tethers within the cardiocyte, Z-disc proteins serve as molecular platforms for signal transduction and initiation of several intracellular signaling cascades, particularly those responding to cardiac stretch.²² The first Z-disc mutations associated with HCM were described in *CSRP*-encoded muscle LIM protein²³ and *TCAP*-encoded telethonin.^{24,25} Subsequently, *LDB3*-encoded LIM domain binding 3, *ACTN2*-encoded alpha actinin 2,²⁶ *VCL*-encoded vinculin/meta-vinculin,^{27,28} *MYOZ2*-encoded myozenin 2,²⁹ and *ANLRD1*-encoded cardiac ankyrin repeat protein¹⁶ have been identified as rare causes of HCM.


Apart from sarcomeric- and Z-disc-HCM, independent studies have identified rare genetic mutations in genes encoding Ca^{2+} -handling or Ca^{2+} -regulatory proteins, including *JPH2*-encoded junctophilin 2,³⁰ *CALR3*-encoded calreticulin,³¹ and the previously mentioned Ca^{2+} -sensitive *TNNC1*¹⁴ in individuals with myofilament negative-HCM. All told, mutations in these genes explain only a small percentage of

HCM, yet may expand understanding of the role of Ca^{2+} in the pathogenesis of HCM.

Genesis of "Benign" and "Malignant" Mutations

The first studies exploring the possibility of mutations holding prognostic importance came on the heels of the identification of the first HCM mutation. Just 2 years after the first mutation was discovered, Watkins and investigators identified 4 families with mutations in *MYH7* associated with a marked reduction in survival on Kaplan-Meier analysis.³² Family members hosting either the MYH7-R403Q or R453C missense mutation had increased disease-related deaths and sudden deaths compared to those hosting a V606M mutation. The R403Q and R453C were designated "malignant" mutations, while the V606M was considered "benign." At the time, it was postulated that these initial malignant *MYH7* mutations might cluster around a "hot spot" on myosin, a loop that forms part of the binding cleft for actin, imparting a more dramatic functional impact on the protein than mutations seen elsewhere.³³ Anan et al identified a MYH7-F513C "benign" mutation as well as "malignant" MYH7-G716R and R719W.³⁴ In particular, the R719W mutation was associated with decreased survival when compared to a family hosting the "benign" F513C mutation.

Watkins and colleagues went on to identify mutations in additional genes, including 4 in *TNNT2* in 7 families which were associated with decreased life expectancy and a high incidence of SCD despite minimal cardiac hypertrophy.³⁵ Specifically, TNNT2-I79N, R92Q, a deletion of Q160 (delQ160), and a G to A transition in the 5' splice donor site of intron 15 (intervening sequence, IVS15+1 G>A) were found to have reduced survival compared with the "benign" MYH7-V606M yet similar to the previously described "malignant" MYH7-R403Q. Moolman and colleagues identified a TNNT2-R92W by linkage analysis in 2 pedigrees of familial HCM.³⁶ Among the 18 clinically examined individuals hosting this mutation, only 6 demonstrated LV wall



thickness ≥ 13 mm by echocardiographic evaluation. Despite this mild HCM phenotype, the authors show increased mortality, particularly in young males, with TNNT2-R92W positive individuals.

Together, these initial studies nucleated an archetype whereby an individual mutation type could, by itself, be considered a poor prognostic biomarker associated with increased risk of sudden death (MYH7-R403Q, R453C, G716R, and R719W; TNNT2-I79N, R92Q, R92W, delQ160, and IVS15+1 G>A) or a reassuring indicator of reduced mortality risk (MYH7-F513C, V606M).³⁷ Investigators quickly extended these studies beyond individual mutations and attempted to link the entire gene, independent of the specific type of mutation, to a particular genotype-specific clinical HCM manifestation. The following genotype-specific pronouncements emerged: MYH7-HCM and early onset with increased hypertrophy and susceptibility to SCD, MYBPC3-HCM and later onset of diagnosis with lower risk, TNNT2-HCM and youthful disease presentation, particularly among males, SCD-predisposition, and minimal hypertrophy, and ACTC-HCM and benign prognosis and apical LV hypertrophy morphology.^{38–40}

Nonreplication, Contradictions, and Confusion

Just as quickly as the field of prognostically relevant mutations expanded, a series of studies directly contradicted, or at least failed to replicate, many of these initial observations. In a family with the “malignant” MYH7-R403Q mutation, Fananapazir and Epstein found no SCD or disease-related deaths among 6 MYH7-R403Q-positive/HCM phenotype-positive individuals, yielding a 100% survival at 50 years of age⁴¹ compared with 9/44 SCD and 21/44 disease-related deaths previously observed.³² In addition to contradicting the “malignant” label of this mutation, investigators went on to contradict the “benign” nature of the MYH7-V606M mutation in a multigenerational family with HCM and prevalent SCD. Among 8 MYH7-V606M-positive/HCM phenotype-positive individuals, 4 succumbed to SCD between the ages of 15 and 27 years, yielding a 71% cumulative cardiac event rate at 50 years of age. In this way, Fananapazir and Epstein began to challenge the notion that clinical analysis of a single large pedigree was sufficient to designate a discrete mutation as intrinsically “malignant” or “benign.”³²

A second independent group led by Havnrup et al, also noted the “benign” MYH7-V606M in a family with HCM and a high incidence of SCD.⁴² In this multigenerational family demonstrating complete disease penetrance, 3 individuals, including a son of the index case, suffered youthful SCD with pronounced septal hypertrophy. Among the surviving 4 MYH7-V606M-positive/HCM phenotype-positive individuals, all were diagnosed before 18 years of age and 3 demonstrated either nonsustained ventricular tachycardia or atrial fibrillation.

These contradictions were not limited to MYH7-HCM. The “malignant” TNNT2-I79N mutation, which was first associated with the SCD of 4/9 mutation-positive individuals in the sentinel pedigree,³⁵ was identified by Menon et al in a

family with a mixed cardiomyopathic phenotype.⁴³ The mutation was identified in each of the 9 affected family members who held diagnoses of restrictive cardiomyopathy (N=2), dilated cardiomyopathy (N=2), HCM (N=4), and a mixed cardiomyopathy (N=1). Furthermore, although there was a high incidence of atrial tachyarrhythmias, there was no incidence of SCD. Taken together with previous studies, these results began to call into question the associations between individual mutations and patient survival. In addition to survival, contradictory reports began to emerge about the role of specific mutations influencing other aspects of HCM disease manifestation, such as the septal hypertrophy morphology.

In the sentinel study implicating *TNNI3* as an HCM-associated gene, Kimura et al identified a proband hosting a deletion of lysine 183 (TNNI3-delK183), which demonstrated apical-type HCM while his son, hosting the same mutation, had “typical” HCM.¹³ A later study by Kokado et al identified an identical finding in a seemingly independent group of 25 individuals from 7 families associated with a highly variable clinical expression of the deletion.⁴⁴ Indeed, among the 15 individuals demonstrating hypertrophy, only 1 had apical disease while 7 did not exhibit the apical morphology. Similarly, Brito and investigators identified 2 independent families with an identical novel MYH7-I263T mutation.⁴⁵ Despite hosting an identical mutation, family A (N=20 members) had increased LV hypertrophy when compared with family B (N=18 members, mean 23.4 mm versus 16 mm, respectively). Conversely, family B demonstrated reduced disease penetrance (100% versus 33%, respectively) with an increased propensity for SCD (0/20 versus 2/18, respectively) when compared with family A.

Taken together, these kindred studies illustrated the multifactorial nature of HCM in that the distinct mutation cannot be the sole factor that dictates clinical phenotype. Accordingly, one must be extremely cautious in assigning prognostic value to a specific mutation. Indeed, even in families carrying so-called “malignant” mutations, such as TNNT2-I79N and R92Q, there was significantly reduced disease penetrance with 3/9 (33%) and 7/32 (22%) mutation-positive individuals failing to show any clinical signs of HCM, respectively.⁴⁶ A major limitation of each of these studies is the small number of individuals available for analysis in even the largest, multigenerational family. Furthermore, confounding inherited factors that would be retained within a family, independent of the specific HCM-causative mutation, might greatly influence the ultimate expression of disease. What was not appreciated at the time, and may not be fully appreciated now, is the role of genomic and epigenetic influences on disease expression. Approaching the pathogenesis of HCM as a classical monogenic disease process, such as cystic fibrosis, does not fully capture the influence of other disease-modifying genetic variation outside the “gene of interest,” or biologically relevant factors inherited apart from the genome in an epigenetic fashion. These confounding factors might be 1 explanation for contradictory findings in families hosting

identical disease-associated mutations. Lastly, these studies do little to capture the epidemiological relevance of these “malignant” or “benign” mutations, and one is left wondering what the overall clinical impact of these mutations might be, if any, among the HCM population.

A Proband-Based Cohort Approach to Validation

In an attempt to move beyond kindred-based investigations of HCM mutation prognostic relevance and to identify “benign” and “malignant” mutations outside of the confounding influences of the genetic milieu of the family, a proband-based cohort approach was undertaken. Van Driest and colleagues identified 5 probands (1.7%), out of a cohort of nearly 300 consecutive index HCM cases, which hosted 1 of 2 “benign” HCM mutations: MYH7-R719Q or L908V.⁴⁷ Initial investigations by Consevage et al described the R719Q mutation in a large Hispanic family with no history of arrhythmia or SCD among 64 “at risk” family members.⁴⁸ Furthermore, Epstein et al initially identified the L908V mutation in a family with mild HCM with only 2 (4%) of 46 individuals <55 years of age experiencing SCD and only 5 (26%) of 19 individuals ≤30 years of age with a maximal LV thickness >12 mm.⁴⁹ In contrast, each of the “benign” mutation probands identified by Van Driest et al demonstrated significant clinical disease with an average age at diagnosis <30 years and LV outflow tract obstruction requiring surgical myectomy, with 4 necessitating β- and Ca²⁺ channel-blockade post myectomy. Three of the individuals had a positive family history of SCD, although 1 required cardiac transplant in the second decade of life because of end-stage HCM. In each of these cases, genetic counseling pertaining to the identification of each patient’s “benign” mutation would have been sorely amiss.

A follow-up study by Ackerman et al sought to determine the prevalence of purported “malignant” mutations in this cohort of clinically robust HCM probands.⁵⁰ Despite the large size of the cohort, only 3 probands (1%) of the 293 hosted the “malignant” MYH7-R453C, G716R, and TNNT2-R92W, mutations, leaving 98% of the 95 patients within the cohort with a positive family history of HCM, 99% of the 69 patients with family history of SCD, 92% of the 25 patients treated with an implantable cardioverter-defibrillator (ICD), and 89% of the patients with extreme (>30 mm) hypertrophy without a known “malignant” mutation. Furthermore, the TNNT2-R92W mutation proband was identified in a 24-year-old woman who had no family history of SCD despite a mutation-positive family. While limited by an incomplete picture of all possible “malignant” mutations, this study suggests that even if all “malignant” mutations identified in the literature were indeed deleterious to survival, presence of 1 of these mutations accounts for a minute fraction of those patients with phenotypically severe HCM.

Since these initial studies, we have grown the proband-based cohort to a cohort of 1064 cases (≈60% male) diagnosed with HCM at 44.4±18.6 (standard deviation) years with a mean ventricular septal thickness of 20.9±5.9 mm and a mean resting




Figure 1. Pie chart of the proportion of our 1064-proband HCM cohort that hosted literature-described “malignant” (black fill) and “benign” mutations (gray fill). Clinical characteristics of these patients, such as whether the proband underwent myectomy, are noted. FH indicates family history; SCD, sudden cardiac death; ICD, implantable cardioverter-defibrillator.

left ventricular outflow tract gradient of 43.7±43.5 mm Hg. A search of the literature to identify additional “prognostically relevant” HCM mutations not otherwise mentioned in earlier studies listed herein, identified the “malignant” MYH7-V406M,⁵¹ -R723G,⁵² MYL2- R58Q,^{20,53,54} -D166V,^{20,54} TNNT2-R94L,⁵⁵ -A104L,⁵⁶ homozygous -S179F,⁵⁷ and the “benign” MYH7-R663H,⁵¹ MYL2-E22K,⁵³ and heterozygous TNNT2-S179F.⁵⁷ Within this expanded cohort, the 16 previously annotated “malignant” mutations were identified in 8/1064 (0.75%) patients whereas those 10 mutations previously annotated as “benign” were seen in 28 patients (2.63%). As seen among the initial cohort of unrelated patients, many of the owners of a so-called “benign” mutation manifested a severe HCM phenotype and were part of families with SCD-predisposition (Figure 1).

One newly identified host of the “benign” MYH7-V606M mutation was diagnosed with HCM after an aborted SCD attempt and was found to be in ventricular fibrillation prior to resuscitation. A second proband hosting MYH7-V606M suffered the loss of his father from an HCM-related SCD at the age of 40 years as well as the SCD of a first cousin at the age of 10 years. A newly identified proband hosting MYH7-R719Q reported a significant family history of SCD-predisposition with SCD of a nephew at age 10 years, 4 first-degree relatives requiring ICD therapy, and a brother requiring cardiac transplant. In the context of previous studies that cast doubt on the prognostic relevance of individual mutations, this study further supports the conclusion that other factors, which play into the heterogeneous expression of HCM, easily overshadow the effects of specific mutation type or the gene that is mutated. These findings, as well as a summary of all studies directly contradicting the traditional label of “malignant” and “benign,” are summarized in Table 3.

Van Driest and colleagues further challenged the notion that mutations in certain genes, by the nature of their genetic location, might be associated with a particular clinical phenotype.⁵⁸ Investigators dissected a proband-based cohort of nearly 400 consecutive individuals with HCM based on genotype, and were unable to discern a clearly different clinical phenotype

Table 3. Summary of Contradictory Follow Up Studies

Mutation	Family/ Cohort	Phenotype	Reference
MYH7	
R403Q	Family	No sudden death in 6/6 individuals	41
TNNT2	
I79N	Family	No sudden death in 9/9 individuals	43
R92W	Cohort	No family hx of SCD	50
"Benign"			
MYH7	
V606M	Family	SCD <30 years of age in 4/8 individuals	41
V606M	Family	Youthful SCD in three individuals	42
V606M	Cohort	Aborted SCD, family hx of SCD	Mayo
R663H	Cohort	Aborted SCD, family hx of SCD	Mayo
R719Q	Cohort	Family hx of SCD, transplant	47
R719Q	Cohort	Family hx of SCD, family hx of transplant	Mayo
L908V	Cohort	Family hx of SCD	47
L908V	Cohort	Family hx of SCD	Mayo

Family/cohort indicates the type of follow-up study conducted; phenotype, pertinent clinical characteristics of kindred within the family or probands within the cohort studied; hx, history; SCD, sudden cardiac death/arrest; Mayo, results from analysis of the unpublished 1064 Mayo Clinic HCM probands.

among unrelated patients with thick filament (*MYH7*, *MYL2*, and *MYL3*), intermediate filament (*MYBPC3*), and thin filament (*ACTC*, *TPM1*, *TNNT2*, and *TNNI3*) HCM mutations. Indeed, among these 3 myofilament-based subtypes, there were no statistical differences in age at diagnosis, the number presenting with cardiac symptoms, family history of HCM or SCD, LV wall thickness, proportion with severe hypertrophy ≥ 30 mm, LV outflow tract gradient, and the proportion receiving myectomy, pacemaker, or ICD implantation.

For example, although MYBPC-HCM has been correlated with benign disease diagnosed late in life compared with MYH7-HCM, which was associated with extensive hypertrophy, Van Driest identified no statistical difference in age at diagnosis (MYH7, 33.0 ± 17 years versus MYBPC, 37.6 ± 15) or LV wall thickness (23.5 ± 7 mm versus 22.5 ± 5 mm, respectively, Figure 2). In addition to being unable to replicate many of the genotype-phenotype correlations that had been identified from multigenerational families, this study also highlighted that the 2 most common genetic subtypes of HCM, MYBPC-, and MYH7-HCM, were clinically similar.

A Rose by Any Other Name

Although it seems clear that a particular HCM-causative mutation is not inherently "malignant" or "benign," there is still a clear diagnostic role for HCM genetic testing in

identifying occult disease. Identification of a disease-associated mutation in a proband with HCM provides the genetic biomarker for the systematic genetic evaluation of his/her offspring, siblings, parents, and more distant relatives that do not demonstrate clinically apparent disease. Furthermore, a positive genetic test, independent of prognostic value of the discrete mutation, allows for risk stratification of family members based on the presence of this biomarker including: 1) close surveillance of the genotype-positive, preclinical individual and 2) casual observation or dismissal of the genotype-negative/phenotype-negative relative and his/her future progeny.⁵⁹ In this way, identification of a "benign" mutation in a family with HCM should not exclude close follow up of all mutation-positive individuals, nor should identification of a "benign" mutation solely dictate clinical management of a patient with HCM. Indeed, just as Juliet pondered "What's in a name? That which we call a rose by any other name would smell as sweet," in William Shakespeare's *Romeo and Juliet*, HCM mutations should be regarded as such without consideration for old descriptors.

Recent Advances in the Prognostic Role of Genotyping

Although significant controversy surrounds the notion of mutation-specific and genotype-specific risk stratification, recent evidence has emerged revealing that unrelated patients with a positive HCM genetic test have a distinctively more severe phenotype than patients with a negative HCM genetic test. In the Mayo Clinic study, compared with patients with a negative genetic test, patients with a positive genetic test were younger at diagnosis, demonstrated greater hypertrophy, were more likely to have a positive family history of HCM, and were more likely to receive an ICD from their HCM specialist even though the cardiologists were blinded to the patient's genetic status.¹⁹ Furthermore, a positive genetic test conferred a hazard ratio of 4.3 (1.5% to 12.5%, 95% confidence interval which was greater than that of age, degree of outflow tract obstruction, and presence of atrial fibrillation. Subsequently, these same distinguishing features between patients with a positive genetic test and those with a negative genetic test were seen in an independent cohort of patients with HCM from Italy. In this study, Olivotto and colleagues demonstrated a marked difference in outcome among those with a positive myofilament genetic test (Figure 3).⁶⁰ In this collection of 203 index cases, the 126 patients (62%) with a positive HCM genetic test had an increased combined risk of cardiovascular death, nonfatal stroke, or progression to New York Heart Association class III or IV symptoms compared with those with a negative genetic test. Furthermore, myofilament-positive individuals demonstrated a greater probability of severe LV systolic (ejection fraction $< 50\%$) and diastolic (restrictive filling pattern) dysfunction. These studies represent the first time a "positive genetic test" might be associated with adverse outcome in patients with HCM.




Figure 2. Drawing of the proteins that comprise the cardiac myofibrils include *MYBPC3*-encoded myosin binding protein C and *MYH7* encoded β -myosin heavy chain. Clinical characteristics for patients hosting *MYBPC3* and *MYH7* mutations are given.⁵⁸ Drawing adapted from Spirito et al.³⁷ Dx indicates age at diagnosis in years; LVWT, left ventricular wall thickness; FH, family history; SCD, sudden cardiac death.

Thus, although mutation-specific risk stratification is not possible, genetic test-based risk stratification seems clinically informative. Still, the translation of this observation into a clinically meaningful and “actionable” biomarker for the patient with already clinically manifest HCM is unclear. Knowing the difference in natural history and the increased likelihood toward disease progression, should patients with a positive genetic test be seen more frequently, intervened on sooner, and if so, with what interventions? Perhaps, the greatest contribution of the HCM genetic test for the phenotypically positive host will be for the clinician to “relax”

somewhat for the patient with a negative HCM genetic test knowing that collectively, such HCM gene test negative-individuals exhibit a milder disease phenotype and are far less likely to progress. Perhaps HCM patients with a negative HCM genetic test will ultimately require less frequent evaluations and cardiac tests.

Conclusion

It is clear that HCM is a truly complex disease that can present at any age with variable hypertrophy and outflow tract obstruction, and it can progress in an innocuous fashion, or predispose

A

Characteristic	Positive	Negative	P-Value
Age at Dx (yr)	36±17	45±19	<0.001
MLVWT (mm)	23±7	21±6	0.002
FH of HCM	68%	59%	<0.001
ICD	25%	10%	<0.001

Characteristic	HR	95% CI	P-Value
Positive Test	4.3	1.5-12.5	0.008
Age (per yr)	1.03	1.01-1.06	0.017
LVOTO (≥30mmHg)	1.33	0.7-2.7	0.43
Atrial Fibrillation	1.67	0.7-3.8	0.22




Figure 3. A) Top, table of the associations between clinical characteristics and a positive and negative HCM genetic test. Variance measured as standard deviation. Dx indicates diagnosis; MLVWT, maximal left ventricular wall thickness; FH, family history; ICD, implantable cardioverter-defibrillator. Bottom, table of the hazard ratio and 95% confidence interval of a positive HCM genetic test compared with age, left ventricular outflow tract obstruction, and atrial fibrillation. Adapted from Van Driest et al.¹⁹ B) Kaplan-Meier analysis of the probability of cardiovascular (CV) death, nonfatal ischemic stroke, or progression to heart failure with a negative and positive HCM genetic test.

individuals to arrhythmia and SCD. This clinical heterogeneity is matched by the genotypic heterogeneity associated with the pathogenesis of this disease. While the mutated genes that serve as the molecular substrate for this disease are becoming increasingly understood, the mechanistic link between a HCM-susceptibility mutation and disease pathogenesis and expressivity remains a significant challenge to elucidate. In large part, this is due to the multiplicity of factors that can modulate disease phenotype beyond the discrete HCM-causative pathogenic substrate. These additional patient-specific factors, whether intrinsic or extrinsic to the myocardium, can cause the same genetic misspelling to ultimately lead to dramatically different phenotypic variations of HCM in different individuals. Until these factors are more clearly understood, and the explanation for the contradiction and nonreplication of many of the studies trying to link specific HCM mutations to a particular phenotype is understood, there is no prognostic utility of a specific mutation in isolation.

Sources of Funding


Funding was provided by the Mayo Clinic Windland Smith Rice Comprehensive Sudden Cardiac Death Program, "Alliance for Calmodulin Kinase II Signaling in Heart Disease," a Leducq Fondation program grant, and the National Institutes of Health IPO1HL094291.

Disclosures

Michael J. Ackerman is a consultant for Medtronic, St. Jude Medical, Inc., Boston Scientific, and PGxHealth, and he serves on the Scientific Advisory Board for PGxHealth.

References

- International Human Genome Sequencing Consortium. Initial sequencing and analysis of the human genome. *Nature*. 2001;409:860–921.
- Venter JC, Adams MD, Myers EW, et al. The sequence of the human genome. *Science*. 2001;291:1304–1351.
- Maron BJ, Gardin JM, Flack JM, Gidding SS, Kurosaki TT, Bild DE. Prevalence of hypertrophic cardiomyopathy in a general population of young adults: echocardiographic analysis of 4111 subjects in the CARDIA Study. Coronary Artery Risk Development in (Young) Adults. *Circulation*. 1995;92:785–789.
- Maron BJ, Roberts WC, McAllister HA, Rosing DR, Epstein SE. Sudden death in young athletes. *Circulation*. 1980;62:218–229.
- Maron BJ, Epstein SE, Roberts WC. Causes of sudden death in competitive athletes. *J Am Coll Cardiol*. 1986;7:204–214.
- Jarcho JA, McKenna W, Pare JA, Solomon SD, Holcombe RF, Dickie S, Levi T, Donis-Keller H, Seidman JG, Seidman CE. Mapping a gene for familial hypertrophic cardiomyopathy to chromosome 14q1. *N Engl J Med*. 1989;321:1372–1378.
- Geisterfer-Lowrance AA, Kass S, Tanigawa G, Vosberg H, McKenna W, Seidman CE, Seidman JG. A molecular basis for familial hypertrophic cardiomyopathy: a beta cardiac myosin heavy chain gene missense mutation. *Cell*. 1990;62:999–1006.
- Poetter K, Jiang H, Hassanzadeh S, Master SR, Chang A, Dalakas MC, Rayment I, Sellers JR, Fananapazir L, Epstein ND. Mutations in either the essential or regulatory light chains of myosin are associated with a rare myopathy in human heart and skeletal muscle. *Nat Genet*. 1996;13:63–69.
- Olson TM, Karst ML, Whitby FG, Driscoll DJ. Myosin light chain mutation causes autosomal recessive cardiomyopathy with mid-cavitary hypertrophy and restrictive physiology. *Circulation*. 2002;105:2337–2340.
- Watkins H, Conner D, Thierfelder L, Jarcho JA, MacRae C, McKenna WJ, Maron BJ, Seidman JG, Seidman CE. Mutations in the cardiac myosin binding protein-C gene on chromosome 11 cause familial hypertrophic cardiomyopathy. *Nat Genet*. 1995;11:434–437.
- Mogensen J, Klausen IC, Pedersen AK, Egeblad H, Bross P, Kruse TA, Gregersen N, Hansen PS, Daandrup U, Borglum AD. Alpha-cardiac actin is a novel disease gene in familial hypertrophic cardiomyopathy. *J Clin Invest*. 1999;103:R39–R43.
- Thierfelder L, Watkins H, MacRae C, Lamas R, McKenna W, Vosberg HP, Seidman JG, Seidman CE. Alpha-tropomyosin and cardiac troponin T mutations cause familial hypertrophic cardiomyopathy: a disease of the sarcomere. *Cell*. 1994;77:701–712.
- Kimura A, Harada H, Park JE, Nishi H, Satoh M, Takahashi M, Hiroi S, Sasaoka T, Ohbuchi N, Nakamura T, Koyanagi T, Hwang TH, Choo JA, Chung KS, Hasegawa A, Nagai R, Okazaki O, Nakamura H, Matsuzaki M, Sakamoto T, Toshima H, Koga Y, Imaizumi T, Sasazuki T. Mutations in the cardiac troponin I gene associated with hypertrophic cardiomyopathy. *Nat Genet*. 1997;16:379–382.
- Landstrom AP, Parvatiyar MS, Pinto JR, Marquardt ML, Bos JM, Tester DJ, Ommen SR, Potter JD, Ackerman MJ. Molecular and functional characterization of novel hypertrophic cardiomyopathy susceptibility mutations in TNNC1-encoded troponin C. *J Mol Cell Cardiol*. 2008;45:281–288.
- Satoh M, Takahashi M, Sakamoto T, Hiroe M, Marumo F, Kimura A. Structural analysis of the titin gene in hypertrophic cardiomyopathy: identification of a novel disease gene. *Biochem Biophys Res Commun*. 1999;262:411–417.
- Arimura T, Bos JM, Sato A, Kubo T, Okamoto H, Nishi H, Harada H, Koga Y, Moulik M, Doi YL, Towbin JA, Ackerman MJ, Kimura A. Cardiac ankyrin repeat protein gene (ANKRD1) mutations in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2009;54:334–342.
- Mörner S, Richard P, Kazzam E, Hellman U, Hainque B, Schwartz K, Waldenström A. Identification of the genotypes causing hypertrophic cardiomyopathy in northern Sweden. *J Mol Cell Cardiol*. 2003;35:841–849.
- Erdmann J, Daehmlow S, Wischke S, Senyuva M, Werner U, Raible J, Tanis N, Dyachenko S, Hummel M, Hetzer R, Regitz-Zagrosek V. Mutation spectrum in a large cohort of unrelated consecutive patients with hypertrophic cardiomyopathy. *Clin Genet*. 2003;64:339–349.
- Van Driest SL, Ommen SR, Tajik AJ, Gersh BJ, Ackerman MJ. Yield of genetic testing in hypertrophic cardiomyopathy. *Mayo Clin Proc*. 2005;80:739–744.
- Richard P, Charron P, Carrier L, Ledeuil C, Cheav T, Pichereau C, Benaiche A, Isnard R, Dubourg O, Burban M, Gueffet JP, Millaire A, Desnos M, Schwartz K, Hainque B, Komajda M. Hypertrophic cardiomyopathy: distribution of disease genes, spectrum of mutations, and implications for a molecular diagnosis strategy. *Circulation*. 2003;107:2227–2232.
- Tester D, Will M, MJ A. Mutation detection in congenital long QT syndrome. In: Wang Q, ed. *Methods in Molecular Medicine*. Vol 128. Totowa, NJ: Humana Press Inc.; 2006:181–207.
- Hoshijima M. Mechanical stress-strain sensors embedded in cardiac cytoskeleton: Z disk, titin, and associated structures. *Am J Physiol Heart Circ Physiol*. 2006;290:H1313–H1325.
- Geier C, Perrot A, Ozcelik C, Binner P, Counsell D, Hoffmann K, Pilz B, Martiniak Y, Gehmlich K, van der Ven PF, Fürst DO, Vorwald A, von Hohenberg E, Nürnberg P, Scheffold T, Dietz R, Osterziel KJ. Mutations in the human muscle LIM protein gene in families with hypertrophic cardiomyopathy. *Circulation*. 2003;107:1390–1395.
- Hayashi T, Arimura T, Itoh-Satoh M, Ueda K, Hohda S, Inagaki N, Takahashi M, Hori H, Yasunami M, Nishi H, Koga Y, Nakamura H, Matsuzaki M, Choi BY, Bae SW, You CW, Han KH, Park JE, Knoll R, Hoshijima M, Chien KR, Kimura A. Tcap gene mutations in hypertrophic cardiomyopathy and dilated cardiomyopathy. *J Am Coll Cardiol*. 2004;44:2192–2201.
- Bos JM, Poley RN, Ny M, Tester DJ, Xu X, Vatta M, Towbin JA, Gersh BJ, Ommen SR, Ackerman MJ. Genotype-phenotype relationships involving hypertrophic cardiomyopathy-associated mutations in titin, muscle LIM protein, and telethonin. *Mol Genet Metab*. 2006;88:78–85.
- Theis JL, Bos JM, Bartleson VB, Will ML, Binder J, Vatta M, Towbin JA, Gersh BJ, Ommen SR, Ackerman MJ. Echocardiographic-determined septal morphology in Z-disc hypertrophic cardiomyopathy. *Biochem Biophys Res Commun*. 2006;351:896–902.
- Vasile VC, Ommen SR, Edwards WD, Ackerman MJ. A missense mutation in a ubiquitously expressed protein, vinculin, confers susceptibility to hypertrophic cardiomyopathy. *Biochem Biophys Res Commun*. 2006;345:998–1003.

- 
28. Vasile VC, Will ML, Ommen SR, Edwards WD, Olson TM, Ackerman MJ. Identification of a metavinculin missense mutation, R975W, associated with both hypertrophic and dilated cardiomyopathy. *Mol Genet Metab*. 2006;87:169–174.
29. Osio A, Tan L, Chen SN, Lombardi R, Nagueh SF, Shete S, Roberts R, Willerson JT, Marian AJ. Myozenin 2 is a novel gene for human hypertrophic cardiomyopathy. *Circ Res*. 2007;100:766–768.
30. Landstrom AP, Weisleder N, Batalden KB, Martijn Bos J, Tester DJ, Ommen SR, Wehrens XHT, Claycomb WC, Ko J-K, Hwang M, Pan Z, Ma J, Ackerman MJ. Mutations in JPH2-encoded junctophilin-2 associated with hypertrophic cardiomyopathy in humans. *J Mol Cell Cardiol*. 2007;42:1026–1035.
31. Chiu C, Tebo M, Ingles J, Yeates L, Arthur JW, Lind JM, Semsarian C. Genetic screening of calcium regulation genes in familial hypertrophic cardiomyopathy. *J Mol Cell Cardiol*. 2007;43:337–343.
32. Watkins H, Rosenzweig A, Hwang DS, Levi T, McKenna W, Seidman CE, Seidman JG. Characteristics and prognostic implications of myosin missense mutations in familial hypertrophic cardiomyopathy. *N Engl J Med*. 1992;326:1108–1114.
33. Watkins H, Seidman JG, Seidman CE. Familial hypertrophic cardiomyopathy: a genetic model of cardiac hypertrophy. *Hum Mol Genet*. 1995;4:1721–1727.
34. Anan R, Greve G, Thierfelder L, Watkins H, McKenna WJ, Solomon S, Vecchio C, Shono H, Nakao S, Tanaka H, Mares AJ, Towbin JA, Spirito P, Roberts R, Seidman JG, Seidman CE. Prognostic implications of novel beta cardiac myosin heavy chain gene mutations that cause familial hypertrophic cardiomyopathy. *J Clin Invest*. 1994;93:280–285.
35. Watkins H, McKenna WJ, Thierfelder L, Suk HJ, Anan R, O'Donoghue A, Spirito P, Matsumori A, Moravec CS, Seidman JG, Seidman CE. Mutations in the genes for cardiac troponin T and alpha-tropomyosin in hypertrophic cardiomyopathy. *N Engl J Med*. 1995;332:1058–1064.
36. Moolman JC, Corfield VA, Posen B, Ngumbela K, Seidman C, Brink PA, Watkins H. Sudden death due to troponin T mutations. *J Am Coll Cardiol*. 1997;29:549–555.
37. Spirito P, Seidman CE, McKenna WJ, Maron BJ. The management of hypertrophic cardiomyopathy. *N Engl J Med*. 1997;336:775–785.
38. Arad M, Penas-Lado M, Monserrat L, Maron BJ, Sherrid M, Ho CY, Barr S, Karim A, Olson TM, Kamisago M, Seidman JG, Seidman CE. Gene mutations in apical hypertrophic cardiomyopathy. *Circulation*. 2005;112:2805–2811.
39. Alcalai R, Seidman JG, Seidman CE. Genetic basis of hypertrophic cardiomyopathy: from bench to the clinics. *J Cardiovasc Electrophysiol*. 2008;19:104–110.
40. Varnava AM, Elliott PM, Baboonian C, Davison F, Davies MJ, McKenna WJ. Hypertrophic cardiomyopathy: histopathological features of sudden death in cardiac troponin T disease. *Circulation*. 2001;104:1380–1384.
41. Fananapazir L, Epstein ND. Genotype-phenotype correlations in hypertrophic cardiomyopathy. Insights provided by comparisons of kindreds with distinct and identical beta-myosin heavy chain gene mutations. *Circulation*. 1994;89:22–32.
42. Havndrup O, Bundgaard H, Andersen PS, Larsen LA, Vuust J, Kjeldsen K, Christiansen M. The Val606Met mutation in the cardiac beta-myosin heavy chain gene in patients with familial hypertrophic cardiomyopathy is associated with a high risk of sudden death at young age. *Am J Cardiol*. 2001;87:1315–1317.
43. Menon S, Michels V, Pellikka P, Ballew J, Karst M, Herron K, Nelson S, Rodeheffer R, Olson T. Cardiac troponin T mutation in familial cardiomyopathy with variable remodeling and restrictive physiology. *Clin Genet*. 2008;74:445–454.
44. Kokado H, Shimizu M, Yoshio H, Ino H, Okeie K, Emoto Y, Matsuyama T, Yamaguchi M, Yasuda T, Fujino N, Ito H, Mabuchi H. Clinical features of hypertrophic cardiomyopathy caused by a Lys183 deletion mutation in the cardiac troponin I gene. *Circulation*. 2000;102:663–669.
45. Brito D, Richard P, Isnard R, Pipa J, Komajda M, Madeira H. Familial hypertrophic cardiomyopathy: the same mutation, different prognosis. Comparison of two families with a long follow-up. *Rev Port Cardiol*. 2003;22:1445–1461.
46. Watkins H, McKenna WJ, Thierfelder L, Suk HJ, Anan R, O'Donoghue A, Spirito P, Matsumori A, Moravec CS, Seidman JG. Mutations in the genes for cardiac troponin T and alpha-tropomyosin in hypertrophic cardiomyopathy. *N Engl J Med*. 1995;332:1058–1064.
47. Van Driest SV, Ackerman MJ, Ommen SR, Shakur R, Will ML, Nishimura RA, Tajik AJ, Gersh BJ. Prevalence and severity of “benign” mutations in the beta myosin heavy chain, cardiac troponin-T, and alpha tropomyosin genes in hypertrophic cardiomyopathy. *Circulation*. 2002;106:3085–3090.
48. Consevage MW, Salada GC, Baylen BG, Ladda RL, Rogan PK. A new missense mutation, Arg719Gln, in the beta-cardiac heavy chain myosin gene of patients with familial hypertrophic cardiomyopathy. *Hum Mol Genet*. 1994;3:1025–1026.
49. Epstein ND, Cohn GM, Cyran F, Fananapazir L. Differences in clinical expression of hypertrophic cardiomyopathy associated with two distinct mutations in the beta-myosin heavy chain gene. A 908Leu—Val mutation and a 403Arg—Gln mutation. *Circulation*. 1992;86:345–352.
50. Ackerman MJ, Van Driest SV, Ommen SR, Will ML, Nishimura RA, Tajik AJ, Gersh BJ. Prevalence and age-dependence of malignant mutations in the beta-myosin heavy chain and troponin T gene in hypertrophic cardiomyopathy: a comprehensive outpatient perspective. *J Am Coll Cardiol*. 2002;39:2042–2048.
51. Greber-Platzer S, Marx M, Fleischmann C, Suppan C, Dobner M, Wimmer M. Beta-myosin heavy chain gene mutations and hypertrophic cardiomyopathy in Austrian children. *J Mol Cell Cardiol*. 2001;33:141–148.
52. Enjuto M, Francino A, Navarro-López F, Viles D, Paré J-C, Ballesta AM. Malignant hypertrophic cardiomyopathy caused by the Arg723Gly mutation in beta-myosin heavy chain gene. *J Mol Cell Cardiol*. 2000;32:2307–2313.
53. Kabaeva ZT, Perrot A, Wolter B, Dietz R, Cardim N, Correia JM, Schulte HD, Aldashev AA, Mirrakhimov MM, Osterziel KJ. Systematic analysis of the regulatory and essential myosin light chain genes: genetic variants and mutations in hypertrophic cardiomyopathy. *Eur J Hum Genet*. 2002;10:741–748.
54. Kerrick WG, Kazmierczak K, Xu Y, Wang Y, Szczesna-Cordary D. Malignant familial hypertrophic cardiomyopathy D166V mutation in the ventricular myosin regulatory light chain causes profound effects in skinned and intact papillary muscle fibers from transgenic mice. *FASEB J*. 2009;23:855–865.
55. Varnava A, Baboonian C, Davison F, de Cruz L, Elliott PM, Davies MJ, McKenna WJ. A new mutation of the cardiac troponin T gene causing familial hypertrophic cardiomyopathy without left ventricular hypertrophy. *Heart*. 1999;82:621–624.
56. Nakajima-Taniguchi C, Matsui H, Fujio Y, Nagata S, Kishimoto T, Yamauchi-Takahara K. Novel missense mutation in cardiac troponin T gene found in Japanese patient with hypertrophic cardiomyopathy. *J Mol Cell Cardiol*. 1997;29:839–843.
57. Ho CY, Lever HM, DeSanctis R, Farver CF, Seidman JG, Seidman CE. Homozygous mutation in cardiac troponin T: implications for hypertrophic cardiomyopathy. *Circulation*. 2000;102:1950–1955.
58. Van Driest SL, Vasile VC, Ommen SR, Will ML, Gersh BJ, Nishimura RA, Tajik AJ, Ackerman MJ. Myosin binding protein C mutations and compound heterozygosity in hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2004;44:1903–1910.
59. Bos JM, Towbin JA, Ackerman MJ. Diagnostic, prognostic, and therapeutic implications of genetic testing for hypertrophic cardiomyopathy. *J Am Coll Cardiol*. 2009;54:201–211.
60. Olivetto I, Girolami F, Ackerman MJ, Nistri S, Bos JM, Zachara E, Ommen SR, Theis JL, Vaubel RA, Re F, Armentano C, Poggesi C, Torricelli F, Cecchi F. Myofibrillar protein gene mutation screening and outcome of patients with hypertrophic cardiomyopathy. *Mayo Clin Proc*. 2008;83:630–638.



Response to Landstrom and Ackerman

Carolyn Y. Ho, MD

The crux of this debate is not the prognostic value of specific mutations, but rather the true purpose of genetic testing in HCM. Although their use of a small number of exceptions to invalidate potentially informative genotype-phenotype trends should be considered cautiously, I agree with the authors that prognosis cannot be determined solely on the basis of mutation identity with the current knowledge base. Indeed, genetic heterogeneity and inconsistent clinical expression hinder precise risk stratification in all inherited cardiovascular disorders. However, genetic testing has much more to offer beyond this narrow interpretation. The real utility of genetics is that it provides the best opportunity to understand disease and change medicine. Rather than categorizing all patients as having unexplained LVH, NOS, genotyping can tell you who has disease and what that disease is. This is crucial because prognosis is intimately tied to accurate diagnosis. As the authors acknowledge, patients without sarcomere mutations may have a more favorable outcome than those with mutations. In families with HCM, genotyping is the only way to identify susceptible relatives before disease is entrenched; those without mutations bear no risk.

Moreover, knowledge is evolving. Genotype-phenotype correlations may seem inconsistent now because traditional phenotypes (LVH, heart failure, sudden death) are late manifestations of disease. With better understanding of early manifestations, stronger correlations will emerge, mechanistic pathways will be identified, and disease-modifying treatments can be developed. I look forward to the day when we tell parents that although their child has a mutation, we can change their destiny.