



Correction to: Amitifadine, a triple reuptake inhibitor, reduces self-administration of the opiate remifentanil in rats

Edward D. Levin¹ · Corinne Wells¹ · Andrew Hawkey¹ · Zade Holloway¹ · Graham Blair¹ · Alexander Vierling¹ · Ashley Ko¹ · Caroline Pace¹ · John Modarres¹ · Anthony McKinney² · Amir H. Rezvani¹ · Jed E. Rose¹

Published online: 8 February 2021

© Springer-Verlag GmbH Germany, part of Springer Nature 2021

Correction to: Psychopharmacology

<https://doi.org/10.1007/s00213-020-05489-w>

Our article published in *Psychopharmacology* had a typographical error in the units of remifentanil infusion for self-administration.

The correct infusion dose of remifentanil is 0.3 µg/kg/infusion not 0.3 mg/kg/infusion.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s00213-020-05489-w>

✉ Edward D. Levin
edlevin@duke.edu

¹ Department of Psychiatry and Behavioral Sciences, Duke University School of Medicine, Durham, NC 27710, USA

² Ethismos Research, Inc, Cambridge, MA, USA