

Sparcl1/Hevin drives pathological pain through spinal cord astrocyte and NMDA receptor signaling

Gang Chen^{1*}, Jing Xu¹, Hao Luo¹, Xin Luo¹, Sandeep K. Singh^{2,3}, Juan J. Ramirez^{2,4}, Michael L. James⁵, Joseph P. Mathew⁵, Miles Berger⁵, Cagla Eroglu^{2,4,6,7*}, and Ru-Rong Ji^{1,2,4,7*}

¹Center for Translational Pain Medicine, Department of Anesthesiology, Duke University Medical Center, Durham, NC 27710, USA

²Department of Cell Biology, Duke University Medical Center, Durham, NC 27710, USA

³Department of Biochemistry and Molecular Biology, Virginia Commonwealth University School of Medicine, Richmond, VA 23298, USA

⁴Department of Neurobiology, Duke University Medical Center, Durham, NC 27710, USA

⁵Department of Anesthesiology, Duke University Medical Center, Durham, NC 27710, USA

⁶Howard Hughes Medical Institute, Duke University Medical Center, Durham, NC 27710, USA

⁷Duke Institute for Brain Sciences (DIBS), Durham, NC 27710, USA

Abbreviated title: Hevin regulates astrocyte signaling and pathological pain

Conflict of interest statement: The authors have declared that no conflict of interest exists.

***To whom correspondence may be addressed to:**

Gang Chen (chengang6626@ntu.edu.cn), Center for Basic Medical Research, Medical School of Nantong University, Nantong, Jiangsu Province, 19 QiXiu Rd, Nantong, JiangSu, China, Zip code: 226001, Tel: 86-513-85051776

ORCID: 0000-0003-3669-5687

Cagla Eroglu (cagla.eroglu@duke.edu), Department of Cell Biology, Howard Hughes Medical Institute, Duke University Medical Center, Durham, NC 27710, Tel: 919-6843605

ORCID, 0000-0002-7204-0218

Ru-Rong Ji (ru-rong.ji@duke.edu), Department of Anesthesiology, Duke University Medical Center, Durham, NC27710, Tel: 919-6849387, ORCID: 0000-0002-9355-3688

Abstract

Hevin/Sparg1 is an astrocyte-secreted protein and regulates synapse formation in the brain. Here we show that astrocytic hevin signaling plays a critical role in maintaining chronic pain. Compared to wild-type mice, hevin-null mice exhibited normal mechanical and heat sensitivity but reduced inflammatory pain. Interestingly, hevin-null mice have faster recovery than wild-type mice from neuropathic pain after nerve injury. Intrathecal injection of wild-type hevin was sufficient to induce persistent mechanical allodynia in naïve mice. In hevin-null mice with nerve injury, AAV-mediated re-expression of hevin in GFAP-expressing spinal cord astrocytes could reinstate neuropathic pain. Mechanistically, hevin is crucial for spinal cord NMDA receptor (NMDAR) signaling. Hevin potentiated NMDA currents mediated by the GluN2B-containing NMDARs. Furthermore, intrathecal injection of a neutralizing antibody against hevin alleviated acute and persistent inflammatory pain, postoperative pain, and neuropathic pain. Secreted hevin was detected in mouse cerebrospinal fluid (CSF) and nerve injury significantly increased CSF hevin abundance. Finally, neurosurgery caused rapid and substantial increases in SPARCL1/HEVIN levels in human CSF. Collectively, our findings support a critical role of hevin and astrocytes in the maintenance of chronic pain. Neutralizing of secreted hevin with monoclonal antibody may provide a new therapeutic strategy for treating acute and chronic pain and NMDAR-mediated neurodegeneration.

Keywords: Cerebrospinal fluid (CSF), GluN2B-NMDA receptors, nerve injury, neuro-glial interaction, spinal cord dorsal horn, synaptic plasticity

Introduction

Chronic pain, such as neuropathic pain is a major health problem worldwide and difficult to treat (1, 2). It is generally believed that maladaptive synaptic plasticity in the spinal cord and brain drives chronic pain (1, 3). Activation of NMDA receptors, especially NR2B subunit (GluN2B) plays a crucial role in injury-induced synaptic plasticity and pain pathogenesis (4, 5). Astrocytes are a major glial cell type in the central nervous system (CNS) and critical for maintaining CNS homeostasis, including physiological pain (6-8). Astrocytes make close contacts with synapses, and therefore, are positioned to regulate synapse formation and synaptic plasticity (9-12). During development astrocytes promote synapse formation and regulate synaptic connectivity in the CNS through secreted signals, such as thrombospondin-4 (TSP4) (9, 10, 13) or adhesion molecules, such as neuroligins and hepatocyte cell adhesion molecule (11, 14). Multiple types of neurological and neuropsychiatric disorders, such as Alzheimer's disease and chronic pain, may result from 'gliopathy' of astrocytes (15), leading to an inflammatory and pathological A1-like phenotype (16). Several lines of evidence support an essential role of astrocytes in neuropathic pain development and maintenance (15, 17-21). Following painful nerve injury or chemotherapy-induced neuropathy, astrogliosis is more prominent and persistent than microgliosis in the spinal cord dorsal horn (SDH) (15, 22). Several signaling molecules, such as connexin-43 (Cx43), and chemokines (CCL2 and CXCL1), MAP kinases (ERK/JNK), and JAK-STAT3 have been implicated in astrocyte signaling to maintain neuropathic pain (6, 18, 19, 23). Nerve injury upregulates TSP4 in spinal astrocytes and facilitates neuropathic pain via modulation of excitatory synaptic transmission (24). Gabapentin is a widely used drug for treating neuropathic pain through interaction with calcium channel $\alpha_2\delta_1$ subunit (Cav $\alpha_2\delta_1$), but interestingly, Cav $\alpha_2\delta_1$ has been identified as a thrombospondin receptor and regulates excitatory synaptogenesis (13). TSP4-induced pain can be blocked by gabapentin or Cav $\alpha_2\delta_1$ knockdown (25). Furthermore, a population of astrocytes in superficial SDH is involved in descending noradrenergic control of mechanical pain (26). However, it is not fully understood how astrocytes regulate synaptic plasticity in pathological pain.

Hevin, short for high endothelial venule protein, also known as SPARC-like 1 (SPARCL1) or synaptic cleft 1, SC1 (27), is a member of the SPARC family of glycoproteins that regulate cell-matrix interactions. Like TSP4, hevin is an astrocyte-secreted synaptogenic protein (9, 10), but its

role in chronic pain is unclear. Using a combination of in vitro and in vivo approaches, we demonstrated that hevin assembles Vesicular Glutamate Transporter 2 (VGLUT2) positive thalamocortical synapses by bridging neurexin-1alpha (Nrx1 α , a presynaptic component) and neuroligin-1B (NL1B, a postsynaptic component) (9). Neurexin-1alpha and NL1B do not interact with each other under normal conditions; but hevin can bridge these two adhesion proteins across the synapse, through interactions mapped to a specific synaptogenic domain (9). These interactions of hevin are critical for the formation and plasticity of thalamocortical synapses in the developing visual cortex via specific regulation of GluN2B-containing NMDARs (9). In this study, we investigated the role of hevin in physiological, inflammatory, neuropathic, and postoperative pain. We demonstrated that hevin is sufficient and required for inducing central sensitization (synaptic plasticity in the spinal cord pain circuit) and mechanical pain (tactile allodynia) via NMDAR/GluN2B signaling. Our data also show that hevin is secreted in mouse and human CSF under injury conditions, and neutralization of the secreted hevin protein with a monoclonal antibody effectively alleviates neuropathic pain. Furthermore, we demonstrate that re-expression of hevin in spinal cord astrocytes of knockout mice lacking hevin is sufficient to reinstate neuropathic pain.

Results

Hevin Is Distinctly Required in Physiological Pain, Inflammatory Pain, and Neuropathic Pain

We first set out to assess pain sensitivity in wild-type (WT) and hevin-knockout (KO) mice in physiological and pathological conditions. Compared to WT control, KO mice exhibited normal baseline mechanical sensitivity as measured in von Frey testing (Fig Figure 1A) and thermal sensitivity in Hargreaves test, hot plate, and tail immersion tests (Figure 1, B-D), suggesting that hevin is dispensable for physiological pain. Next, we examined formalin-induced acute inflammatory pain, which is typically divided into phase-1 (0-10 min) and phase-2 (10-45 min). Notably, phase-2 spontaneous pain is driven by NMDAR-mediated spinal neuron sensitization (central sensitization). We found that formalin-induced phase-2, but not phase-1 spontaneous pain was significantly reduced in KO mice (Figure 1E), supporting a role of hevin in central sensitization-mediated pain. Intraplantar carrageenan induced rapid and persistent inflammatory

pain that fully recovered in 3 days in WT mice; but hevin-KO mice showed a faster recovery of this inflammatory pain (Figure 1F). Chronic constriction injury (CCI) of the sciatic nerve induced neuropathic pain, manifested as mechanical allodynia, a reduction in paw withdrawal threshold (PWT), which was maintained at 28 d in WT mice. Strikingly, even though the induction phase of neuropathic pain (5-10 d) was not altered in KO mice compared to WT mice, the CCI-induced mechanical allodynia was fully recovered at 28 d in hevin-KO mice, (Figure 1G). We also measured CCI-induced ongoing pain (3 w after CCI) using a 2-chamber conditioned place preference test (28). The result showed that hevin-KO mice spent less time than WT mice in the analgesic clonidine-treated chamber (Figure 1H), indicating that nerve injury-induced ongoing pain maintenance is reduced in hevin-KO mice. These results show that hevin is required for driving inflammatory and neuropathic pain but dispensable for physiological pain.

Hevin Is Sufficient to Induce Mechanical Allodynia in WT Mice

Next, we evaluated whether administration of exogenous hevin, via intrathecal route to target spinal cord, in naïve mice would elicit mechanical pain. We also included a hevin mutant which cannot bridge Nr1a and NL1B thus cannot induce synapse formation (hevin- Δ DE, lacking a.a. 351-440, Supplemental Fig. 1) as a negative control (9). Intrathecal injection of purified WT hevin (10 μ g, i.t.) but not hevin- Δ DE (10 μ g, i.t.) induced robust and persistent mechanical allodynia in naïve male mice, this effect lasted for more than 3 days and recovered after 5 days (Figure 2A). We also examined whether hevin could exacerbate neuropathic pain. Intrathecal injections of WT but not hevin- Δ DE, given 12 d after CCI, further enhanced mechanical allodynia in male mice with nerve injury (Figure 2B). Moreover, intrathecal injection of purified WT hevin (10 μ g) significantly enhanced formalin-induced spontaneous pain in phase-2 (10-45 min) in male mice (Figure 2C, $P < 0.05$ vs. hevin- Δ DE), female mice (Figure 2D, $P < 0.05$ vs. hevin- Δ DE), and males and females combined (Figure 2E, $P < 0.05$ vs. hevin- Δ DE). To further examine the possible prolongation of formalin-induced pain by WT hevin, we quantified spontaneous pain in phase-2+ (45-60 min) and found significant increase in phase-2+ by WT hevin in male mice (Figure 2C, $P < 0.05$ vs. hevin- Δ DE), female mice (Figure 2D, $P < 0.05$ vs. hevin- Δ DE), and both sexes combined (Figure 2E, $P < 0.001$ vs. hevin- Δ DE). However, phase-1 pain behavior was not affected by intrathecal hevin- Δ DE (10 μ g) in both male and female mice (Figure 2, C-E). These results suggested that hevin is sufficient to induce mechanical allodynia in naïve mice, potentiate

central sensitization, and further enhance pain in neuropathic pain mice. Recent studies have shown sex dimorphism in spinal cord regulation of pain by glial cells and immune cells (29-31). We did not find sex differences in hevin-induced pain, as i.t. hevin potentiated formalin-induced pain (Figure 2, C-E) and induced robust mechanical allodynia in both males and females (Supplemental Figure 2A).

Re-expression of hevin In Spinal Astrocytes Reinstates Neuropathic Pain

To investigate the cellular locations of hevin in the spinal cord, we examined *Sparcl1* (mouse hevin gene) expression in the spinal cord dorsal horn (SDH) by RNA-scope in situ hybridization. We conducted double staining of *Sparcl1* and *Slc32a1* (encoding vesicular inhibitory amino acid transporter) to localize hevin in inhibitory neurons. In order to localize hevin in excitatory neurons, we also performed double staining in VGLUT2: Ai9 mice. We found that *Sparcl1* is partly expressed in VGLUT2+ excitatory neurons and *Slc32a1*+ inhibitory neurons (Figure 3A). Quantitative analysis revealed that *Slc32a1* is evenly distributed in excitatory and inhibitory neurons of SDH (Figure 3B).

Immunohistochemistry (IHC) revealed that hevin is expressed by the majority of astrocytes (GFAP⁺, Figure 4A) and some neurons (NeuN⁺, Supplemental Figure 3, A and B), but not by microglia (Iba1⁺, Supplemental Figure 3, C and D) in SDH. The specificity of the hevin antibody was validated by loss of hevin staining in SDH of hevin-KO mice (Figure 4B). Western blot result also confirmed hevin expression in WT-SDH, which was lost in SDH of KO mice (Supplemental Figure 3E). IHC also showed hevin expression in neurons and satellite glial cells of dorsal root ganglion (DRG) of WT mice, but this staining was absent in hevin KO mice (Supplemental Figure 3, F and G).

To determine a specific role of hevin from spinal astrocytes in neuropathic pain, we evaluated whether hevin re-expression in SDH astrocytes would reinstate neuropathic pain in hevin-KO mice (Figure 2G). To this end, we conducted SDH microinjection of adeno-associated viruses (AAV2/5.GFAP.Hevin-MycHis, termed as hevin-AAV, Supplemental Figure 1B) (9) to re-express hevin in spinal astrocytes in hevin-KO mice. We also included AAV2/5.GFAP.Hevin Δ DE-MycHis virus as a negative control (termed as hevin Δ DE-AAV, Supplemental Figure 1B). AAVs were injected to the ipsilateral SDH of KO mice either prior to CCI nerve injury (pretreatment, Figure 4, C and D) or after CCI (post-treatment, Figure 4, E and

F). The pre-treatment in naïve mice induced mild and transient reduction in PWT in both hevin-AAV and hevin Δ DE-AAV groups, due to spine surgery; but this PWT change fully recovered 5 days after the surgery (Figure 4D). Following CCI, mice treated with hevin Δ DE-AAV exhibited similar time course of PWT change as in hevin-KO mice (Figure 2G), and neuropathic pain began to recover on CCI-8d (Figure 3D). Strikingly, hevin-AAV-treated KO mice showed no sign of neuropathic pain recovery on CCI 8-12d, and mechanical allodynia was significantly prolonged compared to hevin Δ DE-AAV treated mice ($P < 0.05$, Figure 4D). Furthermore, post-treatment of hevin-AAV, given 2 days after CCI (Figure 3E), also significantly decreased PWT compared to the hevin Δ DE-AAV mice ($P < 0.05$), displaying prolonged neuropathic pain on CCI-26d (Figure 4F). Thus, both pre- and post-treatments of hevin-AAV in KO mice restore neuropathic pain by enhancing and sustaining mechanical allodynia after CCI.

We confirmed the specific expression of hevin in SDH astrocytes after the hevin-AAV injection, in the spinal cord sections of hevin-KO mice. We used antibodies against hevin and GFAP, as well as Myc (Figure 4, G and H), as hevin-AAV has a Myc-tag (Supplemental Figure 1B). At 24 d after the ipsilateral SDH hevin-AAV injection, we found many GFAP+ astrocytes in the ipsilateral SDH which were also labeled for hevin and Myc (Figure 4G), especially laminae I-III, a critical region for the transmission of mechanical pain (32, 33). High magnification images further revealed that hevin+ astrocytes also expressed Myc and GFAP in superficial SDH (Figure 4H). We did not observe hevin expression in the contralateral SDH of hevin-KO mice (Figure 4G). As previously reported (34), CCI caused a dramatic increase in the number of GFAP+ cells (astrogliosis) in the ipsilateral SDH. Collectively, these behavioral and histochemical data suggest that re-expression of hevin in spinal cord astrocytes is sufficient to reinstate neuropathic pain.

Hevin Regulates NMDA-Evoked Pain and NMDA Currents in SDH Neurons

Previously we found that hevin strongly enhances NR2B subunit-containing NMDA receptor function by increasing NMDA currents in autaptic neuron-only cultures (9). Thus, here we tested the function of spinal NMDARs in WT and hevin-KO mice using behavioral and electrophysiological approaches. Spinal injection of NMDA (3 nmol, i.t.) induced persistent mechanical allodynia in WT mice, which lasted for >2 weeks and recovered after 3 weeks (Figure 5A). The duration of NMDA-evoked mechanical allodynia was remarkably shortened in hevin-KO mice, showing a full recovery at 6 days post injection (Figure 5A).

We recorded NMDA-induced currents in spinal lamina IIo neurons via bath application of NMDA (100 μ M, 3 sec) to spinal cord slices in WT and hevin-KO mice. Compared to WT neurons, hevin-KO neurons had significantly smaller NMDA-elicited currents ($P < 0.05$, Figure 5B). Since GluN2A and GluN2B containing NMDARs may play different roles in synaptic transmission and synaptic plasticity (5), we further tested the blocking effects of GluN2A antagonist TCN201 (10 μ M) and GluN2B antagonist RO25 (10 μ M) on NMDA currents in spinal cord slices from WT and hevin-KO mice. RO25 produced a significant inhibition of NMDA currents in WT neurons (31%, $P < 0.05$) but only evoked mild inhibition (12%, $p > 0.05$) of NMDA currents in hevin-deficient neurons (Figure 5B). In contrast, TCN (10 μ M) produced significant and comparable inhibition of NMDA currents in both WT (26%, $P < 0.05$) and KO (26%, $P < 0.05$) neurons (Figure 5C). Therefore, we conclude that hevin deficiency primarily affects the GluN2B-mediated NMDA currents in lamina IIo neurons.

Next, we investigated whether perfusion of spinal cord slices with hevin is sufficient to enhance NMDA-evoked currents in WT mice. A brief exposure of SDH neurons to hevin (4 min, 100 ng/ml \approx 0.14 nM) elicited a significant increase in NMDA-induced currents (Before: 450.3 ± 75.5 pA; After: 625.9 ± 87.3 pA, $P < 0.05$, *SI Appendix*, Fig. S4a). Hevin-induced potentiation of NMDA currents was completely blocked by GluN2B antagonist RO25 (10 μ M, Figure 5D). In contrast, GluN2A antagonist TCN (10 μ M) produced a similar inhibition of NMDA currents in ACSF and hevin treated neurons (19% vs. 18% inhibition, $P < 0.05$, Figure 5E). Hevin still evoked significant increase in NMDA currents after TNC-201 treatment ($P < 0.05$, Figure 5E). However, hevin had no effects on AMPA-evoked currents in lamina IIo neurons (fig. S4b). We further assessed evoked excitatory postsynaptic currents (eEPSCs) using dorsal root stimulation. Hevin enhanced the NMDA receptor-mediated eEPSCs in lamina IIo neurons (Supplemental Figure 4C). Collectively, these data suggest that hevin is sufficient and required to regulate GluN2B/NMDAR-mediated currents in SDH neurons. This hypothesis was further supported by our behavioral observation that mechanical allodynia by intrathecal hevin was completely reversed by RO25 in both sexes (Supplemental Figure 2B). Consistent with a previous study (29), we did not find sex differences in NMDAR-mediated pain (Supplemental Figure 2B).

Hevin-Neutralizing Antibody Reduces Inflammatory, Postoperative and Neuropathic Pain in WT Mice

To examine if hevin is secreted from the mouse spinal cord, we measured hevin levels in the cerebrospinal fluid (CSF) using a mouse hevin-specific ELISA kit. We detected a high basal level of hevin secretion (~5 ng/ml) in the CSF of naïve mice (Figure 6A). To determine the role of secreted hevin in inflammatory and neuropathic pain, we tested a monoclonal antibody against mouse hevin which we previously characterized (9). The monoclonal antibody recognizes an epitope mapping to the amino acids 368–419 of hevin (anti-hevin 12:54, Supplemental Figure 1C) and this antibody blocks hevin's synaptogenic function, likely by interfering hevin's ability to bind Nrx1a and/or NL1B (9). Another hevin antibody recognizing a different epitope (anti-hevin 12:155, Supplemental Figure 1C) does not impact hevin's synaptogenic activity, which we used as our control. Intrathecal injection of anti-hevin 12:54 (10 µg) significantly reduced formalin-induced phase II pain, compared with anti-hevin 12:155 treated mice ($P < 0.05$, Figure 6B). We also tested these two antibodies in a chronic inflammatory pain model (induced by complete Freund's adjuvant, CFA, Figure 6C) and a postoperative pain (induced by plantar incision, Figure 6D). Intrathecal injection of anti-hevin 12:54 (10 µg) significantly reduced mechanical allodynia in the CFA model (Figure 6C, $P < 0.05$ at 1, 3, and 5 hours) and incision model (Figure 6D, $P < 0.001$ at 1, 3, and 5 hours), compared with anti-hevin 12:155 treated mice. Furthermore, intrathecal injection of anti-hevin 12:155 (10 µg), but not anti-hevin 12:54 (10 µg), rapidly (< 1 h) reversed mechanical allodynia for > 5 h in CCI mice, in both early-phase (7d after CCI, Figure 6E) and late-phase (21d after CCI, Figure 6F) of neuropathic pain. ELISA analysis showed that CCI resulted in significant increase in CSF hevin levels (Sham: 5.24 ± 0.57 ng/ml; CCI-14d: 8.32 ± 0.89 ng/ml, $P < 0.05$, Figure 6A). These results suggest that 1) hevin is secreted in physiological and pathological conditions and 2) targeting secreted hevin with a neutralizing antibody can potentially alleviate inflammatory and neuropathic pain.

Hevin Is Increased in Human CSF Following Intracranial Surgery

To explore the translational potential of these findings, next we assessed secreted Hevin levels in human CSF samples collected prior to and 12 h after intracranial surgery (e.g., otolaryngology procedures (35), using a human Hevin-specific ELISA kit. In the human CSF, we were able to detect a basal secretion of Hevin, ranging from 0.8-2.7 ng/ml (Figure 7A). Intracranial surgery resulted in a rapid and dramatic increase in CSF Hevin levels, ranging from 3.1-52.9 ng/ml at 12 h (Figure 7A). Strikingly, Hevin showed marked increases in all the CSF samples we analyzed,

ranging from 2.3-50.9 fold ($P < 0.01$, Figure 7B). We also measured total protein levels in these CSF samples and observed mild but significant increases, ranging from 1.3-4.3 fold ($P < 0.001$, Figure 7C). After normalization with respective protein changes, CSF-Hevin levels still exhibited a significant increase (9.0 fold, $P < 0.01$, Figure 7D). These results suggest that human neurosurgery selectively increased the CSF section of hevin, beyond the postoperative increase in CSF total protein levels.

Discussion

Despite recent progress in demonstrating “gliopathy” in the pathogenesis of pain (15) and the previous studies showing an essential role of spinal TSP4 and cortical TSP1 in neuropathic pain sensitization (24, 25, 36), our knowledge is still limited regarding the specific mediators secreted by astrocytes that can drive neuropathic pain. We employed both loss-of-function and gain-of-function approaches to demonstrate a critical role of astrocytic hevin in driving neuropathic pain. Our data showed that nerve injury-induced mechanical allodynia in the maintenance phase, but not the induction phase, is impaired after hevin deficiency, in further support of the notion that astrocytes are crucial for the maintenance of neuropathic pain (15). Nerve injury induces marked and long-lasting astrogliosis in SDH that is correlated with the time course of neuropathic pain (6, 18). It has been shown that hevin is expressed by reactive astrocytes in the brain (37). We demonstrate that re-expression of hevin in SDH reactive astrocytes in hevin-KO mice is sufficient to reinstate neuropathic pain after nerve injury. Furthermore, intrathecal injection of hevin produced persistent mechanical allodynia in WT mice of both sexes.

Mechanistically, we demonstrated that hevin induces central sensitization and mechanical pain through regulation of GluN2B containing NMDARs in SDH neurons, in agreement with our previous report that hevin is crucial for the formation of thalamocortical connections in the visual cortex via specific regulation of GluN2B during development (8). Thus, hevin-induced mechanical pain was completely blocked by the GluN2B antagonist RO25. As expected, hevin deficiency resulted in a significant reduction of NMDA currents in SDH neurons through adulthood. Interestingly, physiological pain in adult mice at the baseline is unaltered in hevin-KO mice, suggesting a specific contribution of NMDARs to synaptic plasticity in pathological pain (3).

Although we did not find significant change in AMPA-induced currents in SDH neurons of hevin-KO mice, a recent study shows that spinal hevin also mediates membrane trafficking of GluA1-containing AMPA receptors in remifentanyl-induced postoperative hyperalgesia (38). Our findings strongly suggest that hevin-induced synaptic plasticity not only occurs during development but also manifests in pathological and neurodegenerative conditions such as neuropathic pain.

Our data also showed that hevin is highly secreted in mouse CSF and this secretion is further increased after nerve injury. Notably, CSF-hevin changes are clinically relevant. We found a rapid and substantial increase of hevin in human CSF samples 12 hours after painful neurosurgical procedures (35). Neurosurgery/otolaryngology procedures are also associated with an increase in the CSF tau levels that have been implicated in the development of dementia (35). It is of great interest to examine CSF-hevin levels in a general surgery population. Thus, CSF hevin could potentially serve as a biomarker for pain following neurosurgery or general surgery, an important question for evaluation in future studies.

It is well-established that overactivation of GluN2B causes neurotoxicity (39). Thus, high levels of hevin production and secretion in the CNS is not only associated with pain but may also underlie surgery-induced neurocognitive injury, such as delirium (40) by producing GluN2B-mediated neurotoxicity. Future studies are warranted to evaluate the involvement of hevin in neurocognitive function under physiological and pathological conditions. Importantly, we have demonstrated that neutralizing secreted hevin with a monoclonal antibody could effectively alleviate inflammatory and neuropathic pain. Thus, targeting secreted hevin may offer new therapeutics to manage chronic pain, post-operative neurocognitive injury, and neurodegenerative diseases.

There are several limitations in this study. First, despite substantial increases of hevin in all the human CSF samples generated in the acute phase of intracranial surgery, the sample size of this study is small and it is difficult to associate increases in Hevin abundance with pain in patients. It is also possible that CSF-hevin increase is a general response to surgery or acute stress. Future studies are needed to collect more CSF samples at multiple time points of surgeries or other painful diseases to correlate hevin levels with acute pain and chronic pain. Second, hevin is produced not only by astrocytes but also by a subset of neurons in the spinal cord and DRG. It remains to be identified whether hevin in the CSF is mainly derived from astrocytes. Deletion of *Sparcl1* in different cell types using conditional knockout mice will help to address this issue. Finally, hevin

may regulate synaptic plasticity and pain via non-NMDA receptors, such as AMPA receptors (38). In addition to neuronal modulation, hevin may regulate microglia activation that has been strongly implicated in chronic pain development (41, 42). It will be of great interest to study hevin-mediated neuron-glial and glia-glial interactions in acute and chronic pain.

Materials and Methods

Reagents. We purchased formalin, carrageenan, complete Freund's adjuvant (CFA), NMDA from Sigma-Aldrich, TCN201 (GluN2A antagonist, Cat. No.4154), Ro25-6981 (GluN2B antagonist, Cat. No.1594) from Tocris, and recombinant hevin from RD system (Cat: 4547-SL). Purified recombinant hevin and hevin- Δ DE (lacking a.a. 351-440) proteins, anti-hevin monoclonal antibody 12:155, and anti-hevin monoclonal antibody 12:54 were from laboratory of Cagla Eroglu at Duke University Medical Center.

Mice

Hevin KO mice were from Eroglu's laboratory (9), and control SVE/129 mice (aged 8-12 weeks of both sexes) were obtained from Taconic Biosciences. These mouse lines were maintained at Duke University animal facility. We also used CD1 mice (Charles River, 8-10 weeks) of both sexes for comparing hevin's or anti-hevin's effects on mechanical pain. VGLUT2: Ai9 mice were made by crossing Ai9 mice (Jackson JAX:007909) with vGlut2 flox mice (JAX:033688) for RNAscope experiment. All animals were housed under a 12-hour light/dark cycle with food and water available ad libitum. Sample sizes were estimated based on our previous studies for similar types of behavioral and electrophysiological analyses (5, 19). Animals were randomly assigned to each experimental group.

Mouse models of pain, CSF collection, and intrathecal injection

To produce inflammatory pain, diluted formalin (5%, 20 μ l), carrageenan (1.5 %, 20 μ l) or CFA (20 μ l) was injected into the plantar surface of a hindpaw (5). Neuropathic pain was induced by chronic constriction injury (CCI) as previously published (19). In brief, after the left sciatic nerve was exposed, three ligatures (LOOK® 6-0 Silk, #SP102) were placed around the nerve proximal to the trifurcation with one millimeter between each ligature. The ligatures were loosely tied until a short flick of the ipsilateral hind limb was observed. Mice in the sham group received surgery identical to those described but without nerve ligation. Mouse cerebrospinal fluid (CSF) was collected from the cisterna magna 14 days after CCI (34). Plantar incision surgery was performed as previously described (43). A 5-mm longitudinal incision was made by #11 blade through the skin and fascia of the plantar foot, starting 2 mm from the proximal edge of the heel. For intrathecal

injection, spinal cord puncture was made with a 30G needle between the L5 and L6 level to deliver reagents (10 μ l) to the cerebral spinal fluid (34).

Spinal injection of AAV2/5 virus

As our previous report (9), full-length hevin cDNA was cloned into the pZac2.1 gfaABC1D-Cyto-GCaMP3 viral vector (Addgene plasmid #44331), between the NheI and NotI restriction sites, replacing the GCaMP3 sequence. AAV2/5 particles were packaged and synthesized by the Penn Vector Core facility (University of Pennsylvania). Intra-spinal injections were performed as described previously (44). In brief, Hevin KO mice were deeply anaesthetized by subcutaneous (s.c.) injection of ketamine (100 μ g/g) and xylazine (10 μ g/g). Paraspinal muscles around the left side of the interspace between Th13 and L1 vertebrae were removed, and the dura mater and the arachnoid membrane were carefully incised using the tip of a 30G needle to make a small window to allow a glass micropipette (diameter of 80 μ m) insert directly into the spinal cord dorsal horn (SDH). The microcapillary was inserted with AAV2/5 virus solution through the small window (approximately 500 μ m lateral from the midline) and inserted into the SDH (250 μ m in depth from the surface of the dorsal root entry zone). Each mouse was given single injection (0.7 μ l, 1×10^{12} GC/ml) of AAV5.GFAP.Hevin-MycHis.SV40 virus or AAV5.GFAP.HevinDDE-MycHis.SV40 virus.

Behavioral testing

Animals were habituated to the environment for at least 2 days before the testing. The room temperature and humidity remained stable for all experiments. All the behaviors were tested blindly. Key behavioral studies were repeated by different lab members. For testing spontaneous pain in formalin model, we measured the time (seconds) mice spent on licking or flinching the affected paw every 5 minutes for 45 or 60 minutes. For testing mechanical sensitivity, we confined mice in boxes placed on an elevated metal mesh floor and stimulated their hindpaws with a series of von Frey hairs with logarithmically increasing stiffness (0.02-2.56g, Stoelting), presented perpendicularly to the central plantar surface. We determined the 50% paw withdrawal threshold by Dixon's up-down method (45). Thermal sensitivity was assessed by tail immersion, hot plate, and radiant heat tests. For tail immersion test, the lower 5.0 cm portion of the tail was dipped in hot water at 48 or 52 $^{\circ}$ C and the tail flick latency was recorded with a cut-off time of 20 seconds.

For hot plate test, mice were placed on the hot plate at 50 or 56°C and the reaction time was scored when the animal began to exhibit signs of pain avoidance such as jumping or paw licking. Animals that did not respond to the noxious heat stimulus after 30 s were removed from the plate. For the radiant heat test, Hargreaves apparatus (IITC Life Science) was used and the basal paw withdrawal latency was adjusted to 9-12 s, with a cutoff of 20 s to prevent tissue damage.

Spontaneous and ongoing pain was assessed by conditioned place preference (CPP) test. We used a single trial conditioning protocol to measure CPP. All mice underwent a 3-day pre-conditioning habituation and animal behavior was video-recorded. Analyses of the pre-conditioning (baseline) behavior showed no pre-existing chamber preference. On the conditioning day, mice received the vehicle (PBS, 10 µl, i.t.) control paired with a randomly chosen chamber in the morning, and clonidine (10 µg in 10 µl PBS, i.t.) paired with the other chamber 4 h later. Chamber pairings were counterbalanced. On the test day (3 w after CCI), 20 h following the afternoon pairing, mice were placed in the CPP box with access to both chambers and the behavior was recorded for 15 min and analyzed by ANY-maze software for chamber preference.

In situ hybridization (ISH), Immunofluorescence, and Imaging

After appropriate survival times after nerve injury, mice were deeply anesthetized with isoflurane and perfused through the ascending aorta with PBS and then followed by 4% paraformaldehyde. The L4–L5 spinal cord segments and L4-L5 DRGs were removed and postfixed in the same fixative overnight. Spinal cord (30 µm, free floating) and DRG tissue sections (14 µm) were cut in a cryostat. Accordance with the manufacturer's instructions of RNA-scope system (Advanced Cell Diagnostics), in situ hybridization was performed by the Multiplex Fluorescent Kit v.2. We used probes directed against mouse *Sparcl1* (catalogue 424641, NM_010097.4) and mouse *Slc32a1* (catalogue 319191-C3, NM_009508.2) on the spinal cord sections from VGLUT2: Ai9 mice. For immunohistochemistry, the sections were blocked with 2% goat or donkey serum for 1 h at room temperature and then incubated overnight at 4°C with the following primary antibodies: anti-hevin (rat anti-hevin monoclonal 12:155, 1 µg/mL, from Eroglu's lab), anti-NeuN (mouse, 1:1000, Millipore, Catalog: MAB377), anti-GFAP (mouse, 1:1000, Millipore, catalog #MAB360), anti-IBA-1 (rabbit, 1:1000, Wako Chemicals, catalog #019-19741), and anti-Myc (mouse, 1:1000, Cell Signaling, Catalog: #2276). Tissue sections were then stained by cyanine 3(Cy3)-, cyanine 5(Cy5)- and/or FITC-conjugated secondary antibodies (1:400; Jackson ImmunoResearch) for 2 h at room

temperature. The stained sections were examined with a Nikon fluorescence microscope, and images were captured with a CCD Spot camera. For high resolution images, sections were also examined under a Zeiss 510 inverted confocal microscope. Quantification of RNAscope images were conducted in 5 spinal cord sections from two mice.

Patch clamp recordings in spinal cord slices

As previously reported (19), we removed a portion of the lumbar spinal cord (L4-L5) from young mice (4-6 weeks old) under urethane anesthesia (1.5 –2.0 g/kg, i.p.) and kept the spinal cord segments in pre-oxygenated ice-cold artificial cerebrospinal fluid (ACSF). Transverse slices (400 μm) were made on a vibrating microslicer and the slices were transferred to ACSF saturated with 95% O_2 and 5% CO_2 at 26°C for at least 1 h prior to experiment. The ACSF contains (in mM): NaCl 125, KCl 2.5, CaCl_2 2, MgCl_2 1, NaH_2PO_4 1.25, NaHCO_3 26, and glucose 25. The whole cell patch-clamp recordings were conducted from lamina IIo neurons in voltage clamp mode. Lamina IIo interneurons are predominately excitatory, make direct connections with lamina I projection neurons, and play critical role in transmitting mechanical pain (32, 33). Patch pipettes were prepared from thin-walled, borosilicate, glass-capillary tubing (1.5 mm o.d., World Precision Instruments) and filled with an internal solution containing following (in mM): K-gluconate 135, KCl 5, CaCl_2 0.5, MgCl_2 2, EGTA 5, HEPES 5, and ATP- Na_2 5. Data were acquired using an Axopatch 700B amplifier and were low pass-filtered at 2 kHz and digitized at 5 kHz. NMDA currents were recorded in IIo neurons by perfusing spinal cord slices with 100 μM NMDA for 3 sec in ACSF with TTX (5 μM). To measure NMDAR-mediated evoked EPSCs (NMDA-eEPSCs) in lamina IIo neurons, dorsal root (~ 5 mm) was stimulated through a suction electrode. NMDA-eEPSCs were pharmacologically isolated in Mg^{2+} -free ACSF containing CNQX (10 μM), strychnine (2 μM) and picrotoxin (100 μM). Neurons were voltage clamped at -40 mV and NMDA-eEPSCs were evoked at 0.05 Hz. QX-314 (5 mM) was added to the pipette solution to prevent discharge of action potentials. Synaptic strength was quantified by the peak amplitudes of eEPSCs. Data were collected with pClamp version 10 software and analyzed using Clampfit version 10.

Western blot

SDH protein samples were prepared as we previously demonstrated (46), and 30 μg of proteins were loaded for each lane and separated on SDS-PAGE gel (4–15%; Bio-Rad). After the transfer,

the blots were incubated overnight at 4°C with anti-hevin (rat anti-hevin monoclonal 12:155, 1 µg/mL). These blots were further incubated with HRP-conjugated secondary antibody, developed in ECL solution (Pierce), and the chemiluminescence was revealed by Bio-Rad ChemiDoc XRS for 1–5 min.

Human CSF collection

Human CSF was collected from the patients enrolled in the Markers of Alzheimer’s disease after Propofol vs. Isoflurane Anesthesia (MAD-PIA) trial. MAD-PIA was a prospective randomized trial registered with <http://www.clinicaltrials.gov> (NCT01640275) on June 20, 2012, by Miles Berger (35). Human CSF samples (10 ml) were obtained from the lumbar drain (AccuDrain INS-8400; Integra Neurosciences, Plainsboro, NJ, USA) at the time of drain placement (0h) and 10-12h after intracranial surgery.

ELISA

Mouse hevin ELISA kit was purchased from LifeSpan BioSciences (Catalog: LS-F12654). Human hevin ELISA kit was from Cloud-Clone Corp (Catalog: SEM267Hu). ELISA was conducted in mouse CSF samples (10 µl) and human CSF (10 µl), according to manufacturer’s instructions. The standard curve was included in each experiment.

Statistical analyses

All the data were expressed as mean ± s.e.m, as indicated in the figure legends. Statistical analyses were completed with Prism GraphPad 8.0. Biochemical and behavioral data were analyzed using two-tailed student’s t-test (two groups, unpaired or paired) or Two-Way ANOVA followed by post-hoc Bonferroni test. Electrophysiological data were tested using one-way ANOVA (for multiple comparisons) or Two-Way ANOVA (for multiple time points) followed by post-hoc Bonferroni test or student’s t-test (two groups, unpaired or paired). The criterion for statistical significance was $P < 0.05$.

Study approval

All the animal procedures were conducted in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care

& Use Committee (IACUC) of Duke University. The human CSF study was approved by the Duke University Institutional Review Board.

Author contributions

G.C., C.E. and R.R.J designed research; G.C., J.X., H.L., and X.L. performed experiments and data analysis; S.K.S and J.R. prepared critical reagents and participated in project discussion; M.L.J., J.P.M. and M.B. provided human CSF samples and contributed to the related experiments. R-.R.J. and C.E. supervised the project; G.C. and R-.R.J. wrote the manuscript, and other authors edited the paper.

Acknowledgements

This work was supported by Duke University Anesthesiology Research Fund and partially supported by NIH grants R01 grant DE17794 and DoD grants W81XWH2110756 and W81XWH2110885. MB acknowledges additional support from NIH grants K76-AG057022, P30-AG028716, and P30-AG072958. JPM was supported by NIH grants R01 HL130443 and AG074185. CE and JJR were supported by NIH grants R01 DA031833 and NS102237. CE is a Howard Hughes Medical Institute Investigator.

References

1. Costigan M, Scholz J, and Woolf CJ. Neuropathic pain: a maladaptive response of the nervous system to damage. *AnnuRevNeurosci*. 2009;32:1-32.
2. Baron R, Binder A, and Wasner G. Neuropathic pain: diagnosis, pathophysiological mechanisms, and treatment. *Lancet Neurol*. 2010;9(8):807-19.
3. Woolf CJ, and Salter MW. Neuronal plasticity: increasing the gain in pain. *Science*. 2000;288(5472):1765-9.
4. Liu XJ, Gingrich JR, Vargas-Caballero M, Dong YN, Sengar A, Beggs S, et al. Treatment of inflammatory and neuropathic pain by uncoupling Src from the NMDA receptor complex. *NatMed*. 2008;14(12):1325-32.
5. Chen G, Xie RG, Gao YJ, Xu ZZ, Zhao LX, Bang S, et al. beta-arrestin-2 regulates NMDA receptor function in spinal lamina II neurons and duration of persistent pain. *Nat Commun*. 2016;7:12531.
6. Ji RR, Donnelly CR, and Nedergaard M. Astrocytes in chronic pain and itch. *Nat Rev Neurosci*. 2019.
7. Foley JC, McIver SR, and Haydon PG. Gliotransmission modulates baseline mechanical nociception. *Mol Pain*. 2011;7:93.
8. Xu Q, Ford NC, He S, Huang Q, Anderson M, Chen Z, et al. Astrocytes contribute to pain gating in the spinal cord. *Sci Adv*. 2021;7(45):eabi6287.
9. Singh SK, Stogsdill JA, Pulimood NS, Dingsdale H, Kim YH, Pilaz LJ, et al. Astrocytes Assemble Thalamocortical Synapses by Bridging NRX1alpha and NL1 via Hevin. *Cell*. 2016;164(1-2):183-96.
10. Kucukdereli H, Allen NJ, Lee AT, Feng A, Ozlu MI, Conatser LM, et al. Control of excitatory CNS synaptogenesis by astrocyte-secreted proteins Hevin and SPARC. *Proc Natl Acad Sci U S A*. 2011;108(32):E440-9.
11. Stogsdill JA, Ramirez J, Liu D, Kim YH, Baldwin KT, Enustun E, et al. Astrocytic neuroligins control astrocyte morphogenesis and synaptogenesis. *Nature*. 2017;551(7679):192-7.
12. Araque A, Parpura V, Sanzgiri RP, and Haydon PG. Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci*. 1999;22(5):208-15.
13. Eroglu C, Allen NJ, Susman MW, O'Rourke NA, Park CY, Ozkan E, et al. Gabapentin receptor alpha2delta-1 is a neuronal thrombospondin receptor responsible for excitatory CNS synaptogenesis. *Cell*. 2009;139(2):380-92.
14. Baldwin KT, Tan CX, Strader ST, Jiang C, Savage JT, Elorza-Vidal X, et al. HepaCAM controls astrocyte self-organization and coupling. *Neuron*. 2021;109(15):2427-42 e10.
15. Ji RR, Berta T, and Nedergaard M. Glia and pain: Is chronic pain a gliopathy? *Pain*. 2013.
16. Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, et al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature*. 2017;541(7638):481-7.
17. Guo W, Wang H, Watanabe M, Shimizu K, Zou S, LaGraize SC, et al. Glial-cytokine-neuronal interactions underlying the mechanisms of persistent pain. *JNeurosci*. 2007;27(22):6006-18.
18. Tsuda M, Kohro Y, Yano T, Tsujikawa T, Kitano J, Tozaki-Saitoh H, et al. JAK-STAT3 pathway regulates spinal astrocyte proliferation and neuropathic pain maintenance in rats. *Brain*. 2011.

19. Chen G, Park CK, Xie RG, Berta T, Nedergaard M, and Ji RR. Connexin-43 induces chemokine release from spinal cord astrocytes to maintain late-phase neuropathic pain in mice. *Brain*. 2014;137(Pt 8):2193-209.
20. Chiang CY, Sessle BJ, and Dostrovsky JO. Role of Astrocytes in Pain. *NeurochemRes*. 2012.
21. Liu X, Bae C, Gelman BB, Chung JM, and Tang SJ. A neuron-to-astrocyte Wnt5a signal governs astrogliosis during HIV-associated pain pathogenesis. *Brain*. 2022.
22. Zhang H, Yoon SY, Zhang H, and Dougherty PM. Evidence that spinal astrocytes but not microglia contribute to the pathogenesis of Paclitaxel-induced painful neuropathy. *JPain*. 2012;13(3):293-303.
23. Zhuang ZY, Wen YR, Zhang DR, Borsello T, Bonny C, Strichartz GR, et al. A peptide c-Jun N-terminal kinase (JNK) inhibitor blocks mechanical allodynia after spinal nerve ligation: respective roles of JNK activation in primary sensory neurons and spinal astrocytes for neuropathic pain development and maintenance. *JNeurosci*. 2006;26(13):3551-60.
24. Kim DS, Li KW, Boroujerdi A, Peter YY, Zhou CY, Deng P, et al. Thrombospondin-4 Contributes to Spinal Sensitization and Neuropathic Pain States. *J Neurosci*. 2012;32(26):8977-87.
25. Park J, Yu YP, Zhou CY, Li KW, Wang D, Chang E, et al. Central Mechanisms Mediating Thrombospondin-4-induced Pain States. *J Biol Chem*. 2016;291(25):13335-48.
26. Kohro Y, Matsuda T, Yoshihara K, Kohno K, Koga K, Katsuragi R, et al. Spinal astrocytes in superficial laminae gate brainstem descending control of mechanosensory hypersensitivity. *Nat Neurosci*. 2020;23(11):1376-87.
27. Girard JP, and Springer TA. High endothelial venules (HEVs): specialized endothelium for lymphocyte migration. *Immunol Today*. 1995;16(9):449-57.
28. King T, Vera-Portocarrero L, Gutierrez T, Vanderah TW, Dussor G, Lai J, et al. Unmasking the tonic-aversive state in neuropathic pain. *Nat Neurosci*. 2009;12(11):1364-6.
29. Sorge RE, Mapplebeck JC, Rosen S, Beggs S, Taves S, Alexander JK, et al. Different immune cells mediate mechanical pain hypersensitivity in male and female mice. *NatNeurosci*. 2015;18(8):1081-3.
30. Mapplebeck JCS, Dalgarno R, Tu Y, Moriarty O, Beggs S, Kwok CHT, et al. Microglial P2X4R-evoked pain hypersensitivity is sexually dimorphic in rats. *Pain*. 2018;159(9):1752-63.
31. Kuhn JA, Vainchtein ID, Braz J, Hamel K, Bernstein M, Craik V, et al. Regulatory T-cells inhibit microglia-induced pain hypersensitivity in female mice. *Elife*. 2021;10.
32. Braz J, Solorzano C, Wang X, and Basbaum AI. Transmitting Pain and Itch Messages: A Contemporary View of the Spinal Cord Circuits that Generate Gate Control. *Neuron*. 2014;82(3):522-36.
33. Duan B, Cheng L, and Ma Q. Spinal Circuits Transmitting Mechanical Pain and Itch. *Neurosci Bull*. 2018;34(1):186-93.
34. Chen G, Park CK, Xie RG, and Ji RR. Intrathecal bone marrow stromal cells inhibit neuropathic pain via TGF-beta secretion. *J ClinInvest*. 2015;125(8):3226-40.
35. Berger M, Nadler JW, Friedman A, McDonagh DL, Bennett ER, Cooter M, et al. The Effect of Propofol Versus Isoflurane Anesthesia on Human Cerebrospinal Fluid Markers

- of Alzheimer's Disease: Results of a Randomized Trial. *J Alzheimers Dis.* 2016;52(4):1299-310.
36. Kim SK, Hayashi H, Ishikawa T, Shibata K, Shigetomi E, Shinozaki Y, et al. Cortical astrocytes rewire somatosensory cortical circuits for peripheral neuropathic pain. *J Clin Invest.* 2016;126(5):1983-97.
 37. Lively S, Moxon-Emre I, and Schlichter LC. SC1/hevin and reactive gliosis after transient ischemic stroke in young and aged rats. *J Neuropathol Exp Neurol.* 2011;70(10):913-29.
 38. Wang Z, Tao Y, Song C, Liu P, Wang C, Li Y, et al. Spinal hevin mediates membrane trafficking of GluA1-containing AMPA receptors in remifentanyl-induced postoperative hyperalgesia in mice. *Neurosci Lett.* 2020;722:134855.
 39. Aarts M, Liu Y, Liu L, Besshoh S, Arundine M, Gurd JW, et al. Treatment of ischemic brain damage by perturbing NMDA receptor- PSD-95 protein interactions. *Science.* 2002;298(5594):846-50.
 40. Berger M, Oyeyemi D, Olurinde MO, Whitson HE, Weinhold KJ, Woldorff MG, et al. The INTUIT Study: Investigating Neuroinflammation Underlying Postoperative Cognitive Dysfunction. *J Am Geriatr Soc.* 2019;67(4):794-8.
 41. Salter MW, and Stevens B. Microglia emerge as central players in brain disease. *Nat Med.* 2017;23(9):1018-27.
 42. Chen G, Zhang YQ, Qadri YJ, Serhan CN, and Ji RR. Microglia in Pain: Detrimental and Protective Roles in Pathogenesis and Resolution of Pain. *Neuron.* 2018;100(6):1292-311.
 43. Pogatzki EM, and Raja SN. A mouse model of incisional pain. *Anesthesiology.* 2003;99(4):1023-7.
 44. Kohro Y, Sakaguchi E, Tashima R, Tozaki-Saitoh H, Okano H, Inoue K, et al. A new minimally-invasive method for microinjection into the mouse spinal dorsal horn. *Sci Rep.* 2015;5:14306.
 45. Dixon WJ. Efficient analysis of experimental observations. *Annu Rev Pharmacol Toxicol.* 1980;20:441-62.
 46. Berta T, Park CK, Xu ZZ, Xie RG, Liu T, Lu N, et al. Extracellular caspase-6 drives murine inflammatory pain via microglial TNF-alpha secretion. *J Clin Invest.* 2014;124(3):1173-86.

Figure legends




Figure 1. Baseline pain, inflammatory pain, and neuropathic pain in WT and hevin-KO mice.

(A-D) There are no significant differences in mechanical and thermal pain threshold between WT and hevin-KO male mice, as shown in von Frey test (A), radiant heat test (B), hot plate test (C) and tail immersion test (D). $P > 0.05$, unpaired t-test, $n = 11$ mice/group. (E) Formalin-induced acute inflammatory pain was significantly reduced in hevin-KO male mice. Left, time-course of licking and flinching behavior following intraplantar injection of 5% formalin. Right, formalin-induced Phase I (1–10 min) and Phase II (10–45 min) responses. $*P < 0.05$, Two-way ANOVA followed by Bonferroni posthoc test. $n = 6$ mice/group. (F) Mechanical allodynia, induced by

intraplantar injection of carrageenan, recovered faster in hevin-KO male mice than in WT male mice. Arrow indicates the time of carrageenan injection. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, compared to BL group, $^{\#}P < 0.05$, Two-way ANOVA followed by Bonferroni posthoc test. $n = 5$ mice/group. Data shown as mean \pm SEM. (G) Hevin-KO male mice recovered faster from CCI-induced persistent mechanical allodynia than WT male mice. $***P < 0.001$, compared to BL group; $^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.001$, Two-way ANOVA followed by Bonferroni posthoc test. $n = 11$ mice/group. (H) Ongoing pain 3 weeks after CCI in WT and hevin-KO male mice were tested using a 2-chamber conditional place preference. Ongoing pain was present in WT mice but absent in hevin-KO mice following clonidine treatment (10 μ g, i.t.). $*P < 0.05$, unpaired t-test, $n = 6$ mice/group. All data are shown as mean \pm SEM.




Figure 2. Intrathecal administration of hevin but not hevin- Δ DE (10 μ g, i.t.) decreases paw withdrawal threshold in naïve mice and mice with nerve injury and enhances spontaneous pain by formalin.

(A) Intrathecal injection of WT hevin but not mutant hevin (hevin- Δ DE) induced mechanical allodynia in naïve mice, lasting > 3 days. $n = 5$ mice/group. $**P < 0.01$, $***P < 0.001$, compared to BL group, $\#P < 0.05$, $\#\#P < 0.01$, $\#\#\#P < 0.001$, Two-way ANOVA followed by Bonferroni posthoc test. Arrow indicates hevin injection on day 0. (B) Repeated intrathecal injections of WT hevin but not mutant hevin (hevin- Δ DE), given at 12 and 13 days after CCI induced, further exacerbated mechanical allodynia. $n = 5$ mice/group. $*P < 0.05$, $**P < 0.01$, compared to CCI-12d baseline group; $\#P < 0.05$, $\#\#P < 0.01$, $\#\#\#P < 0.001$, Two-way ANOVA followed by Bonferroni posthoc test. Arrows indicate the time of hevin injections. (C-E) Intrathecal injection of WT hevin but not mutant hevin (hevin- Δ DE) enhances formalin-induced licking and flinching on phase-2 (10-45 min) and phase-2+ (45-60 min) but not phase-1 (0-10 min) in both male (C) and female (D) mice. $*P < 0.05$, $***P < 0.001$, compared to hevin- Δ DE group, Two-way ANOVA followed by Bonferroni posthoc test. All data are shown as mean \pm SEM.




Figure 3. RNAscope images showing *Sparcl1* expression in excitatory and inhibitory neurons in the spinal cord dorsal horn of VGLUT2: Ai9 mice.

(A) RNA-scope images show co-localization of *Sparcl1* (red) with VGLUT2+ excitatory neurons (green) and *Slc32a1*+ inhibitory neurons (blue). Top, merged low-magnification image. Scale bars, 50 μ m. Bottom, merged and single channel images enlarged from the box in the top panel. Filled arrows show *Sparcl1*+/VGLUT2+ excitatory neurons, open arrows show *Sparcl1*+/ *Slc32a1*+ inhibitory neurons, and arrows with cross show only *Sparcl1*+ cells. Scale bar, 20 μ m. (B) Quantification of the percentage of *Sparcl1*+ cells expressing VGLUT2 or *Slc32a1* and the percentage of VGLUT2+ or *Slc32a1*+ cells expressing *Sparcl1* in the spinal dorsal horn. n = 6 spinal cord sections from three mice. Data are shown as mean \pm SEM.




Figure 4. Expression of hevin in spinal astrocytes by intraspinal hevin-AAV reinstates neuropathic pain in hevin KO mice.

(A) Double immunostaining of hevin (green) and GFAP (red) in SDH. Note hevin is primarily colocalized with GFAP. Scales: 100 μ m (left) and 20 μ m (right). The box is enlarged in the right panels. (B) Absence of hevin immunostaining in SDH in hevin KO mice. Scale, 100 μ m. (C)

Paradigm for measuring mechanical allodynia in hevin KO mice with intra-spinal microinjection of hevin-AAV and hevin Δ DE-AAV, given 6d before CCI. (D) SDH microinjection of AAV induced reduction in paw withdrawal threshold in naïve hevin KO mice. After CCI, mechanical allodynia was significantly prolonged in hevin-AAV treated mice than hevin Δ DE-AAV treated mice. $n = 6$ mice/group. $*P < 0.05$, Two-way ANOVA followed by Bonferroni posthoc test. Green and red arrows indicate the time of virus injection and nerve injury, respectively. (E) Paradigm for measuring mechanical allodynia in hevin KO mice with intra-spinal microinjection of hevin-AAV and hevin- Δ DE-AAV, given 2 days after CCI. (F) SDH microinjection of hevin-AAV, given after nerve injury, significantly enhanced and prolonged mechanical allodynia in hevin KO mice vs. hevin Δ DE-AAV treated mice. $n = 6$ mice/group. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, Two-way ANOVA followed by Bonferroni posthoc test. Arrows indicate the time of virus injection and nerve injury. (G) Triple immunostaining of Myc (red), hevin (green) and GFAP (blue) in SDH in hevin KO mice, 24 days after the ipsilateral SDH hevin-AAV injection. Note hevin expression is absent in the contralateral SDH of hevin-KO mice. Scale, 100 μ m. (H) Enlarged images in the box of F panel G, with additional merged images for Myc/hevin, Myc/GFAP, and hevin/GFAP. Note hevin $^{+}$ astrocytes also Myc $^{+}$ /GFAP $^{+}$ in superficial SDH. Scale, 20 μ m. All data are shown as mean \pm SEM.




Figure 5. Hevin regulates NMDA-evoked pain and enhances NMDA currents in SDH neurons.

(A) The duration of mechanical allodynia, induced by intrathecal injection of NMDA (3 nmol), is significantly shorter in hevin KO mice than in wild-type mice. $n = 6$ mice/group. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, compared to BL group; $\#P < 0.05$, $\#\#P < 0.01$, $\#\#\#P < 0.001$, Two-way ANOVA followed by Bonferroni posthoc test. Arrow indicates the time of NMDA injection. (B, C) Left: representative traces of inward currents in WT and hevin KO mice, induced by NMDA (100 μ M, 3 sec) via bath application. Note smaller NMDA currents in hevin KO mice and different effects of Ro25-6091 (GluN2B antagonist) and TCN-201 (GluN2A antagonist). Right: amplitude of NMDA-induced currents. $n = 9, 10$ neurons per group (shown in each column). $*P < 0.05$, One-Way ANOVA followed by Bonferroni posthoc test. n.s., not significant. (D, E) Left: representative traces of inward currents in ACSF and hevin (100 ng/ml, 4 min) treated spinal cord slices, induced by NMDA (100 μ M, 3 sec) via bath application. Note different effects of Ro25-6091 and TCN-

201. Right: amplitude of NMDA-induced currents. $n = 7$ neurons per group. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, One-Way ANOVA followed by Bonferroni posthoc test. n.s., not significant. Data are shown as mean \pm SEM.




Figure 6. Anti-hevin monoclonal antibody 12:54 reduces inflammatory, postoperative, and neuropathic pain in WT mice. (A) ELISA analysis showing increased hevin level in the CSF 14 days after CCI. n = 5 mice/group. * $P < 0.05$, unpaired t-test. (B) Left, time course of formalin-induced pain in WT male mice treated with intrathecal anti-hevin 12:155 monoclonal antibody (control antibody, 10 μ g) or anti-hevin 12:54 monoclonal antibody (function blocking antibody, 10 μ g). n = 5 mice per group. Right, formalin-induced Phase-1 and Phase-2 responses. * $P < 0.05$, unpaired t-test. (C) Intrathecal injection of anti-hevin 12:54 antibody (10 μ g), given 3 days after

CFA injection, reduced CFA-induced mechanical allodynia for 5h. Arrows indicate the time of antibody injection. n = 5 mice/group. [#]*P*<0.05, [#]*P*<0.01, versus corresponding BL group; **P*<0.05, versus anti-hevin 12:155 group, Two-Way ANOVA followed by Bonferroni posthoc test. (D) Intrathecal injection of anti-hevin 12:54 antibody (10 μg), given 3 hours after plantar incision, reduced incision-induced mechanical allodynia for 5 h in male and female mice. Arrows indicate the time of antibody injection. n = 10 mice/group. ^{###}*P*<0.001, versus corresponding BL group; ^{***}*P*<0.001, versus anti-hevin 12:155 group, Two-Way ANOVA followed by Bonferroni posthoc test. (E, F) Intrathecal injection of anti-hevin 12:54 antibody (10 μg), given 7 days (E) and 21 days (F) after nerve injury, reduced CCI-induced mechanical allodynia for 5h. Arrows indicate the time of antibody injection. n = 5–6 mice/group. [#]*P*<0.05, versus corresponding BL group; [§]*P*<0.05, ^{§ §}*P*<0.01, versus corresponding baseline at CCI 7d or CCI 21d; **P*<0.05, ***P*<0.01, Two-Way ANOVA followed by Bonferroni posthoc test. Data are shown as mean ± SEM.






Figure 7. Intracranial surgery increases hevin levels in human CSF samples.

(A, B) ELISA analysis showing increased hevin levels in human CSF 12 hours after intracranial surgery. The same data were presented unpaired (A) and paired (B). (C) BCA protein assay showing total protein level increased in human CSF 12 hours after surgery. (D) Fold changes of normalized hevin level in human CSF 12 hours after surgery, normalized to total protein changes. $n = 10$ patients. $**P < 0.01$, $***P < 0.001$, paired t-test, Data shown as mean \pm SEM.




Supplemental Figure 1. Schematic of hevin protein (A), hevin-AAV (B), and mapped epitopes (c) for anti-hevin monoclonal antibodies.




Supplemental Figure 2. Hevin and NR2B activation drive mechanical pain in both sexes.

(A) Intrathecal injection of purified WT hevin (5 μ g, i.t.) induced robust mechanical allodynia in both WT naïve male and female mice for more than 1 day. n = 5 mice per group. *** P <0.001, Two-Way ANOVA followed by Bonferroni posthoc test. Data shown as mean \pm SEM. (B) Intrathecal hevin-induced mechanical allodynia was completely reversed by Ro25-6981 (GluN2B antagonist) in both WT male and female mice. n = 5 mice per group. *** P <0.001. Two-Way ANOVA followed by Bonferroni posthoc test. Data shown as mean \pm SEM.



Supplemental Figure 3. Hevin expression in SDH and DRG of WT and KO mice.

(A, B) Double immunostaining of hevin (green, A) and the neuronal marker NeuN (red, B) in the spinal cord dorsal horn in mice. (C, D) Double immunostaining of hevin (green, C) and microglial marker Iba1 (red, D) in the spinal cord dorsal horn in mice. Note hevin is expressed by some neurons (NeuN⁺) but not microglia (Iba1⁺). Right images are enlarged images of the left image. Scale bars: 50 μ m (left) and 10 μ m (right). (E) Western blot showing hevin expression in spinal dorsal horn of WT mice and KO mice. Right, uncut full gel of the left blot. (F) Hevin expression in DRG neurons and satellite glial cells (indicated by arrows) of WT mice. Right, enlarged box of left. Scale, 20 μ m. (G) Hevin expression in DRG is lost in hevin KO mice.



Supplemental Figure 4. Electrophysiological characterization of hevin's actions in lamina IIo neurons of spinal cord slices.

(A, B) Acute effects of hevin (100 ng/mL, 5 min) on NMDA or AMPA-induced current in lamina II neurons. Note hevin enhanced the NMDA-induced currents but no effects on AMPA-evoked currents in lamina IIo neurons. $n = 7$ or 5 neurons per group (shown in each column). $**P < 0.01$, paired t-test, n.s., not significant. Data shown as mean \pm SEM. (C) Effects of hevin (100 ng/mL) on NMDA-eEPSC in lamina IIo neurons evoked by DR stimulation. Note hevin also enhanced the NMDA receptor-mediated eEPSCs. $n = 4$ neurons per group. $**P < 0.01$, paired t-test. Data shown as mean \pm SEM. (D) Hevin superfusion (100 ng/ml) increased spontaneous EPSC frequency and amplitude in lamina IIo neurons of spinal cord slices from CCI mice. $n = 6$ neurons per group. $*P < 0.05$, paired t-test. Data shown as mean \pm SEM.